



Royal College  
of Midwives



Learn, Share & Improve Care  
Inspiring midwives, improving maternity care

E: [midirs.helpdesk@rcm.org.uk](mailto:midirs.helpdesk@rcm.org.uk)

T: 07768 015063

W: [www.midirs.org](http://www.midirs.org)

## Search Pack P200 (2022) Coronavirus (COVID-19) in pregnancy (2022)

Updated 17 June 2022

Records on coronavirus (COVID-19) in pregnancy from 2022 only. For earlier records on this topic see P200 (2020) and P200 (2021). Includes choice and accessibility of maternal health services. Does not include records on COVID-19 vaccination in pregnancy (P201); COVID-19 in the neonate or infant feeding during the pandemic (PN193); the impact of COVID-19 on midwives (M95); COVID-19 in labour, birth and the impact on intrapartum care (L69) or the impact of COVID-19 on postnatal health and care (PN194).

### Search Pack Tips:

1. The results of your Search Pack start below. If the full text of an article is freely available online, we will provide the URL - simply click the live hyperlink to view. MIDIRS cannot take responsibility for external content, and hyperlinks provided by external organisations are subject to change.
2. You can search within the Search Pack PDF for relevant keywords by pressing 'CTRL + F' and entering the terms you wish to find.
3. If a record in the Bespoke Search has been published in MIDIRS Midwifery Digest or Essentially MIDIRS then this can be ordered via the following [link](#)
4. If a record in the Search Pack does not have a full-text URL link and is not from a MIDIRS publication, then unfortunately we are unable to supply. Try your university library, Trust library, or British library for a copy.

MIDIRS is part of RCM Information Services Limited which is a company incorporated in England and Wales under company no.11914882 with registered office at 10-18 Union Street, London SE1 1SZ  
RCM Information Services Limited is a subsidiary of The Royal College of Midwives



The Royal College of Midwives Trust  
10-18 Union Street  
London  
SE1 1SZ



Open 24 hours a day, 7 days a week  
T: 0300 303 0444  
F: +44 20 7312 3536  
E: [info@rcm.org.uk](mailto:info@rcm.org.uk)  
W: [www.rcm.org.uk](http://www.rcm.org.uk)



**Chief Executive:**  
Gill Walton,  
MA, PGDip, BSc Hons, RM  
**President:**  
Rebecca Davies, RM  
**Patron:**  
HRH The Princess Royal

## 2022-05692

**Point-of-care ultrasound for diagnosis of pneumothorax in a pregnant COVID-19 patient in the emergency department.** Gulen M, Satar S, Unlu N, et al (2022), Journal of Obstetrics and Gynaecology 13 June 2022, online

Case report of a pregnant woman admitted to the emergency department with complaints of dyspnoea, chest pain and a cough at 16 weeks' gestation. The patient was subsequently diagnosed with pneumothorax and COVID-19 pneumonia. (LDO)

---

## 2022-05652

**Experiences of women who gave birth during the pandemic.** Hancock D (2022), Journal of Health Visiting vol 10, no 5, May 2022, pp 195-198

The Care Quality Commission surveyed women who had gone through pregnancy and birth during the national lockdown in 2021. Dave Hancock looks at some positive results and key areas for improvement in maternity care. (Author)

---

## 2022-05594

**The outcomes of favipiravir exposure in pregnancy: a case series.** Ertem O, Guner O, Incir C, et al (2022), Archives of Gynecology and Obstetrics 27 May 2022, online

### Purpose

As in vitro and in vivo studies reported antiviral efficacy against RNA viruses, favipiravir, a pyrazinocarboxamide derivative, has become one of the treatment options for COVID-19 in some countries including Turkey. Preclinical studies demonstrated the risk for teratogenicity and embryotoxicity. Hence, the drug is contraindicated during pregnancy. Although limited in numbers, case-based evaluations indicate that favipiravir might not be a major teratogen in human pregnancies. This study aimed to present and analyze the outcomes of favipiravir exposure during pregnancy.

### Methods

In this case series, the outcomes of nine pregnancies that were referred to the Teratology Information Service of Dokuz Eylul University Faculty of Medicine, Department of Medical Pharmacology between 01 April 2020 and 30 November 2021 were retrospectively evaluated.

### Results

One spontaneous abortion, two elective terminations, one preterm live delivery and five term live deliveries were detected. The premature newborn was reported dead on the 5th day of neonatal intensive care unit admission. Physiological jaundice and transient respiratory distress were recorded in two term infants. One term infant was antenatally diagnosed with renal pelviectasis, but the findings resolved postnatally without requiring intervention.

### Conclusion

The data indicate that favipiravir is not likely to be a major teratogen. Yet, it is not possible to draw a definite conclusion due to methodological limitations. Favipiravir exposures during pregnancy should be followed up closely and the outcomes should be reported consistently. (Author)

**Full URL:** <https://doi.org/10.1007/s00404-022-06615-z>

---

MIDIRS is part of RCM Information Services Limited which is a company incorporated in England and Wales under company no.11914882 with registered office at 10-18 Union Street, London SE1 1SZ  
RCM Information Services Limited is a subsidiary of The Royal College of Midwives



**The Royal College of Midwives Trust**  
10-18 Union Street  
London  
SE1 1SZ



**Open 24 hours a day, 7 days a week**  
T: 0300 303 0444  
F: +44 20 7312 3536  
E: [info@rcm.org.uk](mailto:info@rcm.org.uk)  
W: [www.rcm.org.uk](http://www.rcm.org.uk)



**Chief Executive:**  
Gill Walton,  
MA, PGDip, BSc Hons, RM  
**President:**  
Rebecca Davies, RM  
**Patron:**  
HRH The Princess Royal

2022-05554

**“Other risks don't stop”: adapting a youth sexual and reproductive health intervention in Zimbabwe during COVID-19.**

Mackworth-Young CRS, Mavodza C, Nyamwanza R, et al (2022), Sexual & Reproductive Healthcare vol 30, no 1, 2022, 2029338

COVID-19 threatens hard-won gains in sexual and reproductive health (SRH) through compromising the ability of services to meet needs. Youth are particularly threatened due to existing barriers to their access to services. CHIEDZA is a community-based integrated SRH intervention for youth being trialled in Zimbabwe. CHIEDZA closed in March 2020, in response to national lockdown, and reopened in May 2020, categorised as an essential service. We aimed to understand the impact of CHIEDZA's closure and its reopening, with adaptations to reduce COVID-19 transmission, on provider and youth experiences. Qualitative methods included interviews with service providers (n = 22) and youth (n = 26), and observations of CHIEDZA sites (n = 10) and intervention team meetings (n = 7). Analysis was iterative and inductive. The sudden closure of CHIEDZA impeded youth access to SRH services. The reopening of CHIEDZA was welcomed, but the necessary adaptations impacted the intervention and engagement with it. Adaptations restricted time with healthcare providers, heightening the tension between numbers of youths accessing the service and quality of service provision. The removal of social activities, which had particularly appealed to young men, impacted youth engagement and access to services, particularly for males. This paper demonstrates how a community-based youth-centred SRH intervention has been affected by and adapted to COVID-19. We demonstrate how critical ongoing service provision is, but how adaptations negatively impact service provision and youth engagement. The impact of adaptations additionally emphasises how time with non-judgemental providers, social activities, and integrated services are core components of youth-friendly services, not added extras. (Author)

Full URL: <https://doi.org/10.1080/26410397.2022.2029338>

---

2022-05546

**The Convergence of COVID-19 and Systemic Racism: An Evaluation of Current Evidence, Health System Changes, and Solutions Grounded in Reproductive Justice.**

Britt AJ, Carlson NS, Joseph NT, et al (2021), Journal of Midwifery & Women's Health vol 66, no 3, May/June 2021, pp 298-303

The objectives of this commentary are threefold. First, we will review the newest evidence about the intersection of systemic racism, COVID-19, and perinatal disparities. Second, we will explore how changes in health care delivery made in reaction to COVID-19 may exclude people of color who are pregnant and potentially exacerbate these disparities. Finally, we will discuss strategies by which midwives and the health care delivery system might be reimagined within a reproductive justice framework during this time to better meet both the immediate challenges of COVID-19 and the longer-term need to address entrenched systemic racism. (Author, edited)

Full URL: <https://doi.org/10.1111/jmwh.13250>

---

MIDIRS is part of RCM Information Services Limited which is a company incorporated in England and Wales under company no.11914882 with registered office at 10-18 Union Street, London SE1 1SZ  
RCM Information Services Limited is a subsidiary of The Royal College of Midwives



**The Royal College of Midwives Trust**  
10-18 Union Street  
London  
SE1 1SZ



**Open 24 hours a day, 7 days a week**  
T: 0300 303 0444  
F: +44 20 7312 3536  
E: [info@rcm.org.uk](mailto:info@rcm.org.uk)  
W: [www.rcm.org.uk](http://www.rcm.org.uk)



**Chief Executive:**  
Gill Walton,  
MA, PGDip, BSc Hons, RM  
**President:**  
Rebecca Davies, RM  
**Patron:**  
HRH The Princess Royal

2022-05290

**The role of laboratory parameters in predicting severity of COVID-19 disease in pregnant patients.** Sahin ON, Aktoz F, Bagci H, et al (2022), Journal of Obstetrics and Gynaecology 23 May 2022, online

We aimed to examine the relationship between laboratory markers and the severity of the disease in pregnant women diagnosed with coronavirus disease 2019 (COVID-19). Clinical records were retrospectively reviewed for 112 pregnant women. Patients diagnosed with COVID-19 were divided into two groups as mild/moderate and severe. The relationship between predicting the severity of the disease and laboratory parameters was investigated. Neutrophil lymphocyte ratio, C-reactive protein (CRP), ferritin and aspartate aminotransferase levels were significantly higher in severe COVID-19 cases than mild/moderate cases ( $p = .048$ ,  $p = .003$ ,  $p = .015$  and  $p = .035$ , respectively). CRP was found to be the most useful marker in terms of diagnostic performance with a cut off value of 10.8 (sensitivity 80%, specificity 56.1%, NPV 88.5% and PPV 40.0%). The best diagnostic performance was obtained using CRP and ferritin combined with cut-offs of 10.8 mg/L for CRP and 26.5  $\mu\text{g/L}$  for ferritin. Combined CRP and ferritin showed sensitivity, specificity, negative predictive value and positive predictive value of 94.7%, 52.8%, 96.6% and 41.9%, respectively, in predicting severe COVID-19. The combination of CRP and ferritin parameters may be useful in estimating the severity of the disease in pregnant patients who were initially diagnosed with COVID-19.

#### Impact Statement

What is already known about this subject? Coronavirus disease 2019 (COVID-19) can rapidly develop into acute respiratory distress syndrome (ARDS) and result in serious complications in some pregnant patients. Therefore, timely diagnosis of patients is crucial. Most previous reports of COVID-19 laboratory results are based on data from the general population and limited information is available regarding pregnancy status. Although laboratory medicine makes an important contribution to clinical decision making in many infectious diseases, including COVID-19, studies to predict the severity of the disease with laboratory markers are limited and the results are contradictory.

What do the results of this study add? Our study shows that C-reactive protein (CRP), neutrophil lymphocyte ratio (NLR), ferritin and aspartate aminotransferase (AST) are associated with severe disease in pregnant women diagnosed with COVID-19. In addition, the use of combined CRP and ferritin appears to have higher sensitivity and negative predictive value than using other tests alone. Furthermore, this study shows that coagulation markers are not useful in predicting disease severity in pregnancy.

What are the implications of these findings for clinical practice and/or further research? Predicting the severity of COVID-19 disease in pregnancy can prevent unnecessary hospitalisations and allow the implementation of the necessary clinical approach. Further studies can focus on the clinical usefulness of these parameters in predicting severe COVID-19 in pregnancy. (Author)

---

MIDIRS is part of RCM Information Services Limited which is a company incorporated in England and Wales under company no.11914882 with registered office at 10-18 Union Street, London SE1 1SZ  
RCM Information Services Limited is a subsidiary of The Royal College of Midwives



**The Royal College of Midwives Trust**  
10-18 Union Street  
London  
SE1 1SZ



**Open 24 hours a day, 7 days a week**  
T: 0300 303 0444  
F: +44 20 7312 3536  
E: [info@rcm.org.uk](mailto:info@rcm.org.uk)  
W: [www.rcm.org.uk](http://www.rcm.org.uk)



**Chief Executive:**  
Gill Walton,  
MA, PGDip, BSc Hons, RM  
**President:**  
Rebecca Davies, RM  
**Patron:**  
HRH The Princess Royal

**2022-05287**

**Comparison of pregnancy outcome during pandemic of COVID-19 and non-pandemic situations, Yazd, Iran, in 2019–2020.**

Saberi H, Ghorashi Z, Loripoor M (2022), Journal of Obstetrics and Gynaecology 23 May 2022, online

This study aimed to compare the pregnancy outcomes during pandemic and non-pandemic COVID-19 in women referred to health service centres in Yazd. This descriptive study was performed based on the information obtained from all pregnant women referred to comprehensive health service centres in Yazd city using census method, between March 21 2019 and December 21 2019 and between March 20 2020 and December 20 2020. The pregnant mothers' information, including their pregnancy outcome, and maternal and neonatal complications, was extracted from the electronic health information system of Yazd city. The obtained data were analysed by Chi-Square test. No significant difference was found between non-pandemic and pandemic COVID-19 situations in most variables. As well, maternal and neonatal death were equally observed in both non-pandemic and pandemic COVID-19 situations. Wanted pregnancy, post term birth, multiple pregnancy and caesarean section rates were found to be higher in pandemic than non-pandemic COVID-19 ( $p < .001$ ). Reported abortion, screening for foetal aneuploidy in the first and second trimesters as well as the number of episodes of prenatal care during COVID-19 pandemic were significantly lower than those of non-pandemic period ( $p < .001$ ). The outcome of pregnancy during the pandemic was not significantly different from that of non-pandemic situation.

**Impact Statement**

What is already known on this subject? Studies already showed COVID-19 in pregnancy alter the maternal and neonatal outcomes in different degrees compared with pregnant individuals without COVID-19. However, it is not clear that pregnancy outcome dose alter during pandemic of COVID-19 compared to non-pandemic situations in general population?

What do the results of this study add? The results of this study revealed that the outcome of pregnancy during pandemic was not significantly different from that of non-pandemic situation.

What are the implications of these findings for clinical practice and/or further research? According to results of this study, we can ensure pregnant women in the situation of pandemic COVID-19 that they are not in greater risk. We suggest future research should be done for comparison of pregnancy outcome in the situation of delta variant pandemic with non-pandemic COVID-19. (Author)

---

**2022-05270**

**Impact of COVID-19 disease and vaccination on maternal/fetal inflammatory response, placental pathology, and perinatal outcomes.**

Boelig RC, Aghai Z, Chauhdury S, et al (2022), American Journal of Obstetrics & Gynecology (AJOG) 28 May 2022, online

Research letter aiming to evaluate the impact of COVID-19 disease and vaccination on the maternal-fetal unit through the inflammatory cytokine panel at delivery, placental pathology and perinatal outcomes. Results show that COVID-19 disease, but not vaccination, is associated with inflammatory cytokines, placental vascular pathology and preterm delivery. (LDO)

Full URL: <https://doi.org/10.1016/j.ajog.2022.05.049>

---

MIDIRS is part of RCM Information Services Limited which is a company incorporated in England and Wales under company no.11914882 with registered office at 10-18 Union Street, London SE1 1SZ  
RCM Information Services Limited is a subsidiary of The Royal College of Midwives



**The Royal College of Midwives Trust**  
10-18 Union Street  
London  
SE1 1SZ



**Open 24 hours a day, 7 days a week**  
T: 0300 303 0444  
F: +44 20 7312 3536  
E: [info@rcm.org.uk](mailto:info@rcm.org.uk)  
W: [www.rcm.org.uk](http://www.rcm.org.uk)



**Chief Executive:**  
Gill Walton,  
MA, PGDip, BSc Hons, RM  
**President:**  
Rebecca Davies, RM  
**Patron:**  
HRH The Princess Royal

2022-05219

**Outcomes of Pregnant Patients Treated with REGEN-COV during the COVID-19 Pandemic.** Levey NH, Forrest AD, Spielman DW, et al (2022), American Journal of Obstetrics & Gynecology MFM 3 June 2022, online. 100673

Background: Pregnant patients with SARS-CoV-2 infection are at increased risk for severe disease including hospitalization, intensive care admission, ventilatory support and death. Although pregnant patients were excluded from investigational trials for pharmacologic treatments for COVID-19 illness, the National Institutes of Health Treatment Guidelines state efficacious treatments should not be withheld from pregnant patients. Infusion of Casirivimab and Imdevimab (REGEN-COV), a monoclonal antibody therapy, was shown to reduce the risk of coronavirus disease 2019 (COVID-19) related hospitalization or death from any cause and resolved symptoms and reduced severe acute respiratory syndrome coronavirus 2 viral load (SARS-CoV-2) more rapidly than placebo. In July of 2021 the Food and Drug Administration released an Emergency Use Authorization for REGEN-COV. Although pregnant persons were not included in the original trials, given the higher risk of morbidity and mortality in the pregnant population, our institution offered REGEN-COV to our pregnant patients beginning in August of 2021. Side effects after REGEN-COV administration are rare and thought to be secondary to COVID infection rather than REGEN-COV.

Objective: The objective of this study was to track the safety and clinical outcomes of unvaccinated pregnant patients who received REGEN-COV and compare these outcomes to a contemporary cohort of patients who tested positive for SARS-CoV-2 and were eligible but did not receive REGEN-COV. Our hypothesis was that REGEN-COV administration during pregnancy is safe, and that pregnant persons who received REGEN-COV would experience less severe COVID-19 respiratory illness by decreasing length of hospital stay, decreasing ICU admission, and decreasing the need for oxygen and other COVID-19 therapeutics.

Study Design: This is a retrospective cohort study of pregnant patients who either tested positive for SARS-CoV-2 or had a known exposure to a COVID-19 positive person, and therefore have been eligible for REGEN-COV at our institution. Within this cohort, we compared those who received REGEN-COV to those who did not receive REGEN-COV between March 2021 and October 2021 at Grady Memorial Hospital in Atlanta, Georgia. The main outcomes studied were perinatal outcomes, safety data and the clinical course of SARS-CoV-2 infection.

Results: From March 2021 to October 2021, 86 pregnant people tested positive for SARS-CoV-2 via real time-PCR or had a confirmed exposure. Among this group, 36 received REGEN-COV and 50 did not. There were no instances of infusion rate adjustment or discontinuation, anaphylaxis, or death among those individuals who received REGEN-COV. One individual experienced worsening shortness of breath over 24 hours after administration which was classified as an infusion-related reaction. There were not any significant differences in perinatal outcomes, length of hospitalization, rates of ICU admission, additional pharmacologic treatment for COVID-19, or oxygen requirement between the two groups.

Conclusions: Administration of REGEN-COV is safe in pregnancy and did not increase adverse maternal, neonatal, or obstetrical outcomes. There was not a statistically significant difference in COVID-19 related outcomes in our high-risk population. Given the likely safety of this drug in pregnancy and its known benefits in the non-pregnant population, we advocate for continued use of this therapy and encourage the development of future studies to enroll a larger and more diverse cohort to explore its efficacy further. (Author)

---

MIDIRS is part of RCM Information Services Limited which is a company incorporated in England and Wales under company no.11914882 with registered office at 10-18 Union Street, London SE1 1SZ  
RCM Information Services Limited is a subsidiary of The Royal College of Midwives



**The Royal College of Midwives Trust**  
10-18 Union Street  
London  
SE1 1SZ



**Open 24 hours a day, 7 days a week**  
T: 0300 303 0444  
F: +44 20 7312 3536  
E: [info@rcm.org.uk](mailto:info@rcm.org.uk)  
W: [www.rcm.org.uk](http://www.rcm.org.uk)



**Chief Executive:**  
Gill Walton,  
MA, PGDip, BSc Hons, RM  
**President:**  
Rebecca Davies, RM  
**Patron:**  
HRH The Princess Royal

**2022-05211**

**Associations between COVID-19 lockdown and post-lockdown on the mental health of pregnant women, postpartum women and their partners from the Queensland family cohort prospective study.** Clifton VL, Kumar S, Borg D, et al (2022), BMC Pregnancy and Childbirth vol 22, no 468, 4 June 2022

#### Background

There are very few developed countries where physical isolation and low community transmission has been reported for COVID-19 but this has been the experience of Australia. The impact of physical isolation combined with low disease transmission on the mental health of pregnant women is currently unknown and there have been no studies examining the psychological experience for partners of pregnant women during lockdown. The aim of the current study was to examine the impact of the first COVID-19 lockdown in March 2020 and post lockdown from August 2020 on the mental health of pregnant women or postpartum women and their partners.

#### Methods

Pregnant women and their partners were prospectively recruited to the study before 24 weeks gestation and completed various questionnaires related to mental health and general wellbeing at 24 weeks gestation and then again at 6 weeks postpartum. The Depression, Anxiety and Stress Scale (DASS-21) and the Edinburgh Postnatal Depression Scale (EPDS) were used as outcome measures for the assessment of mental health in women and DASS-21 was administered to their partners. This analysis encompasses 3 time points where families were recruited; before the pandemic (Aug 2018-Feb 2020), during lockdown (Mar-Aug 2020) and after the first lockdown was over (Sept-Dec 2020).

#### Results

There was no significant effect of COVID-19 lockdown and post lockdown on depression or postnatal depression in women when compared to a pre-COVID-19 subgroup. The odds of pregnant women or postpartum women experiencing severe anxiety was more than halved in women during lockdown relative to women in the pre-COVID-19 period (OR = 0.47; 95%CI: 0.27–0.81; P = 0.006). Following lockdown severe anxiety was comparable to the pre-COVID-19 women. Lockdown did not have any substantial effects on stress scores for pregnant and postpartum women. However, a substantial decrease of over 70% in the odds of severe stress was observed post-lockdown relative to pre-COVID-19 levels. Partner's depression, anxiety and stress did not change significantly with lockdown or post lockdown.

#### Conclusion

A reproductive age population appear to be able to manage the impact of lockdown and the pandemic with some benefits related to reduced anxiety. (Author)

**Full URL:** <https://doi.org/10.1186/s12884-022-04795-9>

---

MIDIRS is part of RCM Information Services Limited which is a company incorporated in England and Wales under company no.11914882 with registered office at 10-18 Union Street, London SE1 1SZ  
RCM Information Services Limited is a subsidiary of The Royal College of Midwives



**The Royal College of Midwives Trust**  
10-18 Union Street  
London  
SE1 1SZ



**Open 24 hours a day, 7 days a week**  
T: 0300 303 0444  
F: +44 20 7312 3536  
E: [info@rcm.org.uk](mailto:info@rcm.org.uk)  
W: [www.rcm.org.uk](http://www.rcm.org.uk)



**Chief Executive:**  
Gill Walton,  
MA, PGDip, BSc Hons, RM  
**President:**  
Rebecca Davies, RM  
**Patron:**  
HRH The Princess Royal

2022-05199

**Maternity care during COVID-19: a qualitative evidence synthesis of women's and maternity care providers' views and experiences.** Flaherty SJ, Delaney H, Matvienko-Sikar K, et al (2022), BMC Pregnancy and Childbirth vol 22, no 438, 26 May 2022

Background

As COVID-19 continued to impact society and health, maternity care, as with many other healthcare sectors across the globe, experienced tumultuous changes. These changes have the potential to considerably impact on the experience of maternity care. To gain insight and understanding of the experience of maternity care during COVID-19, from the perspectives of women and maternity care providers, we undertook a qualitative evidence synthesis (QES).

Methods

The population of interest for the QES were pregnant and postpartum women, and maternity care providers, who provided qualitative data on their experiences of maternity care during COVID-19. The electronic databases of MEDLINE, CINAHL, EMBASE, PsycINFO and the Cochrane COVID study register were systematically searched from 01 Jan 2020 to 13 June 2021. The methodological quality of the included studies was appraised using a modified version of the quality assessment tool, based on 12-criteria, designed by the Evidence for Policy and Practice Information coordinating Centre (EPPI-Centre). Data were extracted by two reviewers independently and synthesised using the Thomas and Harden framework. Confidence in the findings was assessed using the Grading of Recommendations Assessment, Development and Evaluation-Confidence in the Evidence from Reviews of Qualitative research (GRADE-CERQual).

Results

Fifty records relating to 48 studies, involving 9,348 women and 2,538 maternity care providers, were included in the QES. The methodological quality of the studies varied from four studies meeting all 12 quality criteria to two studies meeting one quality criterion only. The synthesis revealed eight prominent themes. Five of these reflected women's experiences: 1) Altered maternity care (women), 2) COVID-related restrictions, 3) Infection prevention and risk, 4) 'the lived reality' – navigating support systems, and 5) Interactions with maternity services. Three themes reflected maternity care providers' experiences: 6) Altered maternity care (providers), 7) Professional and personal impact, and 8) Broader structural impact. Confidence in the findings was high or moderate.

Conclusion

Although some positive experiences were identified, overall, this QES reveals that maternity care during COVID-19 was negatively experienced by both women and maternity care providers. The pandemic and associated changes evoked an array of emotive states for both populations, many of which have the potential to impact on future health and wellbeing. Resource and care planning to mitigate medium- and longer-term adverse sequelae are required.

PROSPERO registration

CRD42021232684. (Author)

Full URL: <https://doi.org/10.1186/s12884-022-04724-w>

MIDIRS is part of RCM Information Services Limited which is a company incorporated in England and Wales under company no.11914882 with registered office at 10-18 Union Street, London SE1 1SZ  
RCM Information Services Limited is a subsidiary of The Royal College of Midwives



**The Royal College of Midwives Trust**  
10-18 Union Street  
London  
SE1 1SZ



**Open 24 hours a day, 7 days a week**  
T: 0300 303 0444  
F: +44 20 7312 3536  
E: [info@rcm.org.uk](mailto:info@rcm.org.uk)  
W: [www.rcm.org.uk](http://www.rcm.org.uk)



**Chief Executive:**  
Gill Walton,  
MA, PGDip, BSc Hons, RM  
**President:**  
Rebecca Davies, RM  
**Patron:**  
HRH The Princess Royal



**2022-05170**

**Effects of the SARS-CoV-2 pandemic on perinatal activity in Yorkshire and the Humber region during 2020: an interrupted time series analysis.** Morgan AS, Bradford CM, Farrow H, et al (2022), Archives of Disease in Childhood: Fetal and Neonatal Edition 11 May 2022, online

Objective To assess the impact of public health measures taken during the COVID-19 pandemic on perinatal health indicators.

Design Interrupted time series analysis comparing periods of the pandemic with the previous 5 years.

Setting Yorkshire and the Humber region, England (2015–2020).

Main outcome measures Relative risk (RR) of stillbirth, extreme preterm (EPT, <27 weeks' gestational age) delivery, hypoxic ischaemic encephalopathy (HIE) and meconium aspiration syndrome (MAS), antenatal transfer for threatened EPT delivery and postnatal transfer for EPT birth, HIE or MAS.

Results Stillbirths fell from 3.7/1000 deliveries prepandemic to 2.9/1000 afterwards; EPT births decreased from 2.5/1000 to 1.8/1000 live births. Following adjustment, during the first lockdown there were decreased antenatal transfers (RR 0.74, 95% CI 0.57 to 0.94) with non-statistically significant increased stillbirth (RR 1.08, 95% CI 0.78 to 1.51) and decreased EPT admissions (RR 0.88, 95% CI 0.60 to 1.29). Over the entire pandemic period, antenatal transfer (RR 0.64, 95% CI 0.55 to 0.76) and EPT birth (RR 0.73, 95% CI 0.56 to 0.94) decreased; stillbirths showed non-statistically significant increases overall (RR 1.21, 95% CI 0.98 to 1.49) but with increasing trend through the pandemic (RR 1.11, 95% CI 1.00 to 1.22). No changes were seen for HIE, MAS, postnatal transfers or in subgroup analyses by ethnicity.

Conclusions Lower rates of antenatal transfer and extreme preterm birth were identified, alongside an apparent increase in stillbirth over time. The findings provide evidence that effects on perinatal activity related to the pandemic changed over time. (Author)

**Full URL:** <http://dx.doi.org/10.1136/archdischild-2021-323466>

---

MIDIRS is part of RCM Information Services Limited which is a company incorporated in England and Wales under company no.11914882 with registered office at 10-18 Union Street, London SE1 1SZ  
RCM Information Services Limited is a subsidiary of The Royal College of Midwives



**The Royal College of Midwives Trust**  
10-18 Union Street  
London  
SE1 1SZ



**Open 24 hours a day, 7 days a week**  
T: 0300 303 0444  
F: +44 20 7312 3536  
E: [info@rcm.org.uk](mailto:info@rcm.org.uk)  
W: [www.rcm.org.uk](http://www.rcm.org.uk)



**Chief Executive:**  
Gill Walton,  
MA, PGDip, BSc Hons, RM  
**President:**  
Rebecca Davies, RM  
**Patron:**  
HRH The Princess Royal

**2022-05163**

**Preterm birth rates were slightly lower in Denmark during the first year of the COVID-19 pandemic compared with the previous 4 years.** Hansen BM, Cueto H, Petersen JP, et al (2022), Acta Paediatrica 9 May 2022, online

**Aim**

Our aim was to investigate the rates of preterm births, live births and stillbirths in Denmark during the first year of the COVID-19 pandemic.

**Methods**

This was a national, cross-sectional registry-based study that used the Danish Newborn Quality database, which covers all births in Denmark. The proportions of preterm births were compared between the COVID-19 pandemic period of 1 March 2020 to 28 February 2021 and the preceding 4-year pre-pandemic period.

**Results**

We studied 60 323 and 244 481 newborn infants from the pandemic and pre-pandemic periods, respectively. The proportion of preterm live births and stillbirths declined slightly, from 6.29% during the pre-pandemic period to 6.02% during the pandemic period. This corresponded to a relative risk (RR) of 0.96, with a 95% confidence interval (CI) of 0.93–0.99 during the pandemic. The RRs for extremely preterm, very preterm and moderately preterm infants were 0.88 (95% CI 0.76–1.02), 0.91 (95% CI 0.82–1.02) and 0.97 (95% CI 0.93–1.01), respectively.

**Conclusion**

This comparative study showed a small reduction in just over 4%, from 6.29 to 6.02% in the proportion of all preterm births during the pandemic period, compared with the previous four pandemic-free years. There were no differences between subcategories of preterm births. (Author)

---

MIDIRS is part of RCM Information Services Limited which is a company incorporated in England and Wales under company no.11914882 with registered office at 10-18 Union Street, London SE1 1SZ  
RCM Information Services Limited is a subsidiary of The Royal College of Midwives



**The Royal College of Midwives Trust**  
10-18 Union Street  
London  
SE1 1SZ



**Open 24 hours a day, 7 days a week**  
T: 0300 303 0444  
F: +44 20 7312 3536  
E: [info@rcm.org.uk](mailto:info@rcm.org.uk)  
W: [www.rcm.org.uk](http://www.rcm.org.uk)



**Chief Executive:**  
Gill Walton,  
MA, PGDip, BSc Hons, RM  
**President:**  
Rebecca Davies, RM  
**Patron:**  
HRH The Princess Royal

**2022-05145**

**Association between intimate partner violence and prenatal anxiety and depression in pregnant women: a cross-sectional survey during the COVID-19 epidemic in Shenzhen, China.** Wu F, Zhou L, Chen C, et al (2022), *BMJ Open* vol 12, no 5, May 2022, e055333

Objectives Intimate partner violence (IPV) against women remains a major global public health problem with harmful consequences for individuals and society. People's lifestyles have been greatly affected by the COVID-19 pandemic. This study investigated the prevalence of and relationship between IPV and anxiety and depression in pregnant Chinese women during the pandemic.

Design Cross-sectional study.

Setting This investigation was conducted in Shenzhen City, Guangdong Province, China from 15 September to 15 December 2020.

Participants A total of 3434 pregnant women were screened with the Abuse Assessment Screen Questionnaire to evaluate IPV and General Anxiety Disorder and Patient Health Questionnaire to evaluate symptoms of anxiety and depression, respectively. Pregnant women with perinatal health records at Shenzhen District Maternity and Child Healthcare Hospitals who consented to participate were enrolled. Women with psychotic disorders such as schizophrenia, mania or substance dependence and pregnant women who refused to participate were excluded. Data were analysed with the  $\chi^2$  test and by logistic regression analysis.

Results The prevalence of IPV among pregnant women was 2.2%. Mental violence was the most common type of violence (2.2%), followed by physical (0.6%) and sexual (0.7%) violence. The prevalence of anxiety and depression symptoms was 9.8% and 6.9%, respectively. After adjusting for covariates, there was a statistically significant association between IPV and prenatal anxiety (OR=4.207, 95% CI: 2.469 to 7.166) and depression (OR=3.864, 95% CI: 2.095 to 7.125).

Conclusions IPV increased the risk of prenatal anxiety and depression in pregnant women in China during the COVID-19 pandemic. Efforts should be made by the government and civil society to promote long-lasting antenatal interventions to ensure the safety and protect the mental health of pregnant women. (Author)

**Full URL:** <http://dx.doi.org/10.1136/bmjopen-2021-055333>

---

MIDIRS is part of RCM Information Services Limited which is a company incorporated in England and Wales under company no.11914882 with registered office at 10-18 Union Street, London SE1 1SZ  
RCM Information Services Limited is a subsidiary of The Royal College of Midwives



**The Royal College of Midwives Trust**  
10-18 Union Street  
London  
SE1 1SZ



**Open 24 hours a day, 7 days a week**  
T: 0300 303 0444  
F: +44 20 7312 3536  
E: [info@rcm.org.uk](mailto:info@rcm.org.uk)  
W: [www.rcm.org.uk](http://www.rcm.org.uk)



**Chief Executive:**  
Gill Walton,  
MA, PGDip, BSc Hons, RM  
**President:**  
Rebecca Davies, RM  
**Patron:**  
HRH The Princess Royal

**2022-05125**

**First-time mothers' experiences of antenatal education during the COVID-19 pandemic.** Cassar E, Spiteri G (2022), MIDIRS Midwifery Digest vol 32, no 2, June 2022, pp 196-200

**Background:** The COVID-19 global pandemic has resulted in a change in pregnant women's habits and in the models of care that support them during their pregnancies. Social restrictions and quarantine have changed the way antenatal education has been accessed and delivered.

**Aim and objectives:** This study aimed to explore mothers' lived experiences of antenatal education during the COVID-19 pandemic. The objectives of the study were to explore: the experience of antenatal education in relation to pregnancy; the influence of antenatal education on the childbirth experience, and the experience of antenatal education in relation to the postpartum period.

**Methods:** The qualitative paradigm was used to investigate this phenomenon among a purposive sample of nine primiparous mothers who were selected from a breastfeeding clinic at the main state hospital in Malta via an intermediary. A semi-structured interview schedule was specifically designed for the purpose of the study and was used to elicit the data. The interviews were held virtually and were audio-recorded with the participants' consent.

**Results:** Antenatal classes were considered an essential source of information, especially during the COVID-19 pandemic. Despite this, participants highlighted how antenatal education did not fully prepare them for the realities of their labour, birth and their transition to parenthood. Moreover, it transpired that the COVID-19 pandemic created a heightened sense of anxiety, fear and stress among expectant women.

**Conclusions:** This study suggests that antenatal education should start earlier during pregnancy to offer gestation-specific advice and education should also continue well into the postpartum. Since virtual antenatal education classes have offered support to many childbearing women during this unprecedented time more resources that support and enhance web-based antenatal education should be allocated to this important niche of care. (Author)

---

MIDIRS is part of RCM Information Services Limited which is a company incorporated in England and Wales under company no.11914882 with registered office at 10-18 Union Street, London SE1 1SZ  
RCM Information Services Limited is a subsidiary of The Royal College of Midwives



**The Royal College of Midwives Trust**  
10-18 Union Street  
London  
SE1 1SZ



**Open 24 hours a day, 7 days a week**  
T: 0300 303 0444  
F: +44 20 7312 3536  
E: [info@rcm.org.uk](mailto:info@rcm.org.uk)  
W: [www.rcm.org.uk](http://www.rcm.org.uk)



**Chief Executive:**  
Gill Walton,  
MA, PGDip, BSc Hons, RM  
**President:**  
Rebecca Davies, RM  
**Patron:**  
HRH The Princess Royal

**2022-05098**

**Reduction in preterm birth rates during and after the COVID-19 lockdown in Queensland Australia.** Jasper B, Stillerova T, Anstey C, et al (2022), Australian and New Zealand Journal of Obstetrics and Gynaecology (ANZJOG) 17 May 2022, online

#### Background

Preventative strategies for preterm birth are lacking. Recent evidence proposed COVID-19 lockdowns may have contributed to changes in preterm birth.

#### Aims

To determine the prevalence of preterm birth and birth outcomes during and after the COVID-19 lockdown at the Sunshine Coast University Hospital and the overall state of Queensland, Australia.

#### Methods

Retrospective cohort analysis of all births in Queensland including the Sunshine Coast University Hospital, during two epochs, April 1–May 31, 2020 (lockdown) and June 1–July 31, 2020 (post-lockdown), compared to antecedent calendar-matched periods in 2018–2019. Prevalence of preterm birth, stillbirth, and late terminations were examined.

#### Results

There were 64 989 births in Queensland from April to July 2018–2020. At the Sunshine Coast University Hospital, there was a significantly higher chance of birth at term during both lockdown (odds ratio (OR) 1.81, 95% CI 1.17, 2.79;  $P = 0.007$ ) and post-lockdown (OR 2.01, 95% CI 1.27, 3.18;  $P = 0.003$ ). At the same centre, prevalence of preterm birth was 5.5% (30/547) during lockdown, compared to 9.1% (100/1095) in previous years, a 40.0% relative reduction ( $P = 0.016$ ). At this centre during lockdown, emergency caesareans concurrently decreased ( $P < 0.01$ ) and instrumental vaginal births increased ( $P < 0.01$ ). In Queensland overall, there was a nonsignificant decrease in the prevalence of preterm birth during lockdown.

#### Conclusions

There is a link between lockdown and a reduction in the prevalence of preterm birth on the Sunshine Coast. The cause is speculative at present, although increased influenza vaccination rates, decreased transmission of infections, and improved air quality may have been favourable in reducing preterm birth. Further research is needed to determine a causal link. (Author)

**Full URL:** <https://doi.org/10.1111/ajo.13538>

---

MIDIRS is part of RCM Information Services Limited which is a company incorporated in England and Wales under company no.11914882 with registered office at 10-18 Union Street, London SE1 1SZ  
RCM Information Services Limited is a subsidiary of The Royal College of Midwives



**The Royal College of Midwives Trust**  
10-18 Union Street  
London  
SE1 1SZ



**Open 24 hours a day, 7 days a week**  
T: 0300 303 0444  
F: +44 20 7312 3536  
E: [info@rcm.org.uk](mailto:info@rcm.org.uk)  
W: [www.rcm.org.uk](http://www.rcm.org.uk)



**Chief Executive:**  
Gill Walton,  
MA, PGDip, BSc Hons, RM  
**President:**  
Rebecca Davies, RM  
**Patron:**  
HRH The Princess Royal

**2022-05070**

**Expectant parents' perceptions of healthcare and support during COVID-19 in the UK: a thematic analysis.** Aydin E, Glasgow KA, Weiss SM, et al (2022), Journal of Reproductive and Infant Psychology 17 May 2022, online

#### Background

In response to the COVID-19 pandemic, expectant parents experienced changes in the availability and uptake of both National Health Service (NHS) community and hospital-based healthcare.

#### Objective

To examine how COVID-19 and its societal related restrictions have impacted the provision of healthcare support for pregnant women during the COVID-19 pandemic.

#### Method

A thematic analysis using an inductive approach was undertaken using data from open-ended responses to the national COVID in Context of Pregnancy, Infancy and Parenting (CoCoPIP) Study online survey (n = 507 families).

#### Findings

The overarching theme identified was the way in which the changes to healthcare provision increased parents' anxiety levels, and feelings of not being supported. Five sub-themes, associated with the first wave of the pandemic, were identified: (1) rushed and/or fewer antenatal appointments, (2) lack of sympathy from healthcare workers, (3) lack of face-to-face appointments, (4) requirement to attend appointments without a partner, and (5) requirement to use PPE. A sentiment analysis, that used quantitative techniques, revealed participant responses to be predominantly negative (50.1%), with a smaller proportion of positive (21.8%) and neutral (28.1%) responses found.

#### Conclusion

This study provides evidence indicating that the changes to healthcare services for pregnant women during the pandemic increased feelings of anxiety and have left women feeling inadequately supported. Our findings highlight the need for compensatory social and emotional support for new and expectant parents while COVID-19 related restrictions continue to impact on family life and society. (Author)

**Full URL:** <https://doi.org/10.1080/02646838.2022.2075542>

---

MIDIRS is part of RCM Information Services Limited which is a company incorporated in England and Wales under company no.11914882 with registered office at 10-18 Union Street, London SE1 1SZ  
RCM Information Services Limited is a subsidiary of The Royal College of Midwives



**The Royal College of Midwives Trust**  
10-18 Union Street  
London  
SE1 1SZ



**Open 24 hours a day, 7 days a week**  
T: 0300 303 0444  
F: +44 20 7312 3536  
E: [info@rcm.org.uk](mailto:info@rcm.org.uk)  
W: [www.rcm.org.uk](http://www.rcm.org.uk)



**Chief Executive:**  
Gill Walton,  
MA, PGDip, BSc Hons, RM  
**President:**  
Rebecca Davies, RM  
**Patron:**  
HRH The Princess Royal

**2022-04978**

**Navigating uncertainty alone: A grounded theory analysis of women's psycho-social experiences of pregnancy and childbirth during the COVID-19 pandemic in London.** Montgomery E, De Backer K, Easter A, et al (2022), *Women and Birth: Journal of the Australian College of Midwives* 16 May 2022, online

**Problem**

Maternity care underwent substantial reconfiguration in the United Kingdom during the COVID-19 pandemic.

**Background**

COVID-19 posed an unprecedented public health crisis, risking population health and causing a significant health system shock.

**Aim**

To explore the psycho-social experiences of women who received maternity care and gave birth in South London during the first 'lockdown'.

**Methods**

We recruited women (N = 23) to semi-structured interviews, conducted virtually. Data were recorded, transcribed, and analysed by hand. A Classical Grounded Theory Analysis was followed including line-by-line coding, focused coding, development of super-categories followed by themes, and finally the generation of a theory.

**Findings**

Iterative and inductive analysis generated six emergent themes, sorted into three dyadic pairs: 1 & 2: Lack of relational care vs. Good practice persisting during the pandemic; 3 & 4: Denying the embodied experience of pregnancy and birth vs. Trying to keep everyone safe; and 5 & 6: Removed from support network vs. Importance of being at home as a family. Together, these themes interact to form the theory: 'Navigating uncertainty alone'.

**Discussion**

Women's pregnancy and childbirth journeys during the pandemic were reported as having positive and negative experiences which would counteract one-another. Lack of relational care, denial of embodied experiences, and removal from support networks were counterbalanced by good practice which persisted, understanding staff were trying to keep everyone safe, and renewed importance in the family unit.

**Conclusion**

Pregnancy can be an uncertain time for women. This was compounded by having to navigate their maternity journey alone during the COVID-19 pandemic. (Author)

**Full URL:** <https://doi.org/10.1016/j.wombi.2022.05.002>

---

**2022-04923**

**Changes in women's health service seeking behaviours and the impact of telehealth during COVID-19: Insights from the 1800MyOptions service.** Subasinghe AK, Mogharbel C, Hill D, et al (2021), *Australian and New Zealand Journal of Obstetrics and Gynaecology (ANZJOG)* vol 61, no 5, October 2021, pp e26-e27

Letter to the editor presenting an audit of the 1800MyOptions service operated by Women's Health Victoria about contraception, pregnancy options and sexual health. The audit revealed an increased demand for pregnancy options counselling and an increased proportion of callers at a later gestation. (LDO)

**Full URL:** <https://doi.org/10.1111/ajo.13409>

---

**2022-04921**

**Pregnancy and the risk of severe coronavirus disease 2019 infection: methodological challenges and research recommendations.** Savitz D, Bengtson AM, Hardy E, et al (2022), *BJOG: An International Journal of Obstetrics and Gynaecology* vol 129, no 2, January 2022, pp 192-195

Commentary addressing the methodological challenges encountered in examining the impact of pregnancy on severity of COVID-19 infection. Highlights factors which increase the risk of bias including confounding, increased surveillance and enhanced clinical response to illness. (LDO)

**Full URL:** <https://doi.org/10.1111/1471-0528.16935>

---

MIDIRS is part of RCM Information Services Limited which is a company incorporated in England and Wales under company no.11914882 with registered office at 10-18 Union Street, London SE1 1SZ  
RCM Information Services Limited is a subsidiary of The Royal College of Midwives



**The Royal College of Midwives Trust**  
10-18 Union Street  
London  
SE1 1SZ



**Open 24 hours a day, 7 days a week**  
T: 0300 303 0444  
F: +44 20 7312 3536  
E: [info@rcm.org.uk](mailto:info@rcm.org.uk)  
W: [www.rcm.org.uk](http://www.rcm.org.uk)



**Chief Executive:**  
Gill Walton,  
MA, PGDip, BSc Hons, RM  
**President:**  
Rebecca Davies, RM  
**Patron:**  
HRH The Princess Royal

2022-04850

**Neighborhood Characteristics and Racial Disparities in Severe Acute Respiratory Syndrome Coronavirus 2 (SARS-CoV-2) Seropositivity in Pregnancy.** Burris HH, Mullin AM, Dhudasia MB, et al (2022), *Obstetrics & Gynecology* 5 May 2022, online

**OBJECTIVE:**

To quantify the extent to which neighborhood characteristics contribute to racial and ethnic disparities in severe acute respiratory syndrome coronavirus 2 (SARS-CoV-2) seropositivity in pregnancy.

**METHODS:**

This cohort study included pregnant patients who presented for childbirth at two hospitals in Philadelphia, Pennsylvania from April 13 to December 31, 2020. Seropositivity for SARS-CoV-2 was determined by measuring immunoglobulin G and immunoglobulin M antibodies by enzyme-linked immunosorbent assay in discarded maternal serum samples obtained for clinical purposes. Race and ethnicity were self-reported and abstracted from medical records. Patients' residential addresses were geocoded to obtain three Census tract variables: community deprivation, racial segregation (Index of Concentration at the Extremes), and crowding. Multivariable mixed effects logistic regression models and causal mediation analyses were used to quantify the extent to which neighborhood variables may explain racial and ethnic disparities in seropositivity.

**RESULTS:**

Among 5,991 pregnant patients, 562 (9.4%) were seropositive for SARS-CoV-2. Higher seropositivity rates were observed among Hispanic (19.3%, 104/538) and Black (14.0%, 373/2,658) patients, compared with Asian (3.2%, 13/406) patients, White (2.7%, 57/2,133) patients, and patients of another race or ethnicity (5.9%, 15/256) ( $P < .001$ ). In adjusted models, per SD increase, deprivation (adjusted odds ratio [aOR] 1.16, 95% CI 1.02–1.32) and crowding (aOR 1.15, 95% CI 1.05–1.26) were associated with seropositivity, but segregation was not (aOR 0.90, 95% CI 0.78–1.04). Mediation analyses revealed that crowded housing may explain 6.7% (95% CI 2.0–14.7%) of the Hispanic–White disparity and that neighborhood deprivation may explain 10.2% (95% CI 0.5–21.1%) of the Black–White disparity.

**CONCLUSION:**

Neighborhood deprivation and crowding were associated with SARS-CoV-2 seropositivity in pregnancy in the prevaccination era and may partially explain high rates of SARS-CoV-2 seropositivity among Black and Hispanic patients. Investing in structural neighborhood improvements may reduce inequities in viral transmission. (Author)

---

MIDIRS is part of RCM Information Services Limited which is a company incorporated in England and Wales under company no.11914882 with registered office at 10-18 Union Street, London SE1 1SZ  
RCM Information Services Limited is a subsidiary of The Royal College of Midwives



**The Royal College of Midwives Trust**  
10-18 Union Street  
London  
SE1 1SZ



**Open 24 hours a day, 7 days a week**  
T: 0300 303 0444  
F: +44 20 7312 3536  
E: [info@rcm.org.uk](mailto:info@rcm.org.uk)  
W: [www.rcm.org.uk](http://www.rcm.org.uk)



**Chief Executive:**  
Gill Walton,  
MA, PGDip, BSc Hons, RM  
**President:**  
Rebecca Davies, RM  
**Patron:**  
HRH The Princess Royal



**2022-04848**

**Recruitment of Pregnant Women to Randomised Trials of COVID 19 Treatments, and Pharmaceutical Treatments Received Outside such Trials: A Research Article.** Green O, Young EM, Oberman J, et al (2022),

European Journal of Obstetrics & Gynecology and Reproductive Biology 13 May 2022, online

**Objectives**

To document how many pregnant women with COVID-19 reported in the literature had participated in randomised trials, what treatments they received outside such trials and compare the latter with evidence-based treatment recommendations.

**Study Design:** a systematic review of observational studies.

**Methods**

Two clinical trial registries were searched to identify COVID-19 trials open to pregnant women. Studies were then extracted from a regularly updated list of scientific case reports and case series of confirmed or suspected maternal COVID-19 in pregnancy to identify the number of women enrolled into a trial and the pharmaceutical treatments they received outside such trials.

**Results**

156 studies (case reports, case series and registries) reporting 43,185 pregnant women with COVID-19, after de-duplication. Of these 2,671 (6.2%) were potentially eligible for a randomised trial but only seven women (0.26%) were reported to have enrolled.

For 2,839 women the papers included information on treatment received, 1515/2829 (54%) women had received  $\geq 1$  treatment and in total a COVID-19 pharmaceutical treatment was administered 1,296 times outside of a trial. In 566 (44%) cases the treatments administered to the pregnant women were not recommended by the National Institutes of Health (NIH) at the time of administration.

Of 179 case reports of women with COVID 19 in pregnancy, 109/179 women received  $\geq 1$  COVID-19 pharmaceutical treatment and in total COVID-19 experimental pharmaceutical treatments were administered 274 times.

**Conclusion**

During the early phase of the COVID-19 pandemic, pregnant women excluded from randomised trials did not avoid unproven or ineffective treatments. (Author)

**Full URL:** <https://doi.org/10.1016/j.ejogrb.2022.05.009>

---

MIDIRS is part of RCM Information Services Limited which is a company incorporated in England and Wales under company no.11914882 with registered office at 10-18 Union Street, London SE1 1SZ  
RCM Information Services Limited is a subsidiary of The Royal College of Midwives



**The Royal College of Midwives Trust**  
10-18 Union Street  
London  
SE1 1SZ



**Open 24 hours a day, 7 days a week**  
T: 0300 303 0444  
F: +44 20 7312 3536  
E: [info@rcm.org.uk](mailto:info@rcm.org.uk)  
W: [www.rcm.org.uk](http://www.rcm.org.uk)



**Chief Executive:**  
Gill Walton,  
MA, PGDip, BSc Hons, RM  
**President:**  
Rebecca Davies, RM  
**Patron:**  
HRH The Princess Royal

2022-04847

**The impact of mitigation measures on perinatal outcomes during the first nine months of the COVID-19 pandemic: A systematic review with meta-analysis.** Hawco S, Rolnik DL, Woolner A, et al (2022),

European Journal of Obstetrics & Gynecology and Reproductive Biology 13 May 2022, online

Worldwide reports have produced conflicting data on perinatal outcomes during the COVID-19 pandemic. This systematic review and meta-analysis addressed the effect of mitigation measures against COVID-19 on preterm birth, stillbirth, low birth weight, and NICU admission during the first nine months of the pandemic.

A search was performed using MEDLINE, Embase and SCOPUS for manuscripts published up until 24th May 2021. Studies that reported perinatal outcomes (preterm birth, stillbirth, low birth weight, NICU admission) during the COVID-19 pandemic with a pre-pandemic control period were included. Risk of bias assessment was performed using ROBINS-I tool. RevMan5 was used to perform meta-analysis with random-effects models. A score of the stringency of mitigation measures was calculated from the Oxford COVID-19 Government Response Tracker.

Thirty-eight studies of moderate to serious risk of bias were included, with varied methodology, analysis and regional mitigation measures, using stringency index scores. There was no overall effect on preterm birth at less than 37 weeks (OR 0.96, 95% CI 0.92–1.00). However, there was a reduction in preterm birth at less than 37 weeks (OR 0.89, 95% CI 0.81–0.98) and 34 weeks (OR 0.56, 95% CI 0.37–0.83) for iatrogenic births and in singleton pregnancies. There was also a significant reduction in preterm births at less than 34 weeks in studies with above median stringency index scores (OR 0.71, 95% CI 0.58–0.88). There was no effect on risk of stillbirth (OR 1.04, 95% CI 0.90–1.19) or birth weight. NICU admission rates were significantly reduced in studies with above median stringency index scores (OR 0.87, 95% CI 0.78–0.97). The reduction in preterm births in regions with high mitigation measures against SARS-CoV-2 infection is likely driven by a reduction in iatrogenic births. Variability in study design and cohort characteristics need to be considered for future studies to allow further investigation of population level health measures of perinatal outcomes. (Author)

**Full URL:** <https://doi.org/10.1016/j.ejogrb.2022.05.007>

---

MIDIRS is part of RCM Information Services Limited which is a company incorporated in England and Wales under company no.11914882 with registered office at 10-18 Union Street, London SE1 1SZ  
RCM Information Services Limited is a subsidiary of The Royal College of Midwives



**The Royal College of Midwives Trust**  
10-18 Union Street  
London  
SE1 1SZ



**Open 24 hours a day, 7 days a week**  
T: 0300 303 0444  
F: +44 20 7312 3536  
E: [info@rcm.org.uk](mailto:info@rcm.org.uk)  
W: [www.rcm.org.uk](http://www.rcm.org.uk)



**Chief Executive:**  
Gill Walton,  
MA, PGDip, BSc Hons, RM  
**President:**  
Rebecca Davies, RM  
**Patron:**  
HRH The Princess Royal

**2022-04806**

**Analysis of online antenatal education class use via a mobile terminal app during the COVID-19 pandemic.** Chen X-W, Jiang L-Y, Chen Y, et al (2022), BMC Pregnancy and Childbirth vol 22, no 412, 16 May 2022

### Objective

To understand the use of online antenatal education classes accessed via the Mother and Child Health Handbook app during the COVID-19 pandemic in order to provide a basis and suggestions for optimizing Internet education during pregnancy under public health emergencies.

### Methods

We compared and analyzed the use of online antenatal education classes via the Mother and Child Health Handbook app in Hangzhou in 2019 and 2020 (during the COVID-19 pandemic).

### Results

Between January 1, 2019, and December 31, 2020, a total of 229,794 pregnant women created files and registered for the app, including 124,273 women in 2019 and 105,521 women in 2020. More pregnant women participated in online antenatal education learning ( $n = 36,379/34.5\%$  vs.  $29,226/23.5\%$ ,  $p = 0.000$ ) in 2020 than in 2019. The proportion of pregnant women in the 18–34-year-old group who participated in online learning was higher than that in the advanced age group, and the difference was statistically significant (2019: 24.3% vs. 18.8%,  $p = 0.000$ ) (2020: 35.7% vs. 27.4%,  $p = 0.000$ ). More pregnant women accessed online antenatal education during early pregnancy ( $n = 13,463/37.0\%$  vs.  $9088/31.1\%$ ,  $p = 0.000$ ) in 2020 than in 2019. Similar percentages of pregnant women participated in online antenatal education during mid-pregnancy ( $n = 15,426/52.8\%$  vs.  $19,269/53.0\%$ ,  $p = 0.639$ ) in 2019 and 2020. Fewer pregnant women accessed online antenatal education during late pregnancy ( $n = 10,246/28.2\%$  vs.  $9476/32.4\%$ ,  $p = 0.000$ ) in 2020 than in 2019. Fewer pregnant women choose to take 'Puerperal Health' courses in 2020 than in 2019 (early pregnancy: 36.20% vs. 42.79%,  $p = 0.000$ ; mid-pregnancy: 41.65% vs. 48.19%,  $p = 0.000$ ; late pregnancy: 55.31% vs. 58.41%,  $p = 0.000$ ). Fewer pregnant women choose to take 'Psychological Adjustment' courses in 2020 than in 2019 (early pregnancy: 21.59% vs. 29.60%,  $p = 0.000$ ; mid-pregnancy: 26.20% vs. 40.50%,  $p = 0.000$ ; late pregnancy: 12.79% vs. 42.53%,  $p = 0.000$ ). More pregnant women choose to study 'Nutrition and Exercise' in 2020 than in 2019 (early pregnancy: 44.48% vs. 25.95%,  $p = 0.000$ ; mid-pregnancy: 47.77% vs. 40.75%,  $p = 0.000$ ; late pregnancy: 55.94% vs. 42.99%,  $p = 0.000$ ). "Pregnancy Care and Fetal Development" was the most selected course by pregnant women in early pregnancy (2019: 67.50%; 2020: 71.39%) and middle pregnancy (2019: 67.01%; 2020: 82.05%), and the proportion in 2020 was higher than it was in 2019. "Baby care" was the most selected course by pregnant women in late pregnancy, and the proportion in 2020 was higher than it was in 2019 (78.31% vs. 72.85%).

### Conclusion

During the COVID-19 pandemic, online antenatal education was well-used by pregnant women. More women participated in the online antenatal education modules during the COVID-19 pandemic than during 2019. The proportion of choosing different courses for pregnant women before and after the COVID-19 epidemic varied, and the learning course needs of pregnant women in different trimesters were different. (Author)

Full URL: <https://doi.org/10.1186/s12884-022-04745-5>

MIDIRS is part of RCM Information Services Limited which is a company incorporated in England and Wales under company no.11914882 with registered office at 10-18 Union Street, London SE1 1SZ  
RCM Information Services Limited is a subsidiary of The Royal College of Midwives



**The Royal College of Midwives Trust**  
10-18 Union Street  
London  
SE1 1SZ



**Open 24 hours a day, 7 days a week**  
T: 0300 303 0444  
F: +44 20 7312 3536  
E: [info@rcm.org.uk](mailto:info@rcm.org.uk)  
W: [www.rcm.org.uk](http://www.rcm.org.uk)



**Chief Executive:**  
Gill Walton,  
MA, PGDip, BSc Hons, RM  
**President:**  
Rebecca Davies, RM  
**Patron:**  
HRH The Princess Royal

**2022-04804**

**Implementing “Online Communities” for pregnant women in times of COVID-19 for the promotion of maternal well-being and mother-to-infant bonding: a pretest–posttest study.** Potharst ES, Schaeffer MA, Gunning C, et al (2022), BMC Pregnancy and Childbirth vol 22, no 415, 18 May 2022

#### Background

The Coronavirus Disease 2019 (COVID-19) pandemic elevated the risk for mental health problems in pregnant women, thereby increasing the risk for long-term negative consequences for mother and child well-being. There was an immediate need for easily accessible interventions for pregnant women experiencing elevated levels of pandemic related stress.

#### Methods

A three-session intervention “Online Communities” (OC) was developed at the beginning of the Dutch lockdown, and implemented by a team of midwives and psychologists specialized in Infant Mental Health. Pretest (N = 34) and posttest (N = 17) measurements of depressive symptoms, worries about COVID-19 and worries in general, and mother-to-infant bonding were administered, as well as a posttest evaluation.

#### Results

At pretest, the OC group was compared to two reference groups of pregnant women from an ongoing pregnancy cohort study: a COVID-19 (N = 209) and pre-COVID-19 reference group (N = 297). OC participants had significantly more depressive symptoms than both reference groups, and less positive feelings of bonding than the COVID-19 but not the pre-COVID-19 reference group. Compared to pretest, significant decreases in depressive symptoms (with significantly less participants scoring above cut-off) and worries about COVID-19 (large effect sizes) and worries in general (moderate to large effect size) were found at posttest for the OC participants. No significant improvement was found in bonding. Participants rated the intervention positively.

#### Conclusions

The current study provides initial evidence supporting the idea that OC is a promising and readily accessible intervention for pregnant women experiencing stress due to the COVID-19 pandemic, and possibly also applicable to other stressors.

#### Trial registration

This intervention was registered in the Netherlands Trial Registration (registration number Trial NL8842, registration date 18/08/2020). (Author)

**Full URL:** <https://doi.org/10.1186/s12884-022-04729-5>

---

MIDIRS is part of RCM Information Services Limited which is a company incorporated in England and Wales under company no.11914882 with registered office at 10-18 Union Street, London SE1 1SZ  
RCM Information Services Limited is a subsidiary of The Royal College of Midwives



**The Royal College of Midwives Trust**  
10-18 Union Street  
London  
SE1 1SZ



**Open 24 hours a day, 7 days a week**  
T: 0300 303 0444  
F: +44 20 7312 3536  
E: [info@rcm.org.uk](mailto:info@rcm.org.uk)  
W: [www.rcm.org.uk](http://www.rcm.org.uk)



**Chief Executive:**  
Gill Walton,  
MA, PGDip, BSc Hons, RM  
**President:**  
Rebecca Davies, RM  
**Patron:**  
HRH The Princess Royal

**2022-04766**

**Shared risk factors for COVID-19 and preeclampsia in the first trimester: An observational study.** Serrano B, Mendoza M, Garcia-Aguilar P, et al (2022), *Acta Obstetrica et Gynecologica Scandinavica* 3 May 2022, online

#### Introduction

The association between preeclampsia and coronavirus disease 2019 (COVID-19) is under study. Previous publications have hypothesized the existence of shared risk factors for both conditions or a deficient trophoblastic invasion as possible explanations for this association. The primary aim of this study was to examine baseline risk factors measured in the first-trimester combined screening for preeclampsia in pregnant women with COVID-19 compared with the general population. A secondary aim of this study was to compare risk factors among patients with mild and severe COVID-19.

#### Material and Methods

This was an observational retrospective study conducted at Vall d'Hebron Hospital Campus (Catalonia, Spain). Study patients were 231 pregnant women undergoing the first-trimester screening for preeclampsia and positive for severe acute respiratory syndrome coronavirus 2 between February 2020 and September 2021. The reference cohort were 13 033 women of the general population from six centers across Catalonia from May 2019 to June 2021. Based on the need for hospitalization, patients were classified in two groups: mild and severe COVID-19. First-trimester screening for preeclampsia included maternal history, mean arterial blood pressure, mean uterine artery pulsatility index (UtAPI), placental growth factor (PIGF), and pregnancy-associated plasma protein-A (PAPP-A).

#### Results

The proportion of cases at high risk for preeclampsia was significantly higher among the COVID-19 group compared with the general population (19.0% and 13.2%, respectively;  $p = 0.012$ ). When analyzing risk factors for preeclampsia individually, women with COVID-19 had higher median body mass index (25.2 vs. 24.5,  $p = 0.041$ ), higher UtAPI multiple of the median (MoM) (1.08 vs. 1.00,  $p < 0.001$ ), higher incidence of chronic hypertension (2.8% vs. 0.9%,  $p = 0.015$ ), and there were fewer smokers (5.7% vs. 11.6%,  $p = 0.007$ ). The MoMs of PIGF and PAPP-A did not differ significantly between both groups (0.96 vs. 0.97,  $p = 0.760$  and 1.00 vs. 1.01,  $p = 0.432$ ; respectively).

#### Conclusions

In patients with COVID-19, there was a higher proportion of women at high risk for preeclampsia at the first-trimester screening than in the general population, mainly because of maternal risk factors, rather than placental signs of a deficient trophoblastic invasion. (Author)

**Full URL:** <https://doi.org/10.1111/aogs.14371>

---

MIDIRS is part of RCM Information Services Limited which is a company incorporated in England and Wales under company no.11914882 with registered office at 10-18 Union Street, London SE1 1SZ  
RCM Information Services Limited is a subsidiary of The Royal College of Midwives



**The Royal College of Midwives Trust**  
10-18 Union Street  
London  
SE1 1SZ



**Open 24 hours a day, 7 days a week**  
T: 0300 303 0444  
F: +44 20 7312 3536  
E: [info@rcm.org.uk](mailto:info@rcm.org.uk)  
W: [www.rcm.org.uk](http://www.rcm.org.uk)



**Chief Executive:**  
Gill Walton,  
MA, PGDip, BSc Hons, RM  
**President:**  
Rebecca Davies, RM  
**Patron:**  
HRH The Princess Royal

**2022-04760**

**Gestational diabetes mellitus and COVID-19: results from the CRONOS study.** Kleinwechter HJ, Weber KS, Mingers N, et al (2022), American Journal of Obstetrics & Gynecology (AJOG) 14 May 2022, online

#### BACKGROUND

Gestational diabetes mellitus is one of the most frequent pregnancy complications with a global prevalence of 13.4% in 2021. Pregnant women with COVID-19 and gestational diabetes mellitus are 3.3 times more likely to be admitted to an intensive care unit compared to women without gestational diabetes mellitus. Data on the association of gestational diabetes mellitus with maternal and neonatal pregnancy outcomes in SARS-CoV-2-infected pregnant women are lacking.

#### OBJECTIVE

The aim of our study was to investigate whether gestational diabetes mellitus is an independent risk factor for adverse maternal and fetal/neonatal outcomes in pregnant women with COVID-19.

#### STUDY DESIGN

CRONOS (COVID-19 Obstetric and Neonatal Outcome Study) is a registry-based multicentric prospective observational study from Germany and Linz/Austria. Pregnant women with clinically confirmed COVID-19 were enrolled between April 3, 2020 and August 24, 2021, at any stage of pregnancy. Obstetricians and neonatologists of 115 hospitals actively provided data to CRONOS. For collecting data a cloud based electronic data platform was developed. Women and neonates were followed until hospital discharge. Information on demographic characteristics, comorbidities, medical history, COVID-19 associated symptoms and treatments, pregnancy and birth outcomes were entered by the local sites. Information on the periconceptional body mass index was collected. A primary combined maternal endpoint was defined as (1) admission to an intensive care unit (including maternal mortality), and/or (2) viral pneumonia, and/or (3) oxygen supplementation. A primary combined fetal/neonatal endpoint was defined as (1) stillbirth >24+0 weeks of pregnancy, and/or (2) neonatal death <7 days postnatal, and/or (3) transfer to a neonatal intensive care unit. Multivariable logistic regression analysis was performed to evaluate the modulating impact of gestational diabetes mellitus on the defined endpoints.

#### RESULTS

Of the 1,490 women with COVID-19 (mean age  $31.0 \pm 5.2$  years, 40.7% nulliparous), 9.4% (n=140) were diagnosed with gestational diabetes mellitus, of these, 42.9% were treated with insulin. Overall, gestational diabetes mellitus was not associated with an adverse maternal outcome (odds Ratio: 1.50; 95% confidence interval: 0.88; 2.57). In overweight/obese women, however, gestational diabetes mellitus was independently associated with the primary maternal outcome (adjusted odds ratio 2.69; 95% confidence interval: 1.43; 5.07). Overweight/obese women with gestational diabetes mellitus requiring insulin treatment were found to have an increased risk for a severe course of COVID-19 (adjusted odds ratio 3.05; 95% confidence interval: 1.38; 6.73). Adverse maternal outcome was more common when COVID-19 was diagnosed with or short after gestational diabetes mellitus diagnosis compared to COVID-19 diagnosis prior to gestational diabetes mellitus diagnosis (19.6% vs. 5.6%,  $p < 0.05$ ). Maternal gestational diabetes mellitus and maternal preconceptional body mass index  $>25 \text{ kg/m}^2$  increased the risk of adverse fetal/neonatal outcome (adjusted odds ratio 1.83; 95% confidence interval: 1.05; 3.18). Furthermore, overweight/obesity (irrespective of gestational diabetes mellitus status) were influential factors for the maternal (adjusted odds ratio 1.87; 95% confidence interval: 1.26; 2.75) and neonatal primary endpoints (1.81; 95% confidence interval: 1.32; 2.48) in comparison to women with underweight or normal weight.

#### CONCLUSION

Gestational diabetes mellitus, combined with periconceptional overweight/obesity, is independently associated with severe maternal course of COVID-19, especially when the mothers require insulin and COVID-19 is diagnosed with or after gestational diabetes mellitus diagnosis. These combined factors exhibited a moderate effect on neonatal outcomes. Women with gestational diabetes mellitus and body mass index  $>25 \text{ kg/m}^2$  are a particularly vulnerable group in the case of COVID-19.

(Author)

---

MIDIRS is part of RCM Information Services Limited which is a company incorporated in England and Wales under company no.11914882 with registered office at 10-18 Union Street, London SE1 1SZ  
RCM Information Services Limited is a subsidiary of The Royal College of Midwives



**The Royal College of Midwives Trust**  
10-18 Union Street  
London  
SE1 1SZ



**Open 24 hours a day, 7 days a week**  
T: 0300 303 0444  
F: +44 20 7312 3536  
E: [info@rcm.org.uk](mailto:info@rcm.org.uk)  
W: [www.rcm.org.uk](http://www.rcm.org.uk)



**Chief Executive:**  
Gill Walton,  
MA, PGDip, BSc Hons, RM  
**President:**  
Rebecca Davies, RM  
**Patron:**  
HRH The Princess Royal

## 2022-04723

### **Histopathological features in advanced abdominal pregnancies co-infected with SARS-CoV-2 and HIV-1 infections: A case evaluation.**

Ramphal S, Govender N, Singh S, et al (2022), European Journal of Obstetrics & Gynecology and Reproductive Biology: X vol 15, August 2022, 100153

#### Objectives

This study aims to provide a semi-qualitative histopathological report of the dual SARS-CoV-2 and HIV infected placentae in the third trimester of Advanced Abdominal Pregnancy (AAP).

#### Study design

Four AAP placentae in the third trimester of pregnancy (two positive for HIV-1 and two positives for SARS-CoV-2) were histologically examined.

#### Results

The SARS-CoV-2+ HIV+ placentae were dysmorphic in shape compared to the flattened disc-like shape noted in the SARS-CoV-2+HIV-, SARS-CoV-2-HIV+ and SARS-CoV-2-HIV- placentae. Diffused syncytial knots and syncytial degeneration were observed in all placentae. Intermittent cytotrophoblast increase, perivillous and intravillous fibrin deposition, mononuclear inflammatory cells with widespread degeneration/necrosis of the syncytiotrophoblast and microcalcification were pronounced in the SARS-CoV-2+HIV+ compared to the SARS-CoV-2+HIV- placentae. Vascular pathological changes included thrombi, ectasis, mural hypertrophy and atherotic vessels.

#### Conclusion

Elevated syncytial trophoblast injury, villitis, microcalcifications and mineralisation of the syncytial basement membrane in the AAP placentae may be due to SARS-CoV-2 viral transgression instead of HIV infection alone. Vascular malperfusion is suggestive of a hypoxic insult arising from a compensatory response to meet the fetal oxygen and nutrient demands of an AAP. Placentae from HIV infected women on antiretroviral treatment were characterised by vascular malperfusion. (Author)

**Full URL:** <https://doi.org/10.1016/j.eurox.2022.100153>

---

## 2022-04718

### **Maternity services: visiting restrictions [written answer].** Northern Ireland Assembly (2022), Hansard Written question AQW 214/22-27, 19 May 2022

The Minister of Health responds to a written question asked by Mr Pádraig Delargy, regarding whether all COVID-19 restrictions have been relaxed in maternity services and settings. (LDO)

**Full URL:** <http://aims.niassembly.gov.uk/questions/printquestionssummary.aspx?docid=373642>

---

## 2022-04662

### **Impact of COVID-19 pandemic on maternal and neonatal morbidities in the United States.** Gulersen M, Lenchner E, Grunebaum A, et al (2022), American Journal of Obstetrics & Gynecology MFM 20 May 2022, online. 100667

Research letter aiming to evaluate whether the COVID-19 pandemic had an impact on overall rates of preterm birth and maternal and neonatal complications in the United States. Results suggest there was no increase in preterm birth during the pandemic, but the likelihood of complications increased such as gestational hypertension, pre-eclampsia and pre-gestational diabetes. (LDO)

**Full URL:** <https://doi.org/10.1016/j.ajogmf.2022.100667>

---

MIDIRS is part of RCM Information Services Limited which is a company incorporated in England and Wales under company no.11914882 with registered office at 10-18 Union Street, London SE1 1SZ  
RCM Information Services Limited is a subsidiary of The Royal College of Midwives



**The Royal College of Midwives Trust**  
10-18 Union Street  
London  
SE1 1SZ



**Open 24 hours a day, 7 days a week**  
T: 0300 303 0444  
F: +44 20 7312 3536  
E: [info@rcm.org.uk](mailto:info@rcm.org.uk)  
W: [www.rcm.org.uk](http://www.rcm.org.uk)



**Chief Executive:**  
Gill Walton,  
MA, PGDip, BSc Hons, RM  
**President:**  
Rebecca Davies, RM  
**Patron:**  
HRH The Princess Royal

**2022-04606**

**Evaluation of maternal-infant dyad inflammatory cytokines in pregnancies affected by maternal SARS-CoV-2 infection in early and late gestation.** Taglauer ES, Dhole Y, Akuamoah-Boateng J, et al (2022), Journal of Perinatology 21 April 2022, online

**Objective**

SARS-CoV-2 infection induces significant inflammatory cytokine production in adults, but infant cytokine signatures in pregnancies affected by maternal SARS-CoV-2 are less well characterized. We aimed to evaluate cytokine profiles of mothers and their infants following COVID-19 in pregnancy.

**Study design**

Serum samples at delivery from 31 mother-infant dyads with maternal SARS-CoV-2 infection in pregnancy (COVID) were examined in comparison to 29 control dyads (Control). Samples were evaluated using a 13-plex cytokine assay.

**Results**

In comparison with controls, interleukin (IL)-6 and interferon gamma-induced protein 10 (IP-10) were higher in COVID maternal and infant samples ( $p < 0.05$ ) and IL-8 uniquely elevated in COVID infant samples ( $p < 0.05$ ). Significant elevations in IL-6, IP-10, and IL-8 were found among both early (1st/2nd Trimester) and late (3rd Trimester) maternal SARS-CoV-2 infections.

**Conclusions**

Maternal SARS-CoV-2 infections throughout gestation are associated with increased maternal and infant inflammatory cytokines at birth with potential to impact long-term infant health. (Author)

**Full URL:** <https://doi.org/10.1038/s41372-022-01391-9>

---

MIDIRS is part of RCM Information Services Limited which is a company incorporated in England and Wales under company no.11914882 with registered office at 10-18 Union Street, London SE1 1SZ  
RCM Information Services Limited is a subsidiary of The Royal College of Midwives



**The Royal College of Midwives Trust**  
10-18 Union Street  
London  
SE1 1SZ



**Open 24 hours a day, 7 days a week**  
T: 0300 303 0444  
F: +44 20 7312 3536  
E: [info@rcm.org.uk](mailto:info@rcm.org.uk)  
W: [www.rcm.org.uk](http://www.rcm.org.uk)



**Chief Executive:**  
Gill Walton,  
MA, PGDip, BSc Hons, RM  
**President:**  
Rebecca Davies, RM  
**Patron:**  
HRH The Princess Royal



**2022-04547**

**The rate of SARS-CoV-2 among asymptomatic non-immunised low-risk parturient women between the two waves.** Al-Hussaini TK, EzzEldin AM, Shaaban OM, et al (2022), Journal of Obstetrics and Gynaecology 29 April 2022, online

This study was conducted to evaluate the rate of severe acute respiratory syndrome coronavirus 2 (SARS-CoV-2) infection and immunity among asymptomatic non-immunised low-risk parturient women and their newborns. A cross-sectional study conducted in a tertiary hospital during the nadir period of new cases in Egypt. All asymptomatic pregnant, low risk and non-immunised women were included. All eligible participants had been subjected to SARS-CoV-2 nasopharyngeal swabs according to CDC and sampling of maternal and umbilical blood to evaluate the presence of coronavirus disease 2019 (COVID-19) IgM and IgG antibodies by immunochromatographic assay. Two cases out of 171 (1.2%) parturient women were tested positive for PCR swab to COVID-19 infection. Furthermore, COVID-19 IgG and IgM antibodies testing showed that 67.8% of women were negative for both IgG and IGM, 24.6% were positive for IgG only, 4.1% were positive for IgM only, while 3.5% were positive for both IgG and IgM. Regarding neonatal testing for immunity, 28.1% of the neonates were positive to IgG only and none for IgM.

The rate of positive PCR patients among asymptomatic low-risk parturient women was 1.2%. About quarter of women had got herd immunity as evident by positive IgG antibodies. IgG antibodies transferred to the neonates in almost all cases.

#### Impact Statement

What is already known on this subject? Coronavirus disease 2019 (COVID-19), caused by severe acute respiratory syndrome coronavirus 2 (SARS-CoV-2), has become a global public health emergency. Asymptomatic pregnant women with coronavirus disease can transmit their infection to their newborn, family members and the health care providers.

What do the results of this study add? The study showed very low (1.2%) prevalence of COVID positive cases among asymptomatic pregnant women admitted to our facility. Only two cases out of 171 parturient women tested PCR positive for COVID-19 infection (1.2%). SARS-Cov-2 IgG and IgM antibodies testing showed, about a quarter (24.6%) were positive for IgG antibodies, 4.1% were positive for IgM antibodies, while 3.5% were positive for both IgG and IgM. On the other hand, 28.1% of the neonates were positive to IgG only and none of the newborns had had IgM antibodies in their cord blood.

What are the implications of these findings for clinical practice and/or further research? The first wave of COVID-19 pandemic in Egypt left behind at least a quarter of pregnant women with a positive antibody denoting some immunity. This immunity is usually transmitted to the neonates in almost all cases. (Author)

---

MIDIRS is part of RCM Information Services Limited which is a company incorporated in England and Wales under company no.11914882 with registered office at 10-18 Union Street, London SE1 1SZ  
RCM Information Services Limited is a subsidiary of The Royal College of Midwives



**The Royal College of Midwives Trust**  
10-18 Union Street  
London  
SE1 1SZ



**Open 24 hours a day, 7 days a week**  
T: 0300 303 0444  
F: +44 20 7312 3536  
E: [info@rcm.org.uk](mailto:info@rcm.org.uk)  
W: [www.rcm.org.uk](http://www.rcm.org.uk)



**Chief Executive:**  
Gill Walton,  
MA, PGDip, BSc Hons, RM  
**President:**  
Rebecca Davies, RM  
**Patron:**  
HRH The Princess Royal

**2022-04321**

**Seroprevalence of SARS-CoV-2 antibodies among first-trimester pregnant women during the second wave of the pandemic in India.** Sharma KA, Singh N, Hillman S, et al (2022), International Journal of Gynecology & Obstetrics 24 March 2022, online

#### Objective

Data on the immune response to SARS-CoV-2 during pregnancy are lacking and the potential role and effect of SARS-CoV-2 vaccination in pregnancy is yet to be completely investigated.

#### Method

This is a cross-sectional observational study wherein pregnant women were tested for SARS-CoV-2 immunoglobulin M and immunoglobulin G levels, irrespective of their infective status or presence or symptomatology.

#### Result

Of the 220 pregnant women tested, 160 (72.7%) were SARS-CoV-2 IgG positive, 37 (16.8%) were SARS-CoV-2 IgM positive and 27 (16.9%) were both IgG and IgM positive. The average antibody titer found was 10.49 BAU/ml ( $\pm 14.0$ ) and 0.6 ( $\pm 0.55$ ) for anti-SARS-CoV-2 IgG and IgM non neutralizing antibodies respectively. ROC analysis for SARS-CoV-2 IgG positivity showed a cut-off value of 1.19 with a sensitivity of 99.3% (0.99 AUC, 95% CI) and specificity of 98.3% (0.99 AUC, 95% CI), respectively. Similarly, ROC analysis for SARS-CoV-2 IgM positivity showed a cut-off value of 1 with a sensitivity of 97.3% (0.99 AUC, 95% CI) and specificity of 98.9% (0.99 AUC, 95% CI), respectively.

#### Conclusion

First trimester sero-molecular screening suggests a high prevalence of COVID antibodies in the study population of pregnant women in the first trimester, without the patients being symptomatic. (Author)

**Full URL:** <https://doi.org/10.1002/ijgo.14189>

---

MIDIRS is part of RCM Information Services Limited which is a company incorporated in England and Wales under company no.11914882 with registered office at 10-18 Union Street, London SE1 1SZ  
RCM Information Services Limited is a subsidiary of The Royal College of Midwives



**The Royal College of Midwives Trust**  
10-18 Union Street  
London  
SE1 1SZ



**Open 24 hours a day, 7 days a week**  
T: 0300 303 0444  
F: +44 20 7312 3536  
E: [info@rcm.org.uk](mailto:info@rcm.org.uk)  
W: [www.rcm.org.uk](http://www.rcm.org.uk)



**Chief Executive:**  
Gill Walton,  
MA, PGDip, BSc Hons, RM  
**President:**  
Rebecca Davies, RM  
**Patron:**  
HRH The Princess Royal

**2022-04229**

**Effectiveness and safety of available treatments for COVID-19 during pregnancy: a critical review.** Favilli A, Gentili MM, Raspa F, et al (2022), Journal of Maternal-Fetal and Neonatal Medicine vol 35, no 11, 2022, pp 2174-2187

#### Background

COVID-19 is a pandemic disease caused by the SARS-CoV-2 and it spread globally in the last few months. The complete lack of specific treatment forced clinicians to use old drugs, chosen for their efficacy against similar viruses or their in vitro activity. Trials on patients are ongoing but the majority of information comes from small case series and single center reports. We aimed to provide a literature review on the putative effectiveness and safety of available treatments for COVID-19 in pregnant women.

#### Methods

We reviewed all the available literature concerning the drugs that have been used in the treatment of COVID-19 during pregnancy and whose safe assumption during pregnancy had been demonstrated by clinical studies (i.e. including studies on other infectious diseases). Drugs contra-indicated during pregnancy or with unknown adverse effects were not included in our review.

#### Results and conclusions

Clinical trials are not often conducted among pregnant patients for safety reasons and this means that drugs that may be effective in general population cannot be used for pregnant women due to the lack of knowledge of side effects in this category of people. The choice to use a specific drug for COVID-19 in pregnancy should take into account benefits and possible adverse events in each single case. In the current situation of uncertainty and poor knowledge about the management of COVID-19 during pregnancy, this present overview may provide useful information for physicians with practical implications. (Author)

**Full URL:** <https://doi.org/10.1080/14767058.2020.1774875>

---

**2022-04194**

**Relationship between pregnancy and coronavirus: what we know.** Forestieri S, Marcialis MA, Migliore L, et al (2022), Journal of Maternal-Fetal and Neonatal Medicine vol 35, no 10, 2022, pp 1997-2008

The identification in China in December 2019 of a new coronavirus (SARS-CoV-2) immediately rekindled the spotlight on a problem also addressed in the past during the epidemics of SARS in 2002–2003 and MERS in 2012: the implications of a possible infection during pregnancy, both for pregnant women and for fetuses and infants. Pregnancy is characterized by some changes involving both the immune system and the pulmonary physiology, exposing the pregnant woman to a greater susceptibility to viral infections and more serious complications. The objective of this review is therefore to analyze the relationship between pregnancy and known coronaviruses, with particular reference to SARS-CoV-2. (Author)

**Full URL:** <https://doi.org/10.1080/14767058.2020.1771692>

---

MIDIRS is part of RCM Information Services Limited which is a company incorporated in England and Wales under company no.11914882 with registered office at 10-18 Union Street, London SE1 1SZ  
RCM Information Services Limited is a subsidiary of The Royal College of Midwives



**The Royal College of Midwives Trust**  
10-18 Union Street  
London  
SE1 1SZ



**Open 24 hours a day, 7 days a week**  
T: 0300 303 0444  
F: +44 20 7312 3536  
E: [info@rcm.org.uk](mailto:info@rcm.org.uk)  
W: [www.rcm.org.uk](http://www.rcm.org.uk)



**Chief Executive:**  
Gill Walton,  
MA, PGDip, BSc Hons, RM  
**President:**  
Rebecca Davies, RM  
**Patron:**  
HRH The Princess Royal

**2022-04171**

**High-risk pregnant women's experiences of the receiving prenatal care in COVID-19 pandemic: a qualitative study.** Mirzakhani K, Shoorab NJ, Akbari A, et al (2022), BMC Pregnancy and Childbirth vol 22, no 363, 26 April 2022

#### Background

Women with high-risk pregnancies are among the most vulnerable groups that require additional precautionary measures against the spread of COVID-19 plus receiving prenatal care. Yet, there is limited information on the status of prenatal care in women with high-risk pregnancies. The purpose of this study was to explore the experiences of women with high-risk pregnancies who were receiving prenatal care during the COVID-19 pandemic.

#### Methods

The present qualitative study was conducted on mothers with high-risk pregnancies from September 2020 to March 2021. Purposeful sampling continued until achieving data saturation. Ghaem, Ommolbanin, and Imam Reza in Mashhad, Iran served as the research environment. Face-to-face and semi-structured interviews were effective data collection methods. Each interview lasted between 20 to 45 min (on average 30). The total number of participants was 31. Data analysis was carried out simultaneously with data collection using the qualitative content analysis method developed by Granheim and Landman (2004).

#### Results

Following the reduction and analysis of data from women in high-risk pregnancies, as well as their perceptions and experiences with health services during the COVID-19 pandemic, eight subcategories and three main categories were identified, including 1) "Negative psychology responses," 2) "Adoption behavior," and 3) "Adjustment of health services in mutual protection."

Fear, anxiety, stress, feelings of loneliness, sadness, depression, guilt, doubt and conflict in receiving services were examples of negative psychological responses. The adaptive behaviors' category reflected the behaviors of women with high-risk pregnancies in the context of the COVID COVID-19 pandemic. The Adjustment of health services in mutual protection indicated that health workers took preventive and protective measures against COVID-19, which, in addition to protecting themselves and their clients against COVID-19, gave women a sense of security.

#### Conclusion

Receiving prenatal care during the COVID-19 pandemic presents challenges for women with high-risk pregnancies, negatively impacting their psychological state and health-seeking behavior. Supportive and preventive care can ensure that women with high-risk pregnancies receive optimal prenatal care that focuses on COVID-19 prevention. We recommend implementing screening, psychological counseling, and education for women with high-risk pregnancies, as well as ensuring that they have access to women-centered health care services. (Author)

**Full URL:** <https://doi.org/10.1186/s12884-022-04676-1>

---

MIDIRS is part of RCM Information Services Limited which is a company incorporated in England and Wales under company no.11914882 with registered office at 10-18 Union Street, London SE1 1SZ  
RCM Information Services Limited is a subsidiary of The Royal College of Midwives



**The Royal College of Midwives Trust**  
10-18 Union Street  
London  
SE1 1SZ



**Open 24 hours a day, 7 days a week**  
T: 0300 303 0444  
F: +44 20 7312 3536  
E: [info@rcm.org.uk](mailto:info@rcm.org.uk)  
W: [www.rcm.org.uk](http://www.rcm.org.uk)



**Chief Executive:**  
Gill Walton,  
MA, PGDip, BSc Hons, RM  
**President:**  
Rebecca Davies, RM  
**Patron:**  
HRH The Princess Royal

**2022-04134**

**Severe maternal morbidity in pregnant patients with SARS-CoV-2 infection.** Gulersen M, Rochelson B, Shan W, et al (2022), American Journal of Obstetrics & Gynecology MFM vol 4, no 4, July 2022, 100636

#### Background

While the increased risk of severe illness and adverse pregnancy outcomes associated with severe acute respiratory syndrome coronavirus 2 (SARS-CoV-2) infection in pregnancy are well described, the association of infection with severe maternal morbidity (SMM) has not been well characterized.

#### Objective

To evaluate the risk of SMM associated with SARS-CoV-2 infection in pregnancy.

#### Study Design

A multicenter retrospective cohort study of all pregnant patients who had SARS-CoV-2 testing and delivered in a New York health system between March 1st, 2020 and March 1st, 2021. Patients with missing test results were excluded. The primary outcome of SMM, derived from the American College of Obstetricians and Gynecologists and Society for Maternal-Fetal Medicine example list of diagnoses and complications, was compared between two groups: patients who tested positive for SARS-CoV-2 during pregnancy versus patients who tested negative. Secondary outcomes included subgroups of SMM. Multivariable logistic regression was used to adjust for potential confounders such as maternal demographics, neighborhood socioeconomic status, hospital location, and pregnancy-related complications. A subanalysis was performed to determine whether the risk of severe obstetrical hemorrhage and hypertension/neurologic morbidity differed based on timing of SARS-CoV-2 infection: those who tested positive for SARS-CoV-2 at their delivery hospitalization (i.e. active infection), and those who tested positive during pregnancy but negative at their delivery hospitalization (i.e. resolved infection).

#### Results

Of the 22,483 patients included, 1,653 (7.4%) tested positive for SARS-CoV-2 infection. Patients with SARS-CoV2 infection were more commonly Black, multiracial, Hispanic, non-English speaking, using Medicaid insurance, multiparous, and from neighborhoods with a lower socioeconomic status. Patients with SARS-CoV-2 infection were at an increased risk of SMM compared to those without infection (9.3 vs. 6.5%; adjusted OR 1.52, 95% CI 1.21-1.88). Patients with SARS-CoV-2 infection were also at an increased risk of severe obstetrical hemorrhage (1.1% vs. 0.5%; aOR 1.78, 95% CI 1.04-2.88), pulmonary morbidity (2.0% vs. 0.5%; aOR 3.90, 95% CI 2.52-5.89), and intensive care unit (ICU) admission (1.8% vs. 0.5%; aOR 3.29, 95% CI 2.09-5.04) compared to those without infection. The risk of hypertension/neurologic morbidity was similar between the two groups. Timing of SARS-CoV-2 infection (whether active or resolved at time of delivery) was not associated with the risk of severe obstetrical hemorrhage or hypertension/neurologic morbidity compared to those without infection.

#### Conclusions

SARS-CoV-2 infection in pregnancy is associated with an increased risk of SMM, severe obstetrical hemorrhage, pulmonary morbidity, and ICU admission. These data highlight the need for obstetrical unit preparedness in caring for patients with SARS-CoV-2 infection, continued public health efforts aimed at minimizing risk of infection and support including this select population in investigational therapy and vaccine trials. (Author)

**Full URL:** <https://doi.org/10.1016/j.ajogmf.2022.100636>

---

MIDIRS is part of RCM Information Services Limited which is a company incorporated in England and Wales under company no.11914882 with registered office at 10-18 Union Street, London SE1 1SZ  
RCM Information Services Limited is a subsidiary of The Royal College of Midwives



**The Royal College of Midwives Trust**  
10-18 Union Street  
London  
SE1 1SZ



**Open 24 hours a day, 7 days a week**  
T: 0300 303 0444  
F: +44 20 7312 3536  
E: [info@rcm.org.uk](mailto:info@rcm.org.uk)  
W: [www.rcm.org.uk](http://www.rcm.org.uk)



**Chief Executive:**  
Gill Walton,  
MA, PGDip, BSc Hons, RM  
**President:**  
Rebecca Davies, RM  
**Patron:**  
HRH The Princess Royal

**2022-04127**

**The influence of structural racism, pandemic stress, and SARS-CoV-2 infection during pregnancy with adverse birth**

**outcomes.** Janevic T, Lieb W, Ibroci E, et al (2022), American Journal of Obstetrics & Gynecology MFM vol 4, no 4, July 2022, 100649

Background: Structural racism and pandemic-related stress from the COVID-19 pandemic may increase risk of adverse birth outcomes.

Objective: Our objective was to examine associations between neighborhood measures of structural racism and pandemic stress with three outcomes: SARS-CoV-2 infection, preterm birth (PTB) and delivering a newborn small-for-gestational-age (SGA). Our secondary objective was to investigate the joint associations of SARS-CoV-2 infection during pregnancy and neighborhood measures on PTB and SGA.

Study Design: We analyzed data for 967 patients from a prospective cohort of pregnant persons in New York City, comprised of 367 White persons (38%), 169 Black persons (17%), 293 Latina persons (30%), 87 Asian persons (9%), 41 persons of unknown race-ethnicity (4%), and 10 of unknown race-ethnicity (1%). We evaluated structural racism (social/built structural disadvantage, racial-economic segregation) and pandemic-related stress (community COVID-19 mortality, community unemployment rate increase) in quartiles by zip code. SARS-CoV-2 serologic enzyme-linked immunosorbent assay was performed on blood samples from pregnant persons. We ascertained preterm birth (PTB) and small-for-gestational age (SGA) from an electronic medical record database. We used log-binomial regression with robust standard error for clustering by zip code to estimate associations of each neighborhood measure separately with three outcomes: SARS-CoV-2 infection, PTB, and SGA. Covariates included maternal age, parity, insurance status, and BMI. Models with PTB and SGA as the dependent variables additionally adjusted for SARS-CoV-2 infection.

Results: 193 (20%) persons were SARS-CoV-2 seropositive, and the overall risk of PTB and SGA were 8.4% and 9.8%, respectively. Among birthing persons in neighborhoods in the highest quartile of structural disadvantage (n=190), 94% were non-White, 50% had public insurance, 41% were obese, 32% were seropositive, 11% delivered preterm, and 12% delivered an infant SGA. Among birthing persons in neighborhoods in the lowest quartile of structural disadvantage (n=360), 39% were non-White, 17% had public insurance, 15% were obese, 9% were seropositive, 6% delivered preterm, and 10% delivered an infant SGA. In adjusted analyses structural racism measures and community unemployment were associated with both SARS-CoV-2 infection and PTB, but not SGA. High vs. low structural disadvantage was associated with an adjusted relative risk (aRR) of 2.6 for infection (95% Confidence Interval (CI)=1.7, 3.9) and 1.7 for PTB (95%CI=1.0, 2.9); high vs. low racial-economic segregation was associated with aRR of 1.9 (95% CI=1.3, 2.8) for infection and 2.0 (95%CI=1.3, 3.2) for PTB; high vs. low community unemployment increase was associated with aRR of 1.7 (95% CI=1.2, 1.5) for infection and 1.6 (95%CI=1.0, 2.8) for PTB. COVID-19 mortality rate was associated with SARS-CoV-2 infection, but not PTB or SGA. SARS-CoV-2 infection was not independently associated with birth outcomes. We found no interaction between SARS-CoV-2 infection and neighborhood measures on PTB or SGA.

Conclusions: Neighborhood measures of structural racism were associated with both SARS-CoV-2 infection and PTB, but these associations were independent and did not have a synergistic effect. Community unemployment rate increases were also associated with an increased risk of PTB independently of SARS-CoV-2 infection. Mitigating these factors might reduce the impact of the pandemic on pregnant people. (Author)

---

MIDIRS is part of RCM Information Services Limited which is a company incorporated in England and Wales under company no.11914882 with registered office at 10-18 Union Street, London SE1 1SZ  
RCM Information Services Limited is a subsidiary of The Royal College of Midwives



**The Royal College of Midwives Trust**  
10-18 Union Street  
London  
SE1 1SZ



**Open 24 hours a day, 7 days a week**  
T: 0300 303 0444  
F: +44 20 7312 3536  
E: [info@rcm.org.uk](mailto:info@rcm.org.uk)  
W: [www.rcm.org.uk](http://www.rcm.org.uk)



**Chief Executive:**  
Gill Walton,  
MA, PGDip, BSc Hons, RM  
**President:**  
Rebecca Davies, RM  
**Patron:**  
HRH The Princess Royal

**2022-04016**

**Effect of restrictions imposed due to COVID-19 pandemic on the antenatal care and pregnancy outcomes: a prospective observational study from rural North India.** Goyal LD, Garg P, Verma M, et al (2022), BMJ Open vol 12, no 4, April 2022, e059701

**Objectives** To assess the difficulties faced by the pregnant women in seeking appropriate antenatal care due to the restrictions imposed during the COVID-19 pandemic; assess the difficulties encountered during delivery and postpartum period; the suitability of the teleconsultation services offered; effect of COVID-19 infection on pregnancy outcomes and the effect of restrictions on the nutrition profile of the pregnant women.

**Design** Prospective observational study.

**Setting and participants** We included 1374 pregnant women from the rural areas of three districts of Punjab, India registered at government health centres before the implementation of lockdown due to the COVID-19 pandemic on 24 March 2020.

**Primary and secondary outcome measures** The primary outcome was the difficulties faced by the women during their pregnancies due to restrictions imposed during the lockdown. The secondary outcomes included the effect of COVID-19 infections on pregnancy outcomes, satisfaction from the telemedicine services and restrictions on the nutrition profile of the pregnant women.

**Results** One-third of the women (38.4%) considered their last pregnancy unplanned. Women faced difficulties due to the restrictions in getting adequate nutrition (76.5%), accessing transportation facilities (35.4%), consultations from doctors (22.4%) or getting an ultrasonography scan (48.7%). One-fifth (21.9%) of women could not access safe abortion services. Only 3.6% of respondents ever took any teleconsultation services offered by the government. Most of them felt unsatisfied compared with routine visits (77.5%). COVID-19-infected women were primarily asymptomatic (76.1%), but there was a high incidence of preterm birth (42.8%). Frontline workers could visit 64.3% of the women in the postpartum period despite restrictions.

**Conclusions** Lockdown compromised the antenatal care in our study area while the frontline workers attempted to minimise the inconvenience. Telemedicine services did not prove to be of many benefits to pregnant women and should only work as a supplement to the existing protocols of antenatal care. (Author)

**Full URL:** <http://dx.doi.org/10.1136/bmjopen-2021-059701>

---

MIDIRS is part of RCM Information Services Limited which is a company incorporated in England and Wales under company no.11914882 with registered office at 10-18 Union Street, London SE1 1SZ  
RCM Information Services Limited is a subsidiary of The Royal College of Midwives



**The Royal College of Midwives Trust**  
10-18 Union Street  
London  
SE1 1SZ



**Open 24 hours a day, 7 days a week**  
T: 0300 303 0444  
F: +44 20 7312 3536  
E: [info@rcm.org.uk](mailto:info@rcm.org.uk)  
W: [www.rcm.org.uk](http://www.rcm.org.uk)



**Chief Executive:**  
Gill Walton,  
MA, PGDip, BSc Hons, RM  
**President:**  
Rebecca Davies, RM  
**Patron:**  
HRH The Princess Royal

**2022-04012**

**Impact of the COVID-19 pandemic on intimate partner violence during pregnancy: evidence from a multimethods study of recently pregnant women in Ethiopia.** Wood SN, Yirgu R, Wondimagegnehu A, et al (2022), *BMJ Open* vol 12, no 4, April 2022, e055790

**Objectives** This multimethods study aimed to: (1) compare the prevalence of intimate partner violence (IPV) during pregnancy pre-COVID-19 and during the COVID-19 pandemic using quantitative data and (2) contextualise pregnant women's IPV experiences during the COVID-19 pandemic through supplemental interviews.

**Design** Quantitative analyses use data from Performance Monitoring for Action-Ethiopia, a cohort of 2868 pregnant women that collects data at pregnancy, 6 weeks, 6 months and 1-year postpartum. Following 6-week postpartum survey, in-depth semistructured interviews contextualised experiences of IPV during pregnancy with a subset of participants (n=24).

**Participants** All pregnant women residing within six regions of Ethiopia, covering 91% of the population, were eligible for the cohort study (n=2868 completed baseline survey). Quantitative analyses were restricted to the 2388 women with complete 6-week survey data (retention=82.7%). A purposive sampling frame was used to select qualitative participants on baseline survey data, with inclusion criteria specifying completion of quantitative 6-week interview after the onset of the COVID-19 pandemic, and indication of IPV experience.

**Interventions** A State of Emergency in Ethiopia was declared in response to the COVID-19 pandemic approximately halfway through 6-week postpartum interview, enabling a natural experiment (n=1405 pre-COVID-19; n=983 during-COVID-19).

**Primary outcome measures** IPV during pregnancy was assessed via the 10-item Revised Conflict and Tactics Scale.

**Results** 1-in-10 women experienced any IPV during pregnancy prior to COVID-19 (10.5%), and prevalence of IPV during pregnancy increased to 15.1% during the COVID-19 pandemic (aOR=1.51; p=0.02). Stratified by residence, odds of IPV during the pandemic increased for urban women only (aOR=2.09; p=0.03), however, IPV prevalence was higher in rural regions at both time points. Qualitative data reveal COVID-19-related stressors, namely loss of household income and increased time spent within the household, exacerbated IPV.

**Conclusions** These multimethods results highlight the prevalent, severe violence that pregnant Ethiopian women experience, with pandemic-related increases concentrated in urban areas. Integration of IPV response and safety planning across the continuum of care can mitigate impact. (Author)

**Full URL:** <http://dx.doi.org/10.1136/bmjopen-2021-055790>

---

MIDIRS is part of RCM Information Services Limited which is a company incorporated in England and Wales under company no.11914882 with registered office at 10-18 Union Street, London SE1 1SZ  
RCM Information Services Limited is a subsidiary of The Royal College of Midwives



**The Royal College of Midwives Trust**  
10-18 Union Street  
London  
SE1 1SZ



**Open 24 hours a day, 7 days a week**  
T: 0300 303 0444  
F: +44 20 7312 3536  
E: [info@rcm.org.uk](mailto:info@rcm.org.uk)  
W: [www.rcm.org.uk](http://www.rcm.org.uk)



**Chief Executive:**  
Gill Walton,  
MA, PGDip, BSc Hons, RM  
**President:**  
Rebecca Davies, RM  
**Patron:**  
HRH The Princess Royal



**2022-04008**

**Effects of the COVID-19 pandemic on antenatal care utilisation in Kenya: a cross-sectional study.** Landrian A, Mboya J, Golub G, et al (2022), *BMJ Open* vol 12, no 4, April 2022, e058312

**Objective** The aim of this study was to assess the effects of COVID-19 on antenatal care (ANC) utilisation in Kenya, including women's reports of COVID-related barriers to ANC and correlates at the individual and household levels.

**Design** Cross-sectional study.

**Setting** Six public and private health facilities and associated catchment areas in Nairobi and Kiambu Counties in Kenya.

**Participants** Data were collected from 1729 women, including 1189 women who delivered in healthcare facilities before the COVID-19 pandemic (from September 2019–January 2020) and 540 women who delivered during the pandemic (from July through November 2020). Women who delivered during COVID-19 were sampled from the same catchment areas as the original sample of women who delivered before to compare ANC utilisation.

**Primary and secondary outcome measures** Timing of ANC initiation, number of ANC visits and adequate ANC utilisation were primary outcome measures. Among only women who delivered during COVID-19 only, we explored women's reports of the pandemic having affected their ability to access or attend ANC as a secondary outcome of interest.

**Results** Women who delivered during COVID-19 had significantly higher odds of delayed ANC initiation (ie, beginning ANC during the second vs first trimester) than women who delivered before (aOR 1.72, 95% CI 1.24 to 2.37), although no significant differences were detected in the odds of attending 4–7 or ≥8 ANC visits versus <4 ANC visits, respectively (aOR 1.12, 95% CI 0.86 to 1.44 and aOR 1.46, 95% CI 0.74 to 2.86). Nearly half (n=255/540; 47%) of women who delivered during COVID-19 reported that the pandemic affected their ability to access ANC.

**Conclusions** Strategies are needed to mitigate disruptions to ANC among pregnant women during pandemics and other public health, environmental, or political emergencies. (Author)

**Full URL:** <http://dx.doi.org/10.1136/bmjopen-2021-060185>

---

MIDIRS is part of RCM Information Services Limited which is a company incorporated in England and Wales under company no.11914882 with registered office at 10-18 Union Street, London SE1 1SZ  
RCM Information Services Limited is a subsidiary of The Royal College of Midwives



**The Royal College of Midwives Trust**  
10-18 Union Street  
London  
SE1 1SZ



**Open 24 hours a day, 7 days a week**  
T: 0300 303 0444  
F: +44 20 7312 3536  
E: [info@rcm.org.uk](mailto:info@rcm.org.uk)  
W: [www.rcm.org.uk](http://www.rcm.org.uk)



**Chief Executive:**  
Gill Walton,  
MA, PGDip, BSc Hons, RM  
**President:**  
Rebecca Davies, RM  
**Patron:**  
HRH The Princess Royal

**2022-03992**

**Factors affecting the mental health of pregnant women using UK maternity services during the COVID-19 pandemic: a qualitative interview study.** McKinlay AR, Fancourt D, Burton A (2022), BMC Pregnancy and Childbirth vol 22, no 313, 12 April 2022

#### Background

People using maternity services in the United Kingdom (UK) have faced significant changes brought on by the COVID-19 pandemic and social distancing regulations. We focused on the experiences of pregnant women using UK maternity services during the pandemic and the impact of social distancing rules on their mental health and wellbeing.

#### Methods

We conducted 23 qualitative semi-structured interviews from June 2020 to August 2021, with women from across the UK who experienced a pregnancy during the pandemic. Nineteen participants in the study carried their pregnancy to term and four had experienced a miscarriage during the pandemic. Interviews took place remotely over video or telephone call, discussing topics such as mental health during pregnancy and use of UK maternity services. We used reflexive thematic analysis to analyse interview transcripts.

#### Results

We generated six higher order themes: [1] Some pregnancy discomforts alleviated by social distancing measures, [2] The importance of relationships that support coping and adjustment, [3] Missed pregnancy and parenthood experiences, [4] The mental health consequences of birth partner and visitor restrictions, [5] Maternity services under pressure, and [6] Lack of connection with staff. Many participants felt a sense of loss over a pregnancy experience that differed so remarkably to what they had expected because of the pandemic. Supportive relationships were important to help cope with pregnancy and pandemic-related changes; but feelings of isolation were compounded for some participants because opportunities to build social connections through face-to-face parent groups were unavailable. Participants also described feeling alone due to restrictions on their partners being present when accessing UK maternity services.

#### Conclusions

Our findings highlight some of the changes that may have affected pregnant women's mental health during the COVID-19 pandemic. Reduced social support and being unable to have a partner or support person present during maternity service use were the greatest concerns reported by participants in this study. Absence of birth partners removed a protective buffer in times of uncertainty and distress. This suggests that the availability of a birth partner or support person must be prioritised wherever possible in times of pandemics to protect the mental health of people experiencing pregnancy and miscarriage.

(Author)

**Full URL:** <https://doi.org/10.1186/s12884-022-04602-5>

**2022-03965**

**Satisfaction of pregnant women in a Facebook group led by midwives with information on prenatal care during the COVID-19 pandemic in Peru.** De La Cruz-Ramirez YM, Olaza-Maguiña AF (2022), The Practising Midwife vol 25, no 5, May 2022, pp 42-46

The objective of this research was to determine the satisfaction of pregnant women in a Facebook group created and directed by midwives to provide information on prenatal care during the COVID-19 pandemic in Peru. A cross-sectional study was carried out, where 226 pregnant women answered an online questionnaire. It was concluded that the majority of pregnant women were fully satisfied with the information provided by midwives on prenatal care, highlighting the credibility of the information. Areas that could have been improved included the variety of topics covered and the way participants were treated. The variety of topics addressed and the cordial treatment among the participants are aspects susceptible to improvement. (Author)

MIDIRS is part of RCM Information Services Limited which is a company incorporated in England and Wales under company no.11914882 with registered office at 10-18 Union Street, London SE1 1SZ  
RCM Information Services Limited is a subsidiary of The Royal College of Midwives



**The Royal College of Midwives Trust**  
10-18 Union Street  
London  
SE1 1SZ



**Open 24 hours a day, 7 days a week**  
T: 0300 303 0444  
F: +44 20 7312 3536  
E: [info@rcm.org.uk](mailto:info@rcm.org.uk)  
W: [www.rcm.org.uk](http://www.rcm.org.uk)



**Chief Executive:**  
Gill Walton,  
MA, PGDip, BSc Hons, RM  
**President:**  
Rebecca Davies, RM  
**Patron:**  
HRH The Princess Royal

## 2022-03948

**Impact of new diagnostic pathway for gestational diabetes in time of COVID-19.** Walker B, Edey J, Hall L, et al (2022), *Obstetric Medicine* 19 April 2022, online

### Background

In April 2020, the diagnostic criteria for gestational diabetes mellitus (GDM) changed in Queensland, with the goal of reducing exposure of pregnant women to COVID-19.

### Methods

A retrospective clinical audit was conducted at a regional hospital to compare the incidence of GDM, and specific maternal and neonatal outcomes four months before and after the change in guidelines was implemented.

### Results

Less than 50% of diagnostic tests were performed according to new guidelines. There was a non-significant increase in the incidence of GDM (13.3% to 15.3%), and pharmacological treatments. Instrumental deliveries ( $p = 0.01$ ) and shoulder dystocia ( $p = 0.04$ ) increased following the change in guidelines. There were no differences in the incidence of elective and emergency caesarean delivery, macrosomia and fetal weight. Maternal pre-pregnancy body mass index (BMI) was higher in the COVID-19 GDM cohort ( $p = 0.02$ ).

### Conclusions

Despite the change in guidelines, there was a non-significant increase in the incidence of diagnosis of gestational diabetes.  
(Author)

---

MIDIRS is part of RCM Information Services Limited which is a company incorporated in England and Wales under company no.11914882 with registered office at 10-18 Union Street, London SE1 1SZ  
RCM Information Services Limited is a subsidiary of The Royal College of Midwives



**The Royal College of Midwives Trust**  
10-18 Union Street  
London  
SE1 1SZ



**Open 24 hours a day, 7 days a week**  
T: 0300 303 0444  
F: +44 20 7312 3536  
E: [info@rcm.org.uk](mailto:info@rcm.org.uk)  
W: [www.rcm.org.uk](http://www.rcm.org.uk)



**Chief Executive:**  
Gill Walton,  
MA, PGDip, BSc Hons, RM  
**President:**  
Rebecca Davies, RM  
**Patron:**  
HRH The Princess Royal

**2022-03888**

**Clinical-pathological features in placentas of pregnancies with SARS-CoV-2 infection and adverse outcome: case series with and without congenital transmission.** Zaigham M, Gisselsson D, Sand A, et al (2022), BJOG: An International Journal of Obstetrics and Gynaecology 3 March 2022, online

**Objective**

To correlate clinical outcomes to pathology in SARS-CoV-2 infected placentas in stillborn and live-born infants presenting with fetal distress.

**Design**

Retrospective, observational.

**Setting**

Nationwide.

**Population**

Five stillborn and nine live-born infants from 13 pregnant women infected with SARS-CoV-2 seeking care at seven different maternity units in Sweden.

**Methods**

Clinical outcomes and placental pathology were studied in 14 cases (one twin pregnancy) of maternal SARS-CoV-2 infection with impaired fetal outcome. Outcomes were correlated to placental pathology in order to investigate the impact of virus-related pathology on the villous capillary endothelium, trophoblast and other cells.

**Main outcome measures**

Maternal and fetal clinical outcomes and placental pathology in stillborn and live-born infants.

**Results**

Reduced fetal movements were reported (77%) and time from onset of maternal COVID-19 symptoms to signs of fetal distress among live-born infants was 6 (3–12) days and to diagnosis of stillbirth 11 (2–25) days. Two of the live-born infants died during the postnatal period. Signs of fetal distress led to emergency caesarean section in all live-born infants with umbilical cord blood gases and low Apgar scores confirming intrauterine hypoxia. Five stillborn and one live-born neonate had confirmed congenital transmission. Massive perivillous fibrinoid deposition, intervillitis and trophoblast necrosis were associated with SARS-CoV-2 placental infection and congenital transmission.

**Conclusions**

SARS-CoV-2 can cause rapid placental dysfunction with subsequent acute fetal hypoxia leading to intrauterine fetal compromise. Associated placental pathology included massive perivillous fibrinoid deposition, intervillitis and trophoblast degeneration. (Author)

**Full URL:** <https://doi.org/10.1111/1471-0528.17132>

---

MIDIRS is part of RCM Information Services Limited which is a company incorporated in England and Wales under company no.11914882 with registered office at 10-18 Union Street, London SE1 1SZ  
RCM Information Services Limited is a subsidiary of The Royal College of Midwives



**The Royal College of Midwives Trust**  
10-18 Union Street  
London  
SE1 1SZ



**Open 24 hours a day, 7 days a week**  
T: 0300 303 0444  
F: +44 20 7312 3536  
E: [info@rcm.org.uk](mailto:info@rcm.org.uk)  
W: [www.rcm.org.uk](http://www.rcm.org.uk)



**Chief Executive:**  
Gill Walton,  
MA, PGDip, BSc Hons, RM  
**President:**  
Rebecca Davies, RM  
**Patron:**  
HRH The Princess Royal

**2022-03880**

**Monitored home-based with or without face-to-face exercise for maternal mental health during the COVID-19 pandemic.**

Veisy A, Mohammad-Alizadeh-Charandabi S, Abbas-Alizadeh S, et al (2022), Journal of Reproductive and Infant Psychology 13 April 2022, online

**Introduction**

Despite the known beneficial effects of exercise, most pregnant women do not exercise regularly. Most studies on exercise have been conducted on supervised exercise and there is limited evidence regarding the adherence and effect of other exercise programs on pregnancy outcomes. Therefore, we aimed to investigate adherence to a face-to-face plus monitored home exercise program versus a monitored home-based exercise program on its own during pregnancy. In addition, effects of these two exercise programs on women's mental health during pregnancy and postpartum (primary outcomes) and on some other maternal and neonatal outcomes (secondary outcomes) will be assessed.

**Methods**

In this superiority trial with three parallel arms, 150 women at 12–18 weeks of gestation will be randomised equally into three groups (face-to-face plus monitored home exercise, only monitored home-based exercise, and control). The exercise programs will be performed up to the 38th week of gestation during which participants will be assessed at specific intervals during the pregnancy, and post-partum and followed up until six months after childbirth. The exercise diary will be used to assess the adherence. The Edinburgh Depression Scale and the Positive and Negative Affect Schedule will be used to assess prenatal and postnatal depression and affect, respectively.

**Discussion**

This study reflects the feasibility and acceptance of two exercise programs for pregnant women and their effects on important outcomes. If these programs are followed properly and effectively, pregnant women's health can be improved using these methods at a lower cost compared to the conventional supervised exercise program. (Author)

---

MIDIRS is part of RCM Information Services Limited which is a company incorporated in England and Wales under company no.11914882 with registered office at 10-18 Union Street, London SE1 1SZ  
RCM Information Services Limited is a subsidiary of The Royal College of Midwives



**The Royal College of Midwives Trust**  
10-18 Union Street  
London  
SE1 1SZ



**Open 24 hours a day, 7 days a week**  
T: 0300 303 0444  
F: +44 20 7312 3536  
E: [info@rcm.org.uk](mailto:info@rcm.org.uk)  
W: [www.rcm.org.uk](http://www.rcm.org.uk)



**Chief Executive:**  
Gill Walton,  
MA, PGDip, BSc Hons, RM  
**President:**  
Rebecca Davies, RM  
**Patron:**  
HRH The Princess Royal

**2022-03861**

**Pregnancy and the Risk of In-Hospital Coronavirus Disease 2019 (COVID-19) Mortality.** Pineles BL, Goodman KE, Pineles L, et al (2022), *Obstetrics & Gynecology* vol 139, no 5, May 2022, pp 846-854

**OBJECTIVE:**

To evaluate whether pregnancy is an independent risk factor for in-hospital mortality among patients of reproductive age hospitalized with coronavirus disease 2019 (COVID-19) viral pneumonia.

**METHODS:**

We conducted a retrospective cohort study (April 2020–May 2021) of 23,574 female inpatients aged 15–45 years with an International Classification of Diseases, Tenth Revision, Clinical Modification diagnosis code for COVID-19 discharged from 749 U.S. hospitals in the Premier Healthcare Database. We used a viral pneumonia diagnosis to select for patients with symptomatic COVID-19. The associations between pregnancy and in-hospital mortality, intensive care unit (ICU) admission, and mechanical ventilation were analyzed using propensity score–matched conditional logistic regression. Models were matched for age, marital status, race and ethnicity, Elixhauser comorbidity score, payer, hospital number of beds, season of discharge, hospital region, obesity, hypertension, diabetes mellitus, chronic pulmonary disease, deficiency anemias, depression, hypothyroidism, and liver disease.

**RESULTS:**

In-hospital mortality occurred in 1.1% of pregnant patients and 3.5% of nonpregnant patients hospitalized with COVID-19 and viral pneumonia (propensity score–matched odds ratio [OR] 0.39, 95% CI 0.25–0.63). The frequency of ICU admission for pregnant and nonpregnant patients was 22.0% and 17.7%, respectively (OR 1.34, 95% CI 1.15–1.55). Mechanical ventilation was used in 8.7% of both pregnant and nonpregnant patients (OR 1.05, 95% CI 0.86–1.29). Among patients who were admitted to an ICU, mortality was lower for pregnant compared with nonpregnant patients (OR 0.33, 95% CI 0.20–0.57), though mechanical ventilation rates were similar (35.7% vs 38.3%, OR 0.90, 95% CI 0.70–1.16). Among patients with mechanical ventilation, pregnant patients had a reduced risk of in-hospital mortality compared with nonpregnant patients (0.26, 95% CI 0.15–0.46).

**CONCLUSION:**

Despite a higher frequency of ICU admission, in-hospital mortality was lower among pregnant patients compared with nonpregnant patients with COVID-19 viral pneumonia, and these findings persisted after propensity score matching. (Author)

**Full URL:** <https://doi.org/10.1097/AOG.0000000000004744>

---

MIDIRS is part of RCM Information Services Limited which is a company incorporated in England and Wales under company no.11914882 with registered office at 10-18 Union Street, London SE1 1SZ  
RCM Information Services Limited is a subsidiary of The Royal College of Midwives



**The Royal College of Midwives Trust**  
10-18 Union Street  
London  
SE1 1SZ



**Open 24 hours a day, 7 days a week**  
T: 0300 303 0444  
F: +44 20 7312 3536  
E: [info@rcm.org.uk](mailto:info@rcm.org.uk)  
W: [www.rcm.org.uk](http://www.rcm.org.uk)



**Chief Executive:**  
Gill Walton,  
MA, PGDip, BSc Hons, RM  
**President:**  
Rebecca Davies, RM  
**Patron:**  
HRH The Princess Royal

## 2022-03720

**Association of SARS-CoV-2 Infection During Pregnancy With Maternal and Perinatal Outcomes.** McClymont E, Albert AY, Alton GD, et al (2022), JAMA (Journal of the American Medical Association) vol 327, no 20, 24/31 May 2022, pp 1983-1991

**Importance** There are limited high-quality, population-level data about the effect of SARS-CoV-2 infection on pregnancy using contemporaneous comparator cohorts.

**Objectives** To describe maternal and perinatal outcomes associated with SARS-CoV-2 infection in pregnancy and to assess variables associated with severe disease in the pregnant population.

**Design, Setting, and Participants** CANCOVID-Preg is an observational surveillance program for SARS-CoV-2-affected pregnancies in Canada. This analysis presents exploratory, population-level data from 6 Canadian provinces for the period of March 1, 2020, to October 31, 2021. A total of 6012 pregnant persons with a positive SARS-CoV-2 polymerase chain reaction test result at any time in pregnancy (primarily due to symptomatic presentation) were included and compared with 2 contemporaneous groups including age-matched female individuals with SARS-CoV-2 and unaffected pregnant persons from the pandemic time period.

**Exposure** SARS-CoV-2 infection during pregnancy. Incident infections in pregnancy were reported to CANCOVID-Preg by participating provinces/territories.

**Main Outcomes and Measures** Maternal and perinatal outcomes associated with SARS-CoV-2 infection as well as risk factors for severe disease (ie, disease requiring hospitalization, admission to an intensive care unit/critical care unit, and/or oxygen therapy).

**Results** Among 6012 pregnant individuals with SARS-CoV-2 in Canada (median age, 31 [IQR, 28-35] years), the greatest proportion of cases were diagnosed at 28 to 37 weeks' gestation (35.7%). Non-White individuals were disproportionately represented. Being pregnant was associated with a significantly increased risk of SARS-CoV-2-related hospitalization compared with SARS-CoV-2 cases among all women aged 20 to 49 years in the general population of Canada (7.75% vs 2.93%; relative risk, 2.65 [95% CI, 2.41-2.88]) as well as an increased risk of intensive care unit/critical care unit admission (2.01% vs 0.37%; relative risk, 5.46 [95% CI, 4.50-6.53]). Increasing age, preexisting hypertension, and greater gestational age at diagnosis were significantly associated with worse maternal outcomes. The risk of preterm birth was significantly elevated among SARS-CoV-2-affected pregnancies (11.05% vs 6.76%; relative risk, 1.63 [95% CI, 1.52-1.76]), even in cases of milder disease not requiring hospitalization, compared with unaffected pregnancies during the same time period.

**Conclusions and Relevance** In this exploratory surveillance study conducted in Canada from March 2020 to October 2021, SARS-CoV-2 infection during pregnancy was significantly associated with increased risk of adverse maternal outcomes and preterm birth. (Author)

**Full URL:** <https://doi.org/10.1001/jama.2022.5906>

---

## 2022-03586

**A case of temporary anhydramnios after COVID-19 infection.** Kasuga Y, Sou Y, Fukuoka M, et al (2022), European Journal of Obstetrics & Gynecology and Reproductive Biology vol 272, May 2022, pp 255-256

Herein, we describe a case of temporary anhydramnios after a COVID-19 infection. (Author, edited)

**Full URL:** <https://doi.org/10.1016/j.ejogrb.2022.04.002>

---

MIDIRS is part of RCM Information Services Limited which is a company incorporated in England and Wales under company no.11914882 with registered office at 10-18 Union Street, London SE1 1SZ  
RCM Information Services Limited is a subsidiary of The Royal College of Midwives



**The Royal College of Midwives Trust**  
10-18 Union Street  
London  
SE1 1SZ



**Open 24 hours a day, 7 days a week**  
T: 0300 303 0444  
F: +44 20 7312 3536  
E: [info@rcm.org.uk](mailto:info@rcm.org.uk)  
W: [www.rcm.org.uk](http://www.rcm.org.uk)



**Chief Executive:**  
Gill Walton,  
MA, PGDip, BSc Hons, RM  
**President:**  
Rebecca Davies, RM  
**Patron:**  
HRH The Princess Royal

## 2022-03581

**“Never let a good crisis go to waste”: Positives from disrupted maternity care in Australia during COVID-19.** Kluwgant D, Homer C, Dahlen H (2022), *Midwifery* vol 110, July 2022, 103340

### Objective

Due to the COVID-19 pandemic, a number of changes to maternity care were rapidly introduced in all countries, including Australia, to reduce the risk of infection for pregnant women and their care providers. While many studies have reported on the negative effects of these changes, there is a paucity of evidence on factors which women and their providers perceived as positive and useful for future maternity care.

### Design

Data was analysed from the Birth in the time of COVID-19 (BITTOC 2020) study survey. Conventional content analysis and descriptive statistics were used to analyse the data and examine which aspects of COVID-amended care women experienced as positive. Data from women were compared to data from midwives.

### Setting

This project took place in Australia in 2020-2021.

### Participants

The survey was distributed to women who gave birth and midwives who worked in Australia during the COVID-19 pandemic (March 2020 onwards).

### Measurements and findings

Women reported a variety of positives from their maternity care during COVID-19. These included both care-related factors as well as contextual factors. The most commonly mentioned positives for pregnant and postnatal women were care-related, namely fewer visitors in hospital, having increased access to telehealth services. These were also the most commonly reported positives by midwives. Having midwifery continuity of care models, giving birth at home and having their partner work from home were also highlighted by women as positives.

### Key conclusions

Despite the negative effect of COVID-19-related restrictions on maternity care, a variety of changes were viewed as positive by both women and midwives, with strong agreement between the two groups.

### Implications for practice

These findings provide evidence to support the inclusion of these positive elements of care and ensure that the lessons learned from the pandemic are utilised to improve maternity care in Australia going forward. (Author)

**Full URL:** <https://doi.org/10.1016/j.midw.2022.103340>

---

## 2022-03575

**Trends in Maternal Outcomes During the COVID-19 Pandemic in Alabama From 2016 to 2021.** Shukla VV, Rahman F, Shen X, et al (2022), *JAMA Network Open* vol 5, no 4, April 2022, e222681

This cohort study assesses whether the COVID-19 pandemic is associated with an increase in the risk of maternal morbidity and mortality in Alabama from 2016 to 2021. (Author)

**Full URL:** <https://doi.org/10.1001/jamanetworkopen.2022.2681>

---

MIDIRS is part of RCM Information Services Limited which is a company incorporated in England and Wales under company no.11914882 with registered office at 10-18 Union Street, London SE1 1SZ  
RCM Information Services Limited is a subsidiary of The Royal College of Midwives



**The Royal College of Midwives Trust**  
10-18 Union Street  
London  
SE1 1SZ



**Open 24 hours a day, 7 days a week**  
T: 0300 303 0444  
F: +44 20 7312 3536  
E: [info@rcm.org.uk](mailto:info@rcm.org.uk)  
W: [www.rcm.org.uk](http://www.rcm.org.uk)



**Chief Executive:**  
Gill Walton,  
MA, PGDip, BSc Hons, RM  
**President:**  
Rebecca Davies, RM  
**Patron:**  
HRH The Princess Royal



2022-03573

**Prevalence of postpartum depression and antenatal anxiety symptoms during COVID-19 pandemic: An observational prospective cohort study in Greece.** Micha G, Hyphantis T, Staikou C, et al (2022), European Journal of Midwifery vol 6, April 2022, p 23

**Introduction:**

A significant proportion of pregnant women and women in the early postpartum period suffer from mental health problems. The COVID-19 pandemic represents a unique stressor during this period and many studies across the world have shown elevated rates of postpartum depression (PPD).

**Methods:**

In this multicenter two-phase observational prospective cohort study, we aim to assess the prevalence of anxiety prior to labor (Generalized Anxiety Disorder-7), as well as PPD at 6–8 weeks postpartum using the Edinburgh Postnatal Depression Scale (EPDS).

**Results:**

Of the 330 women analyzed, 13.2% reported symptoms of depression using EPDS cut-off score  $\geq 13$ . High antenatal levels of anxiety (24.8% scored  $\geq 10$  in GAD-7) were documented. A significant proportion of postpartum women reported a decrease in willingness to attend antenatal education courses (36%) and fewer antenatal visits to their obstetrician (34%) due to pandemic. Higher antenatal anxiety increased the odds of being depressed at 6–8 weeks postpartum (EPDS  $\geq 13$ ).

**Conclusions:**

Compared to reported prevalence of PPD from previous studies before the COVID-19 era in Greece, we did not find elevated rates during the first wave of the pandemic. High anxiety levels were observed indicating that there is a need for close monitoring in pregnancy during the pandemic and anxiety screening to identify women who need support in the pandemic era. A well-planned maternity program should be employed by all the associated care providers to maintain the proper antenatal care adjusted to the pandemic strains as well as a follow-up after labor. (Author)

**Full URL:** <https://doi.org/10.18332/ejm/146233>

---

MIDIRS is part of RCM Information Services Limited which is a company incorporated in England and Wales under company no.11914882 with registered office at 10-18 Union Street, London SE1 1SZ  
RCM Information Services Limited is a subsidiary of The Royal College of Midwives



**The Royal College of Midwives Trust**  
10-18 Union Street  
London  
SE1 1SZ



**Open 24 hours a day, 7 days a week**  
T: 0300 303 0444  
F: +44 20 7312 3536  
E: [info@rcm.org.uk](mailto:info@rcm.org.uk)  
W: [www.rcm.org.uk](http://www.rcm.org.uk)



**Chief Executive:**  
Gill Walton,  
MA, PGDip, BSc Hons, RM  
**President:**  
Rebecca Davies, RM  
**Patron:**  
HRH The Princess Royal

**2022-03458**

**Systematic review and critical evaluation of quality of clinical practice guidelines on the management of SARS-CoV-2 infection in pregnancy.** Di Girolamo R, Khalil A, Rizzo G, et al (2022), American Journal of Obstetrics & Gynecology MFM 2 May 2022, online. 100654

Objective: To systematically identify and critically assess the quality of clinical practice guidelines (CPGs) for the management of SARS-CoV-2 infection in pregnancy.

Data Source: Medline, Scopus and ISI Web of Science databases were searched until 15 th of February.

Study eligibility criteria: Inclusion criteria were CPGs on the management of SARS-CoV-2 infection in pregnancy. The risk of bias and quality assessment of the included CPGs were performed using "The Appraisal of Guidelines for REsearch and Evaluation (AGREE II)" tool, which is considered as the gold standard for CPG quality assessment. To define a CPG as of good quality we adopted the cut-off score according to Amer et al.: if the overall guideline score was >60%, CPGs was recommended.

Study appraisal and synthesis methods: The following clinical points related to the management of pregnant women with SARS-CoV-2 infection were addressed: criteria for maternal hospitalization, recommendations for follow-up fetal growth scan, specific recommendations against invasive procedures, management of labor, timing of delivery, postpartum care and vaccination strategy.

Results: Twenty-eight CPGs were included. All of them recommended hospitalization only for severe disease. Forty-six percent (6/13) of CPGs suggested a fetal growth scan after SARS-COV-2 infection while 23.1% (3/13) did not support this practice. Thromboprophylaxis with low molecular weight heparin (LMWH) was recommended in symptomatic women by 77.1% (7/9) of the CPGs. None of the CPGs recommended to administer corticosteroids only for the presence of SARS-CoV-2 infection in preterm gestation, unless specific obstetric indication exists. Elective induction of labor from 39 weeks of gestation was suggested by 18.1% (2/11) of the CPGs included in the present review, while 45.4% (5/11) did not recommend elective induction unless other obstetric indications co-existed. Twenty-seven percent (3/11) of the CPGs suggested shortening of the second stage of labor and active pushing was supported by 18.1% (2/11) of them. A general agreement was found among the CPGs in not recommending Cesarean Section (CS) only for the presence of maternal infection and recommending vaccine booster at least 6 months after the primary series of vaccination. The AGREE II standardized domain scores for the first overall assessment (OA1) of CPGs had a mean of 50% (SD±21.82%) and 9 CPGs scored more than 60%.

Conclusions: A significant heterogeneity was found in some major aspects of the main aspects of the management of SARS-CoV-2 infection in pregnancy reported by the published CPGs. (Author)

Full URL: <https://doi.org/10.1016/j.ajogmf.2022.100654>

MIDIRS is part of RCM Information Services Limited which is a company incorporated in England and Wales under company no.11914882 with registered office at 10-18 Union Street, London SE1 1SZ  
RCM Information Services Limited is a subsidiary of The Royal College of Midwives



**The Royal College of Midwives Trust**  
10-18 Union Street  
London  
SE1 1SZ



**Open 24 hours a day, 7 days a week**  
T: 0300 303 0444  
F: +44 20 7312 3536  
E: [info@rcm.org.uk](mailto:info@rcm.org.uk)  
W: [www.rcm.org.uk](http://www.rcm.org.uk)



**Chief Executive:**  
Gill Walton,  
MA, PGDip, BSc Hons, RM  
**President:**  
Rebecca Davies, RM  
**Patron:**  
HRH The Princess Royal

**2022-03408**

**Impact of SARS-CoV-2 infection on risk of prematurity, birthweight and obstetric complications: A multivariate analysis from a nationwide, population-based retrospective cohort study.** Simon E, Gouyon J-B, Cottenet J, et al (2022), BJOG: An International Journal of Obstetrics and Gynaecology vol 129, no 7, June 2022, pp 1084-1094

**Objective**

To determine the impact of maternal coronavirus disease 2019 (COVID-19) on prematurity, birthweight and obstetric complications.

**Design**

Nationwide, population-based retrospective cohort study.

**Setting**

National Programme de Médicalisation des Systèmes d'Information database in France.

**Population**

All single births from March to December 2020: 510 387 deliveries, including 2927 (0.6%) with confirmed COVID-19 in the mother and/or the newborn.

**Methods**

The group with COVID-19 was compared with the group without COVID-19 using the chi-square test or Fisher's exact test, and the Student's t test or Mann–Whitney U test. Logistic regressions were used to study the effect of COVID-19 on the risk of prematurity or macrosomia (birthweight  $\geq 4500$  g).

**Main outcome measures**

Prematurity less than 37, less than 28, 28–31, or 32–36 weeks of gestation; birthweight; obstetric complications.

**Results**

In singleton pregnancies, COVID-19 was associated with obstetric complications such as hypertension (2.8% versus 2.0%,  $p < 0.01$ ), pre-eclampsia (3.6% versus 2.0%,  $p < 0.01$ ), diabetes (18.8% versus 14.4%,  $p < 0.01$ ) and caesarean delivery (26.8% versus 19.7%,  $p < 0.01$ ). Among pregnant women with COVID-19, there was more prematurity between 28 and 31 weeks of gestation (1.3% versus 0.6%,  $p < 0.01$ ) and between 32 and 36 weeks of gestation (7.7% versus 4.3%,  $p < 0.01$ ), and more macrosomia (1.0% versus 0.7%,  $p = 0.04$ ), but there was no difference in small-for-gestational-age newborns (6.3% versus 8.7%,  $p = 0.15$ ). Logistic regression analysis for prematurity showed an adjusted odds ratio (aOR) of 1.77 (95% CI 1.55–2.01) for COVID-19. For macrosomia, COVID-19 resulted in non-significant aOR of 1.38 (95% CI 0.95–2.00).

**Conclusions**

COVID-19 is a risk factor for prematurity, even after adjustment for other risk factors. (Author)

**Full URL:** <https://doi.org/10.1111/1471-0528.17135>

---

MIDIRS is part of RCM Information Services Limited which is a company incorporated in England and Wales under company no.11914882 with registered office at 10-18 Union Street, London SE1 1SZ  
RCM Information Services Limited is a subsidiary of The Royal College of Midwives



**The Royal College of Midwives Trust**  
10-18 Union Street  
London  
SE1 1SZ



**Open 24 hours a day, 7 days a week**  
T: 0300 303 0444  
F: +44 20 7312 3536  
E: [info@rcm.org.uk](mailto:info@rcm.org.uk)  
W: [www.rcm.org.uk](http://www.rcm.org.uk)



**Chief Executive:**  
Gill Walton,  
MA, PGDip, BSc Hons, RM  
**President:**  
Rebecca Davies, RM  
**Patron:**  
HRH The Princess Royal

**2022-03396**

**Knowledge, attitudes, and practices related to the COVID-19 pandemic among pregnant women in Bangkok, Thailand.** Kunno J, Yubonpant P, Supawattanbodee B, et al (2022), BMC Pregnancy and Childbirth vol 22, no 357, 23 April 2022

#### Background

Pregnancy is associated with increased risk for severe COVID-19. Few studies have examined knowledge, attitudes, and practices (KAP) related to pregnancy during the pandemic. This study investigated the association between socio-demographic characteristics and KAP related to COVID-19 among pregnant women in an urban community in Thailand.

#### Methods

A cross-sectional online survey was distributed among pregnant women in Bangkok, Thailand from July–August 2021. Binary logistic regression was conducted to test the association between socio-demographic characteristics and KAP related to COVID-19, and a Spearman’s analysis tested correlations between KAP scores.

#### Results

A total of 150 pregnancy survey responses were received. Most participants were third trimester (27–40 weeks gestation; 68.0%). Pregnancy had never been risked contracting COVID-19 (84.7%). Most expressed concerns about being infected with COVID-19 during pregnancy and following birth (94.0 and 70.0%, respectively). The results of binary logistic regression analysis found associations between knowledge and marital status (OR = 4.983, 95%CI 1.894–13.107). In addition, having a bachelor’s degree or higher was associated with higher attitude scores (OR = 2.733, 95%CI 1.045–7.149), as was being aged 26–30 (OR = 2.413 95%CI 0.882–6.602) and 31–35 years of age (OR = 2.518–2.664, 95%CI 0.841–8.442). Higher practice scores were associated with having a bachelor’s degree or higher (OR = 2.285 95%CI 1.110–6.146), and income  $\geq$ 15,001 bath (OR = 4.747 95%CI 1.588–14.192). Correlation analysis found a weak positive correlation between knowledge and practice scores ( $r = 0.210$ ,  $p$ -value = 0.01).

#### Conclusion

Participants overall had high KAP scores. This study can guide public health strategies regarding pregnant women and COVID-19. We recommend that interventions to improve and attitude and practice scores. Knowledge on pregnancy and COVID-19 should focus on reducing fear and improving attitudes toward the care of patients as well as the promotion of preventive practices. (Author)

**Full URL:** <https://doi.org/10.1186/s12884-022-04612-3>

---

MIDIRS is part of RCM Information Services Limited which is a company incorporated in England and Wales under company no.11914882 with registered office at 10-18 Union Street, London SE1 1SZ  
RCM Information Services Limited is a subsidiary of The Royal College of Midwives



**The Royal College of Midwives Trust**  
10-18 Union Street  
London  
SE1 1SZ



**Open 24 hours a day, 7 days a week**  
T: 0300 303 0444  
F: +44 20 7312 3536  
E: [info@rcm.org.uk](mailto:info@rcm.org.uk)  
W: [www.rcm.org.uk](http://www.rcm.org.uk)



**Chief Executive:**  
Gill Walton,  
MA, PGDip, BSc Hons, RM  
**President:**  
Rebecca Davies, RM  
**Patron:**  
HRH The Princess Royal

## 2022-03369

**Case series of COVID-19 infection in pregnancy complicated by ketoacidosis and symptomatic breathlessness.** Thorne I, Steele S, Martineau M, et al (2022), *Obstetric Medicine* vol 15, no 1, March 2022, pp 50-53

### Background

The differential diagnosis of acute shortness of breath in a pregnant woman with COVID-19 is broad. Pregnancy is a ketosis-prone state, which can result in metabolic acidosis and tachypnoea.

### Methods

We describe four pregnant women with COVID-19 and breathlessness where ketoacidosis was found to contribute to symptomatic tachypnoea.

### Results

One patient did not have associated COVID-19 pneumonitis, but presented with severe tachypnoea and metabolic acidosis; three women had pneumonitis and metabolic acidosis. Corrective treatment for the metabolic abnormalities resulted in resolution of the ketoacidosis in all cases. No women had coexistent diabetes.

### Conclusion

This is the first series of COVID-19 in pregnancy complicated by ketoacidosis and symptomatic tachypnoea. Ketoacidosis associated with COVID-19 is an important cause of tachypnoea requiring specific treatment, which should not be overlooked. Potential mechanisms for this are discussed with a framework for interpretation of blood gas results during pregnancy. (Author)

Full URL: <https://doi.org/10.1177%2F1753495X211024511>

---

## 2022-03361

**Considerations for women with COVID-19 admitted to hospital.** Coad F, Frise C (2022), *Obstetric Medicine* 16 March 2022, online

The number of pregnant women being admitted with severe COVID-19 infection and dying has increased with each wave of the pandemic. These women often present unique challenges to the medical and obstetric teams given the changes in physiology that occur in pregnancy, affecting assessment and management, as well as the practical difficulties such as the ideal location of care. Whilst the basis of treatment remains the same, there are nuances to caring for pregnant women that need considerable thought and multidisciplinary collaboration. Obstetricians, neonatologists, midwives, intensivists, anaesthetists and physicians may all be involved at some point, depending on the gestation and severity of illness. Implementing a COVID-19 in pregnancy guideline or checklist for your hospital will help ensure pregnant women are managed in a safe and timely manner. Here described are some key recommendations to help in the management of pregnant women admitted with COVID-19. (Author)

---

## 2022-03335

**COVID-19: clinical presentation and implications. A primer for obstetricians.** Dahan MH, Steiner N (2022), *Journal of Maternal-Fetal & Neonatal Medicine* vol 35, no 12, 2022, pp 2424-2426

Although, many obstetricians (OBs) are fighting the war on COVID-19 as we speak, others have not yet had contact with these patients, practicing in areas where the infection rates are low and therefore, the clinical presentation remains unknown. This article was developed to shed light on this enigma. It is based on published studies and physician experience treating these patients. (Author)

Full URL: <https://doi.org/10.1080/14767058.2020.1777274>

---

MIDIRS is part of RCM Information Services Limited which is a company incorporated in England and Wales under company no.11914882 with registered office at 10-18 Union Street, London SE1 1SZ  
RCM Information Services Limited is a subsidiary of The Royal College of Midwives



**The Royal College of Midwives Trust**  
10-18 Union Street  
London  
SE1 1SZ



**Open 24 hours a day, 7 days a week**  
T: 0300 303 0444  
F: +44 20 7312 3536  
E: [info@rcm.org.uk](mailto:info@rcm.org.uk)  
W: [www.rcm.org.uk](http://www.rcm.org.uk)



**Chief Executive:**  
Gill Walton,  
MA, PGDip, BSc Hons, RM  
**President:**  
Rebecca Davies, RM  
**Patron:**  
HRH The Princess Royal

**2022-03321**

**Removal of Covid-19 guidance on protecting pregnant nurses branded 'reckless'.** Ford M (2022), Nursing Times 22 April 2022

Health unions and organisations have raised serious concerns about the government's "reckless" removal of Covid-19 guidance on how pregnant staff should be protected in the workplace. (Author)

---

**2022-03294**

**Diagnosis and management of covid-19 in pregnancy.** Nana M, Hodson K, Lucas N, et al (2022), BMJ vol 377, 26 April 2022, e069739

Pregnant women with covid-19 are at greater risk of severe disease than their non-pregnant peers, and yet they are frequently denied investigations or treatments because of unfounded concerns about risk to the fetus. The basic principles of diagnosing and managing covid-19 are the same as for non-pregnant patients, and a multidisciplinary, expert team approach is essential to ensure optimal care. During pregnancy, treatment with corticosteroids should be modified to use non-fluorinated glucocorticoids. Il-6 inhibitors and monoclonal antibodies, together with specific antiviral therapies, may also be considered. Prophylaxis against venous thromboembolism is important. Women may require respiratory support with oxygen, non-invasive ventilation, ventilation in a prone position (either awake or during invasive ventilation), intubation and ventilation, and extracorporeal membrane oxygenation (ECMO). Pregnancy is not a contraindication for any of these supportive therapies, and the criteria for providing them are the same as in the general population. Decisions regarding timing, place, and mode of delivery should be taken with a multidisciplinary team including obstetricians, physicians, anesthetists, and intensivists experienced in the care of covid-19 in pregnancy. Ideally these decisions should take place in consultation with centers that have experience and expertise in all these specialties. (Author)

**Full URL:** <https://doi.org/10.1136/bmj-2021-069739>

---

MIDIRS is part of RCM Information Services Limited which is a company incorporated in England and Wales under company no.11914882 with registered office at 10-18 Union Street, London SE1 1SZ  
RCM Information Services Limited is a subsidiary of The Royal College of Midwives



**The Royal College of Midwives Trust**  
10-18 Union Street  
London  
SE1 1SZ



**Open 24 hours a day, 7 days a week**  
T: 0300 303 0444  
F: +44 20 7312 3536  
E: [info@rcm.org.uk](mailto:info@rcm.org.uk)  
W: [www.rcm.org.uk](http://www.rcm.org.uk)



**Chief Executive:**  
Gill Walton,  
MA, PGDip, BSc Hons, RM  
**President:**  
Rebecca Davies, RM  
**Patron:**  
HRH The Princess Royal

**2022-03106**

**Maternal and neonatal outcomes of pregnancies with COVID-19 after medically assisted reproduction – results from the prospective Covid-19-Related Obstetric and Neonatal Outcome Study (CRONOS).** Ziert Y, Abou-Dakn M, Backes C, et al (2022), American Journal of Obstetrics & Gynecology (AJOG) 19 April 2022, online

#### Background

Severe acute respiratory syndrome coronavirus type 2 infections in pregnancy have been associated with maternal morbidity, admission to intensive care, and adverse perinatal outcomes such as preterm birth, stillbirth and hypertensive disorders of pregnancy. It is unclear whether in women with COVID-19 medically assisted reproduction additionally affects maternal and neonatal outcomes.

#### Objective

To evaluate the effect of medically assisted reproduction on maternal and neonatal outcomes of women with COVID-19 in pregnancy.

#### Study design

A total of 1,485 women with COVID-19 registered in the Covid-19 Related Obstetric and Neonatal Outcome Study, a multicentric prospective observational cohort study, were included. Maternal and neonatal outcomes of 65 pregnancies achieved with medically assisted reproduction and 1,420 spontaneously conceived pregnancies were compared. We used univariate and multivariate (multinomial) logistic regressions to estimate (un)adjusted odds ratios and 95% confidence intervals for adverse outcomes.

#### Results

Compared to women after spontaneous conceptions with COVID-19, the incidence of COVID-19 associated adverse outcomes (e.g. pneumonia, admission to intensive care, death) was not different than in women after medically assisted reproduction pregnancies. Yet, the risk of obstetric and neonatal complications was higher in pregnancies achieved through medically assisted reproduction. However, medically assisted reproduction was not the primary risk factor for adverse maternal and neonatal outcomes, e.g. pregnancy-related hypertensive disorders, gestational diabetes mellitus, cervical insufficiency, peripartum hemorrhage, delivery by caesarean section, preterm birth or admission to neonatal intensive care. Maternal age, multiple pregnancies, nulliparity, BMI >30 (before pregnancy) and multiple gestation contributed differently to the increased risks of adverse pregnancy outcomes in women with COVID-19, independent of medically assisted reproduction.

#### Conclusion

Although women with COVID-19 who conceived through fertility treatment experienced a higher incidence of adverse obstetric and neonatal complications than women with spontaneous conceptions, medically assisted reproduction was not the primary risk factor. (Author)

---

MIDIRS is part of RCM Information Services Limited which is a company incorporated in England and Wales under company no.11914882 with registered office at 10-18 Union Street, London SE1 1SZ  
RCM Information Services Limited is a subsidiary of The Royal College of Midwives



**The Royal College of Midwives Trust**  
10-18 Union Street  
London  
SE1 1SZ



**Open 24 hours a day, 7 days a week**  
T: 0300 303 0444  
F: +44 20 7312 3536  
E: [info@rcm.org.uk](mailto:info@rcm.org.uk)  
W: [www.rcm.org.uk](http://www.rcm.org.uk)



**Chief Executive:**  
Gill Walton,  
MA, PGDip, BSc Hons, RM  
**President:**  
Rebecca Davies, RM  
**Patron:**  
HRH The Princess Royal

## 2022-03104

**Increase in preterm stillbirths in association with reduction in iatrogenic preterm births during COVID-19 lockdown in Australia: a multi-centre cohort study.** Hui L, Marzan MB, Potenza S, et al (2022), American Journal of Obstetrics & Gynecology (AJOG) 19 April 2022, online

### Background

The COVID-19 pandemic has been associated with a worsening of perinatal outcomes in many regions. Melbourne, Australia, had one of the longest and most stringent lockdowns in the world in 2020, while recording only rare instances of COVID-19 infection in pregnant women.

### Objective

The aim of this study was to compare stillbirth and preterm birth rates in women who were exposed or unexposed to lockdown restrictions during pregnancy.

### Study design

Retrospective multi-centre cohort study of perinatal outcomes in Melbourne before and during COVID-19 lockdown. Lockdown period was defined as 23 March 2020 to 14 March 2021. Routinely-collected maternity data from all 12 public hospitals in Melbourne were obtained on singleton pregnancies > 24 weeks gestation without congenital anomalies. We defined the lockdown-exposed cohort as those women for whom weeks 20-40 of gestation occurred during lockdown, and the unexposed control group as women from the corresponding calendar periods 12 and 24 months prior. The main outcome measures were: stillbirth, preterm birth, fetal growth restriction (birth weight < 3rd centile), and iatrogenic preterm birth for fetal compromise. We performed multivariable logistic regression analysis to compare the odds of stillbirth, preterm birth, fetal growth restriction, and iatrogenic preterm birth for fetal compromise, adjusting for multiple covariates.

### Results

There were 24,817 births in the exposed and 50,017 births in the control group. There was a significantly higher risk of preterm stillbirth in the exposed group compared with the control group (0.26% vs 0.18%, aOR 1.49, 95%CI 1.08-2.05, P = 0.015). There was also a significant reduction in preterm birth of live infants < 37 weeks (5.68% vs 6.07%, aOR 0.93, 95%CI 0.87-0.99, P = 0.02), largely mediated by a significant reduction in iatrogenic preterm birth (3.01% vs 3.27%, aOR 0.91, 95%CI 0.83-0.99, P = 0.03), including iatrogenic preterm birth for fetal compromise (1.25% vs 1.51%, aOR 0.82, 95%CI 0.71-0.93, P = 0.003). There were also significant reductions in special care nursery admissions during lockdown (11.53% vs 12.51%, aOR 0.90, 95%CI 0.86-0.95, P < 0.0001). There was a trend to fewer spontaneous preterm births < 37 weeks in the exposed group of a similar magnitude to that reported in other countries (2.69% vs 2.82%, aOR 0.95, 95%CI 0.87-1.05, P = 0.32).

### Conclusions

Lockdown restrictions in Melbourne, Australia were associated with a significant reduction in iatrogenic preterm birth for fetal compromise and a significant increase in preterm stillbirths. This raises concerns that pandemic conditions in 2020 may have led to a failure to identify and appropriately care for pregnant women at increased risk of antepartum stillbirth. Further research is required to understand the relationship between these two findings and to inform our ongoing responses to the pandemic. (Author)

---

MIDIRS is part of RCM Information Services Limited which is a company incorporated in England and Wales under company no.11914882 with registered office at 10-18 Union Street, London SE1 1SZ  
RCM Information Services Limited is a subsidiary of The Royal College of Midwives



**The Royal College of Midwives Trust**  
10-18 Union Street  
London  
SE1 1SZ



**Open 24 hours a day, 7 days a week**  
T: 0300 303 0444  
F: +44 20 7312 3536  
E: [info@rcm.org.uk](mailto:info@rcm.org.uk)  
W: [www.rcm.org.uk](http://www.rcm.org.uk)



**Chief Executive:**  
Gill Walton,  
MA, PGDip, BSc Hons, RM  
**President:**  
Rebecca Davies, RM  
**Patron:**  
HRH The Princess Royal



**2022-03089**

**Vitamin D may prevent COVID-19 induced pregnancy complication.** Al-Kaleel A, Al-Gailani L, Demir M, et al (2022), Medical Hypotheses vol 158, January 2022, 110733

SARS-CoV-2 enters target cells via the ACE2 receptor and downregulates it. ACE2 exhibits high catalytic activity to produce Angiotensin 1–7 (Ang-1–7), which has a vasodilator effect and also inactivates the vasoconstrictor Angiotensin II. In normal pregnancy ACE2 expression is raising in the uterus and placenta. Ang-1–7 levels in plasma are significantly higher in third-trimester pregnant women when compared to non-pregnant women. This may be contributing to systemic vasodilation and reduced blood pressure and modulating hemodynamics during pregnancy. Interestingly, Ang-1–7 plasma levels are lower in pregnancies complicated by pre-eclampsia than normal pregnancies. COVID-19 infection increased the inflammatory cytokines and reduced ACE2 level. This may lead to pre-eclampsia or hypertensive pregnancies, then increasing the perinatal and maternal mortality and morbidity. Vitamin D increased ACE2 expression and Ang-1–7 plasma levels and also decreased Ang II level in plasma. Moreover, Vitamin D reduced the inflammatory cytokine storm. So, Vitamin D supplementation can prevent the risk of preeclampsia or hypertension in pregnant women with COVID-19. (Author)

**Full URL:** <https://doi.org/10.1016/j.mehy.2021.110733>

---

MIDIRS is part of RCM Information Services Limited which is a company incorporated in England and Wales under company no.11914882 with registered office at 10-18 Union Street, London SE1 1SZ  
RCM Information Services Limited is a subsidiary of The Royal College of Midwives



**The Royal College of Midwives Trust**  
10-18 Union Street  
London  
SE1 1SZ



**Open 24 hours a day, 7 days a week**  
T: 0300 303 0444  
F: +44 20 7312 3536  
E: [info@rcm.org.uk](mailto:info@rcm.org.uk)  
W: [www.rcm.org.uk](http://www.rcm.org.uk)



**Chief Executive:**  
Gill Walton,  
MA, PGDip, BSc Hons, RM  
**President:**  
Rebecca Davies, RM  
**Patron:**  
HRH The Princess Royal

## 2022-03041

**Differences and similarities in endothelial and angiogenic profiles of preeclampsia and COVID-19 in pregnancy.** Palomo M, Youssef L, Ramos A, et al (2022), American Journal of Obstetrics & Gynecology (AJOG) 26 March 2022, online

### Background

COVID-19 presents a spectrum of signs and symptoms in pregnant women that might resemble preeclampsia. Differentiation between severe COVID-19 and preeclampsia is difficult in some cases.

### Objective

To study biomarkers of endothelial damage, coagulation, innate immune response, and angiogenesis in preeclampsia and COVID-19 in pregnancy in addition to in vitro alterations in endothelial cells exposed to sera from pregnant women with preeclampsia and COVID-19.

### Study Design

Plasma and sera samples were obtained from pregnant women with COVID-19 infection classified into mild (n=10) or severe (n=9) and from women with normotensive pregnancies as controls (n=10) and patients with preeclampsia (n=13). A panel of plasmatic biomarkers was assessed, including vascular cell adhesion molecule-1, soluble tumor necrosis factor-receptor I, heparan sulfate, von Willebrand factor antigen (activity and multimeric pattern),  $\alpha$ 2-antiplasmin, C5b9, neutrophil extracellular traps, placental growth factor, soluble fms-like tyrosine kinase-1, and angiopoietin 2. In addition, microvascular endothelial cells were exposed to patients' sera, and changes in the cell expression of intercellular adhesion molecule 1 on cell membranes and von Willebrand factor release to the extracellular matrix were evaluated through immunofluorescence. Changes in inflammation cell signaling pathways were also assessed by of p38 mitogen-activated protein kinase phosphorylation. Statistical analysis included univariate and multivariate methods.

### Results

Biomarker profiles of patients with mild COVID-19 were similar to those of controls. Both preeclampsia and severe COVID-19 showed significant alterations in most circulating biomarkers with distinctive profiles. Whereas severe COVID-19 exhibited higher concentrations of vascular cell adhesion molecule-1, soluble tumor necrosis factor- $\alpha$  receptor I, heparan sulfate, von Willebrand factor antigen, and neutrophil extracellular traps, with a significant reduction of placental growth factor compared with controls, preeclampsia presented a marked increase in vascular cell adhesion molecule-1 and soluble tumor necrosis factor- $\alpha$  receptor I (significantly increased compared with controls and patients with severe COVID-19), with a striking reduction in von Willebrand factor antigen, von Willebrand factor activity, and  $\alpha$ 2-antiplasmin. As expected, reduced placental growth factor, increased soluble fms-like tyrosine kinase-1 and angiopoietin 2, and a very high soluble fms-like tyrosine kinase-1 to placental growth factor ratio were also observed in preeclampsia. In addition, a significant increase in C5b9 and neutrophil extracellular traps was also detected in preeclampsia compared with controls. Principal component analysis demonstrated a clear separation between patients with preeclampsia and the other groups (first and second components explained 42.2% and 13.5% of the variance), mainly differentiated by variables related to von Willebrand factor, soluble tumor necrosis factor-receptor I, heparan sulfate, and soluble fms-like tyrosine kinase-1. Von Willebrand factor multimeric analysis revealed the absence of von Willebrand factor high-molecular-weight multimers in preeclampsia (similar profile to von Willebrand disease type 2A), whereas in healthy pregnancies and COVID-19 patients, von Willebrand factor multimeric pattern was normal. Sera from both preeclampsia and severe COVID-19 patients induced an overexpression of intercellular adhesion molecule 1 and von Willebrand factor in endothelial cells in culture compared with controls. However, the effect of preeclampsia was less pronounced than the that of severe COVID-19. Immunoblots of lysates from endothelial cells exposed to mild and severe COVID-19 and preeclampsia sera showed an increase in p38 mitogen-activated protein kinase phosphorylation. Patients with severe COVID-19 and preeclampsia were statistically different from controls, suggesting that both severe COVID-19 and preeclampsia sera can activate inflammatory signaling pathways.

### Conclusion

Although similar in in vitro endothelial dysfunction, preeclampsia and severe COVID-19 exhibit distinctive profiles of circulating biomarkers related to endothelial damage, coagulopathy, and angiogenic imbalance that could aid in the

MIDIRS is part of RCM Information Services Limited which is a company incorporated in England and Wales under company no.11914882 with registered office at 10-18 Union Street, London SE1 1SZ  
RCM Information Services Limited is a subsidiary of The Royal College of Midwives



**The Royal College of Midwives Trust**  
10-18 Union Street  
London  
SE1 1SZ



**Open 24 hours a day, 7 days a week**  
T: 0300 303 0444  
F: +44 20 7312 3536  
E: [info@rcm.org.uk](mailto:info@rcm.org.uk)  
W: [www.rcm.org.uk](http://www.rcm.org.uk)



**Chief Executive:**  
Gill Walton,  
MA, PGDip, BSc Hons, RM  
**President:**  
Rebecca Davies, RM  
**Patron:**  
HRH The Princess Royal

differential diagnosis of these entities. (Author)

Full URL: <https://doi.org/10.1016/j.ajog.2022.03.048>

---

## 2022-03010

**Guidance for people previously considered clinically extremely vulnerable from COVID-19.** Department of Health and Social Care (2022), 1 April 2022

Government advice for those who had previously received a letter or email identifying them as clinically extremely vulnerable (CEV) during the coronavirus pandemic. Includes advice on vaccination against COVID-19 in pregnancy. (JSM)

Full URL: <https://www.gov.uk/government/publications/guidance-on-shielding-and-protecting-extremely-vulnerable-persons-from-covid-19/guidance-on-shielding-and-protecting-extremely-vulnerable-persons-from-covid-19>

---

## 2022-02975

**Comparison of clinical features and perinatal outcomes between pre-variant and post-variant periods in pregnant women with SARS-CoV-2: analysis of 1935 cases.** Sahin D, Tanacan A, Anuk AT, et al (2022), Archives of Gynecology and Obstetrics 7 March 2022, online

### Purpose

To compare the clinical features and perinatal outcomes of pregnant women with severe acute respiratory syndrome coronavirus 2 (SARS-CoV-2) in the pre-variant and post-variant periods.

### Methods

This prospective cohort study includes pregnant women with SARS-CoV-2 who were followed-up at Ankara City Hospital between 11, March 2020 and 15, September 2021. Demographic features, clinical characteristics and pregnancy outcomes were compared between the pre-variant (n = 1416) and post-variant (n = 519) groups.

### Results

The rates of severe and critical cases significantly increased in the post-variant group (9.7% vs 2%,  $p < 0.001$ ). The rates of respiratory support (26.8% vs 7.3%,  $p < 0.001$ ), ICU admission (12.9% vs 1.8%,  $p < 0.001$ ) and maternal mortality (2.9% vs 0.4%,  $p < 0.001$ ) were significantly higher in the post-variant group. A significant increase was observed for pregnancy complications in the post-variant group (45.6% vs 18.8%,  $p = 0.007$ ). The rates of preterm delivery (26.4% vs 4.4%,  $p < 0.001$ ) and NICU admission (34% vs 18.8%,  $p < 0.001$ ) were significantly higher in the post-variant group. Positive, weak, statistically significant correlations were observed between the post-variant period, disease severity and maternal mortality ( $r = 0.19$ ,  $r = 0.12$  and  $p < 0.001$ ).

### Conclusion

Post-variant COVID-19 period was associated with a severe course of the disease and increased rates of adverse obstetric outcomes in pregnant patients.. (Author)

Full URL: <https://doi.org/10.1007/s00404-022-06493-5>

---

MIDIRS is part of RCM Information Services Limited which is a company incorporated in England and Wales under company no.11914882 with registered office at 10-18 Union Street, London SE1 1SZ  
RCM Information Services Limited is a subsidiary of The Royal College of Midwives



**The Royal College of Midwives Trust**  
10-18 Union Street  
London  
SE1 1SZ



**Open 24 hours a day, 7 days a week**  
T: 0300 303 0444  
F: +44 20 7312 3536  
E: [info@rcm.org.uk](mailto:info@rcm.org.uk)  
W: [www.rcm.org.uk](http://www.rcm.org.uk)



**Chief Executive:**  
Gill Walton,  
MA, PGDip, BSc Hons, RM  
**President:**  
Rebecca Davies, RM  
**Patron:**  
HRH The Princess Royal

**2022-02974**

**Perinatal outcomes of pregnant women with severe COVID-19 requiring extracorporeal membrane oxygenation (ECMO): a case series and literature review.** Clemenza S, Zullino S, Migliavacca C, et al (2022), Archives of Gynecology and Obstetrics vol 305, no 5, May 2022, pp 1135-1142

Purpose

Pregnant women with severe acute respiratory syndrome coronavirus 2 (SARS-CoV-2) infection have a higher risk of hospitalization, admission to intensive care unit (ICU) and invasive ventilation, and of acute respiratory distress syndrome (ARDS). In case of ARDS and critical severe coronavirus disease 2019 (COVID-19), the use of extracorporeal membrane oxygenation (ECMO) is recommended when other respiratory support strategies (oxygen insufflation, non-invasive ventilation [NIV], invasive ventilation through an endotracheal tube) are insufficient. However, available data on ECMO in pregnant and postpartum women with critical COVID-19 are very limited.

Methods

A case series of three critically ill pregnant women who required ECMO support for COVID-19 in pregnancy and/or in the postpartum period.

Results

The first patient tested positive for COVID-19 during the second trimester, she developed ARDS and required ECMO for 38 days. She was discharged in good general conditions and a cesarean-section [CS] at term was performed for obstetric indication. The second patient developed COVID-19-related ARDS at 28 weeks of gestation. During ECMO, she experienced a precipitous vaginal delivery at 31 weeks and 6 days of gestation. She was discharged 1 month later in good general conditions. The third patient, an obese 43-year-old woman, tested positive at 38 weeks and 2 days of gestation. Because of the worsening of clinical condition, a CS was performed, and she underwent ECMO. 143 days after the CS, she died because of sepsis and multiple organ failure (MOF). Thrombosis, hemorrhage and infections were the main complications among our patients. Neonatal outcomes have been positive.

Conclusion

ECMO should be considered a life-saving therapy for pregnant women with severe COVID-19. (Author)

Full URL: <https://doi.org/10.1007/s00404-022-06479-3>

---

**2022-02946**

**Pregnancy affected by SARS-CoV-2 infection: a flash report from Michigan.** Qadri F, Mariona F (2022), Journal of Maternal-Fetal and Neonatal Medicine vol 35, no 9, 2022, pp 1805-1807

The world is currently affected by the invasion of a human to human highly transmissible novel corona virus classified as SARS-CoV-2. It causes a severe acute lower respiratory tract syndrome named corona virus disease (CoVid-19). The virus is detected primarily by RT-PCR. The reproduction number (Ro) has been reported between 2.28 and 5.27]. It is beyond our objective to provide an in-depth discussion of the virus characteristics and its distinct viral clades and pathogenic behavior. On 30 January 2020 the World Health Organization (WHO) declared this outbreak a Public Health Emergency of International Concern, (PHEIC) and on 11 March 2020 WHO declared it a pandemic. There is limited information on the effect of CoVid-19 in pregnancy and the new born. We describe the details of the hospital course of the first 16 cases involving pregnant women, admitted to an urban-suburban community general hospital in Wayne County Michigan, from 26 March to 10 April 2020. At the time of this writing the Covid-19 pandemic has affected 35,291 persons in the state of Michigan (0.37%) making it the third most affected state in the USA (MDHHS). Pregnant women are believed to be at higher risk of Covid-19 infection in association with the known physiologic changes of the immune, cardiorespiratory and metabolic systems during pregnancy. (Author)

Full URL: <https://doi.org/10.1080/14767058.2020.1765334>

---

MIDIRS is part of RCM Information Services Limited which is a company incorporated in England and Wales under company no.11914882 with registered office at 10-18 Union Street, London SE1 1SZ  
RCM Information Services Limited is a subsidiary of The Royal College of Midwives



**The Royal College of Midwives Trust**  
10-18 Union Street  
London  
SE1 1SZ



**Open 24 hours a day, 7 days a week**  
T: 0300 303 0444  
F: +44 20 7312 3536  
E: [info@rcm.org.uk](mailto:info@rcm.org.uk)  
W: [www.rcm.org.uk](http://www.rcm.org.uk)



**Chief Executive:**  
Gill Walton,  
MA, PGDip, BSc Hons, RM  
**President:**  
Rebecca Davies, RM  
**Patron:**  
HRH The Princess Royal

**2022-02944**

**Management of covid-19: a practical guideline for maternal and newborn health care providers in Sub-Saharan Africa.**

Ezenwa BN, Fajolu IB, Akinajo OR, et al (2022), Journal of Maternal-Fetal and Neonatal Medicine vol 35, no 9, 2022, pp 1789-1795

COVID-19 is a pandemic that is currently ravaging the world. Infection rate is steadily increasing in Sub-Saharan Africa. Pregnant women and their infants may suffer severe illnesses due to their lower immunity. This guideline prepares and equips clinicians working in the maternal and newborn sections in the sub-region to manage COVID-19 during pregnancy and childbirth.

(Author)

**Full URL:** <https://doi.org/10.1080/14767058.2020.1763948>

---

**2022-02920**

**Impact of COVID-19 on perinatal care: Perceptions of family physicians in the United States.** Goldstein JT, Eden AR, Taylor MK, et

al (2022), Birth 9 April 2022, online

**Background**

Patient-centered care is the best practice in the care of pregnant and postpartum patients. The COVID-19 pandemic prompted changes in perinatal care policies, which were often reactive, resulting in unintended consequences, many of which made the delivery of patient-centered care more difficult. This study aimed to understand the impact of the COVID-19 pandemic on perinatal health care delivery from the perspective of family physicians in the United States.

**Methods**

From October 5 to November 4, 2020, we surveyed mid- to late-career family physicians who provide perinatal care. We conducted descriptive analyses to measure the impact of COVID-19 on prenatal care, labor and delivery, postpartum care, patient experience, and patient volume. An immersion-crystallization approach was used to analyze qualitative data provided as open-text comments.

**Results**

Of the 1518 survey respondents, 1062 (69.8%) stated that they currently attend births; 595 of those elaborated about the impact of COVID-19 on perinatal care in free-text comments. Eight themes emerged related to the impact of COVID-19 on perinatal care: visitation, patient decisions, testing, personal protective equipment, care continuity, changes in care delivery, reassessment, and volume. The greatest perceived impact of COVID-19 was on patient experience.

**Conclusions**

Family physicians who provided perinatal care during the COVID-19 pandemic noted a considerable impact on patient experience, which particularly affected the ability to deliver patient-centered and family-centered care. Continued research is needed to understand the long-term impact of policies affecting the delivery of patient-centered perinatal care and to inform more evidence-based, proactive policies to be implemented in future pandemic or disaster situations. (Author)

**Full URL:** <https://doi.org/10.1111/birt.12637>

---

MIDIRS is part of RCM Information Services Limited which is a company incorporated in England and Wales under company no.11914882 with registered office at 10-18 Union Street, London SE1 1SZ  
RCM Information Services Limited is a subsidiary of The Royal College of Midwives



**The Royal College of Midwives Trust**  
10-18 Union Street  
London  
SE1 1SZ



**Open 24 hours a day, 7 days a week**  
T: 0300 303 0444  
F: +44 20 7312 3536  
E: [info@rcm.org.uk](mailto:info@rcm.org.uk)  
W: [www.rcm.org.uk](http://www.rcm.org.uk)



**Chief Executive:**  
Gill Walton,  
MA, PGDip, BSc Hons, RM  
**President:**  
Rebecca Davies, RM  
**Patron:**  
HRH The Princess Royal

**2022-02914**

**COVID-19 pneumonia and pregnancy; a systematic review and meta-analysis.** Kasraeian M, Zare M, Vafaei H, et al (2022), Journal of Maternal-Fetal and Neonatal Medicine vol 35, no 9, 2022, pp 1652-1659

#### Background

The new SARS-CoV-2 originated from Wuhan, China is spreading rapidly worldwide. A number of SARS-CoV-2 positive pregnant women have been reported. However, more information is still needed on the pregnancy outcome and the neonates regarding COVID-19 pneumonia.

#### Material and Methods

A systematic search was done and nine articles on COVID-19 pneumonia and SARS-CoV-2 positive pregnant women were extracted. Some maternal-fetal characteristics were extracted to be included in the meta-analysis.

#### Results

The present meta-analysis was conducted on 87 SARS-CoV-2 positive pregnant women. Almost 65% of the patients reported a history of exposure to an infected person, 78% suffered from mild or moderate COVID-19, 99.9% had successful termination, 86% had cough, and 68% had fever ( $p = .022$  and  $p < .001$ ). The overall proportions of vertical transmission, still birth, and neonatal death were zero, 0.002, and, 0.002, respectively ( $p = 1$ ,  $p = .86$ , and  $p = .89$ , respectively). The means of the first- and fifth-minute Apgar scores were 8.86 and 9, respectively ( $p < .001$  for both). The confounding role of history of underlying diseases with an estimated overall proportion of 33% ( $p = .03$ ) resulted in further investigations due to sample size limitation. A natural history of COVID-19 pneumonia in the adult population was presented, as well.

#### Conclusion

Currently, no evidence of vertical transmission has been suggested at least in late pregnancy. No hazards have been detected for fetuses or neonates. Although pregnant women are at an immunosuppressive state due to the physiological changes during pregnancy, most patients suffered from mild or moderate COVID-19 pneumonia with no pregnancy loss, proposing a similar pattern of the clinical characteristics of COVID-19 pneumonia to that of other adult populations. (Author)

**Full URL:** <https://doi.org/10.1080/14767058.2020.1763952>

---

MIDIRS is part of RCM Information Services Limited which is a company incorporated in England and Wales under company no.11914882 with registered office at 10-18 Union Street, London SE1 1SZ  
RCM Information Services Limited is a subsidiary of The Royal College of Midwives



**The Royal College of Midwives Trust**  
10-18 Union Street  
London  
SE1 1SZ



**Open 24 hours a day, 7 days a week**  
T: 0300 303 0444  
F: +44 20 7312 3536  
E: [info@rcm.org.uk](mailto:info@rcm.org.uk)  
W: [www.rcm.org.uk](http://www.rcm.org.uk)



**Chief Executive:**  
Gill Walton,  
MA, PGDip, BSc Hons, RM  
**President:**  
Rebecca Davies, RM  
**Patron:**  
HRH The Princess Royal

**2022-02913**

**Antenatal corticosteroids for pregnant women with COVID-19 infection and preterm prelabor rupture of membranes: a decision analysis.** Zhou CG, Packer CH, Hersh AR, et al (2022), Journal of Maternal-Fetal and Neonatal Medicine vol 35, no 9, 2022, pp 1643-1651

#### Background

While antenatal corticosteroids are routinely used to decrease adverse neonatal outcomes following preterm delivery, corticosteroids are also associated with worse outcomes in patients with viral respiratory infections. Currently in the setting of the COVID-19 pandemic, it is unclear whether antenatal corticosteroids for infant benefit outweigh the potential harm to a pregnant woman with a COVID-19 infection.

#### Objective

To determine at which gestational ages administering antenatal corticosteroids is the optimal management strategy for hospitalized women with preterm prelabor rupture of membranes (PPROM) who have a COVID-19 infection.

#### Methods

We designed a decision-analytic model to assess the maternal and infant outcomes associated with antenatal corticosteroid administration for risk of preterm delivery following rupture of membranes in the setting of a COVID-19 infection. We used a theoretical cohort of 10,000 women at each gestational age between 24 and 32 weeks who were hospitalized with PPRM and found to be COVID-19 positive. Maternal outcomes included intensive care unit admission and death related to COVID-19 infection. The infant outcomes of interest included respiratory distress syndrome, intraventricular hemorrhage, neurodevelopmental delay, and death, and were assessed along with maternal and infant quality-adjusted life years (QALYs). Deterministic and probabilistic sensitivity analyses were used to evaluate model assumptions.

#### Results

In our theoretical cohort of 10,000 women with COVID-19 infection and preterm prelabor rupture of membrane between 24 and 32 weeks, corticosteroid administration resulted in 2,200 women admitted to the ICU and 110 maternal deaths at each gestational age. No antenatal corticosteroid use resulted in 1,500 ICU admissions and 75 maternal deaths at each gestational age. Antenatal corticosteroid administration also resulted in fewer cases of respiratory distress syndrome, intraventricular hemorrhage, and infant death. Overall, we found that between 24 and 30 weeks of gestation, administering antenatal corticosteroids was the optimal management strategy as it resulted in higher combined QALYs than no corticosteroid use. For 31 and 32 weeks of gestation, antenatal corticosteroid administration resulted in lower combined QALYs. On sensitivity analyses, we found that with increasing gestational age, the probability which antenatal corticosteroids was the optimal management strategy decreased.

#### Conclusion

Administration of antenatal corticosteroids was an effective management strategy compared to no corticosteroid administration at gestational ages less than 31 weeks. These results provide data for clinicians to utilize when counseling pregnant patients hospitalized with PPRM and have a COVID-19 infection. (Author)

**Full URL:** <https://doi.org/10.1080/14767058.2020.1763951>

---

MIDIRS is part of RCM Information Services Limited which is a company incorporated in England and Wales under company no.11914882 with registered office at 10-18 Union Street, London SE1 1SZ  
RCM Information Services Limited is a subsidiary of The Royal College of Midwives



**The Royal College of Midwives Trust**  
10-18 Union Street  
London  
SE1 1SZ



**Open 24 hours a day, 7 days a week**  
T: 0300 303 0444  
F: +44 20 7312 3536  
E: [info@rcm.org.uk](mailto:info@rcm.org.uk)  
W: [www.rcm.org.uk](http://www.rcm.org.uk)



**Chief Executive:**  
Gill Walton,  
MA, PGDip, BSc Hons, RM  
**President:**  
Rebecca Davies, RM  
**Patron:**  
HRH The Princess Royal

**2022-02910**

**Coronavirus disease 2019 (COVID-19) and pregnancy: a systematic review.** Yang Z, Wang M, Zhu Z, et al (2022), Journal of Maternal-Fetal and Neonatal Medicine vol 35, no 8, 2022, pp 1619-1622

#### Objective

To summarize currently available evidence on maternal, fetal, and neonatal outcomes of pregnant women infected with Coronavirus Disease 2019 (COVID-19).

#### Material and methods

PubMed, Google Scholar, CNKI, Wanfang Data, VIP, and CBMdisc were searched for studies reporting maternal, fetal, and neonatal outcomes of women infected with COVID-19 published from 1 January 2020 to 26 March 2020. The protocol was registered with the Open Science Framework (DOI: 10.17605/OSF.IO/34ZAV).

#### Results

In total, 18 studies comprising 114 pregnant women were included in the review. Fever (87.5%) and cough (53.8%) were the most commonly reported symptoms, followed by fatigue (22.5%), diarrhea (8.8%), dyspnea (11.3%), sore throat (7.5%), and myalgia (16.3%). The majority of patients (91%) had cesarean delivery due to various indications. In terms of fetal and neonatal outcomes, stillbirth (1.2%), neonatal death (1.2%), preterm birth (21.3%), low birth weight (<2500 g, 5.3%), fetal distress (10.7%), and neonatal asphyxia (1.2%) were reported. There are reports of neonatal infection, but no direct evidence of intrauterine vertical transmission has been found.

#### Conclusions

The clinical characteristics of pregnant women with COVID-19 are similar to those of non-pregnant adults. Fetal and neonatal outcomes appear good in most cases, but available data only include pregnant women infected in their third trimesters. Further studies are needed to ascertain long-term outcomes and potential intrauterine vertical transmission. (Author)

**Full URL:** <https://doi.org/10.1080/14767058.2020.1759541>

---

MIDIRS is part of RCM Information Services Limited which is a company incorporated in England and Wales under company no.11914882 with registered office at 10-18 Union Street, London SE1 1SZ  
RCM Information Services Limited is a subsidiary of The Royal College of Midwives



**The Royal College of Midwives Trust**  
10-18 Union Street  
London  
SE1 1SZ



**Open 24 hours a day, 7 days a week**  
T: 0300 303 0444  
F: +44 20 7312 3536  
E: [info@rcm.org.uk](mailto:info@rcm.org.uk)  
W: [www.rcm.org.uk](http://www.rcm.org.uk)



**Chief Executive:**  
Gill Walton,  
MA, PGDip, BSc Hons, RM  
**President:**  
Rebecca Davies, RM  
**Patron:**  
HRH The Princess Royal



## 2022-02842

**A qualitative study of Swedish fathers' experiences of becoming a father during the COVID-19 pandemic.** Wells MB, Svahn J, Svedlind K, et al (2022), European Journal of Midwifery vol 6, March 2022, p 15

### Introduction:

Expectant fathers want to participate in perinatal care. COVID-19 policies restrict their access to care, but it is unknown how these policies have affected them. The aim of this study is to explore the perinatal care given to and wanted by expectant and new fathers during the COVID-19 pandemic in Sweden.

### Methods:

The current study used an inductive qualitative design where 14 expectant or new fathers participated in a video- or telephone-based semi-structured interview. Interviews lasted 20 minutes, on average. The collected data were analyzed using content analysis.

### Results:

Two main themes were reported: 1) 'Being left out, but trying to remain positive', and 2) immediate consequences related to restrictions. Expectant fathers were not able to attend as many perinatal visits as they wanted to, due to the COVID-19 restrictions on non-birthing parents. Expectant fathers regretted and felt discouraged that they could not support their partner during visits and not follow their baby's growth and development. Furthermore, they faced uncertainties and stress regarding whether or not they could attend the birth of their child. Fathers reported how their exclusion negatively impacted the entire family.

### Conclusions:

Expectant and new fathers felt that their level of participation in prenatal care was negatively impacted by the Swedish policies imposed on them during the COVID-19 pandemic. Fathers were physically and emotionally excluded, resulting in receiving little direct care support, and lacked companionship with other parents. Fathers provided suggestions and alternatives on how to increase their participation with individual midwives and from an organizational level. (Author)

**Full URL:** <https://doi.org/10.18332/ejm/146082>

---

## 2022-02785

**COVID-19 in pregnancy: A UK perspective.** Zayyan S, Frise C (2022), Obstetric Medicine 23 March 2022, online

This article aims to review data collected and analysed so far over the course of the coronavirus pandemic, that examine demographic associations, patterns of disease, severity and outcomes of COVID-19 in pregnancy in the UK. (Author, edited)

---

MIDIRS is part of RCM Information Services Limited which is a company incorporated in England and Wales under company no.11914882 with registered office at 10-18 Union Street, London SE1 1SZ  
RCM Information Services Limited is a subsidiary of The Royal College of Midwives



**The Royal College of Midwives Trust**  
10-18 Union Street  
London  
SE1 1SZ



**Open 24 hours a day, 7 days a week**  
T: 0300 303 0444  
F: +44 20 7312 3536  
E: [info@rcm.org.uk](mailto:info@rcm.org.uk)  
W: [www.rcm.org.uk](http://www.rcm.org.uk)



**Chief Executive:**  
Gill Walton,  
MA, PGDip, BSc Hons, RM  
**President:**  
Rebecca Davies, RM  
**Patron:**  
HRH The Princess Royal

**2022-02781**

**The United Kingdom and the Netherlands maternity care responses to COVID-19: A comparative study.** van den Berg LMM, Balaam M-C, Nowland R, et al (2022), *Women and Birth: Journal of the Australian College of Midwives* 4 April 2022, online

#### Background

The national health care response to coronavirus (COVID-19) has varied between countries. The United Kingdom (UK) and the Netherlands (NL) have comparable maternity and neonatal care systems, and experienced similar numbers of COVID-19 infections, but had different organisational responses to the pandemic. Understanding why and how similarities and differences occurred in these two contexts could inform optimal care in normal circumstances, and during future crises.

#### Aim

To compare the UK and Dutch COVID-19 maternity and neonatal care responses in three key domains: choice of birthplace, companionship, and families in vulnerable situations.

#### Method

A multi-method study, including documentary analysis of national organisation policy and guidance on COVID-19, and interviews with national and regional stakeholders.

#### Findings

Both countries had an infection control focus, with less emphasis on the impact of restrictions, especially for families in vulnerable situations. Differences included care providers' fear of contracting COVID-19; the extent to which community- and personalised care was embedded in the care system before the pandemic; and how far multidisciplinary collaboration and service-user involvement were prioritised.

#### Conclusion

We recommend that countries should 1) make a systematic plan for crisis decision-making before a serious event occurs, and that this must include authentic service-user involvement, multidisciplinary collaboration, and protection of staff wellbeing 2) integrate women's and families' values into the maternity and neonatal care system, ensuring equitable inclusion of the most vulnerable and 3) strengthen community provision to ensure system wide resilience to future shocks from pandemics, or other unexpected large-scale events. (Author)

**Full URL:** <https://doi.org/10.1016/j.wombi.2022.03.010>

---

MIDIRS is part of RCM Information Services Limited which is a company incorporated in England and Wales under company no.11914882 with registered office at 10-18 Union Street, London SE1 1SZ  
RCM Information Services Limited is a subsidiary of The Royal College of Midwives



**The Royal College of Midwives Trust**  
10-18 Union Street  
London  
SE1 1SZ



**Open 24 hours a day, 7 days a week**  
T: 0300 303 0444  
F: +44 20 7312 3536  
E: [info@rcm.org.uk](mailto:info@rcm.org.uk)  
W: [www.rcm.org.uk](http://www.rcm.org.uk)



**Chief Executive:**  
Gill Walton,  
MA, PGDip, BSc Hons, RM  
**President:**  
Rebecca Davies, RM  
**Patron:**  
HRH The Princess Royal

## 2022-02723

**The COVID-19 Pandemic and Psychopathological symptoms in pregnant women in Spain.** Garcia-Leon MA, Martin-Tortosa PL, Cambio-Ledesma A, et al (2022), Journal of Reproductive and Infant Psychology 9 March 2022, online

### Background

During the COVID-19 pandemic, pregnant women are exposed to potentially harmful stressors that might affect their health. The direct consequences that SARS-CoV-2 may have on perinatal mental health are still unknown.

### Objective

The present study aimed to explore the impact of the COVID-19 pandemic on psychopathological symptoms in a sample of Spanish pregnant women.

### Methods

A sample of 186 pregnant women was assessed using the revised Symptoms Check List-90 during the first lockdown in Spain.

### Results

The results showed clinical scores on the obsession and compulsion, anxiety and phobic anxiety subscales, as well as on the severity indexes. Phobic anxiety was the only variable that was inversely correlated with age and the number of previous miscarriages. A linear regression model showed that age was inversely associated with phobic anxiety scores. A younger age was associated with higher levels of phobic anxiety symptoms.

### Conclusions

Our results indicated that younger pregnant women and women in the first trimester of pregnancy were more vulnerable to the effects of stress and concerns about COVID-19. (Author)

---

## 2022-02685

**Concurrent caecum perforation in second trimester pregnant woman complicated with SARS-CoV-2 infection: a case report.**

Turgut ÜK, Erdemoğlu E, Kılçar M, et al (2022), Journal of Obstetrics and Gynaecology 9 March 2022, online

Case report of a 32-year-old pregnant woman presenting with SARS-CoV-2 and caecal perforation requiring urgent surgical intervention. (LDO)

---

## 2022-02684

**First and second waves of SARS-COV-2 infection in the obstetric population.** Sangaletti M, Gibellini D, Diani E, et al (2022), Journal of Obstetrics and Gynaecology 11 March 2022, online

Brief report exploring the impact of the first and second waves of the COVID-19 pandemic on pregnant women. (LDO)

---

MIDIRS is part of RCM Information Services Limited which is a company incorporated in England and Wales under company no.11914882 with registered office at 10-18 Union Street, London SE1 1SZ  
RCM Information Services Limited is a subsidiary of The Royal College of Midwives



**The Royal College of Midwives Trust**  
10-18 Union Street  
London  
SE1 1SZ



**Open 24 hours a day, 7 days a week**  
T: 0300 303 0444  
F: +44 20 7312 3536  
E: [info@rcm.org.uk](mailto:info@rcm.org.uk)  
W: [www.rcm.org.uk](http://www.rcm.org.uk)



**Chief Executive:**  
Gill Walton,  
MA, PGDip, BSc Hons, RM  
**President:**  
Rebecca Davies, RM  
**Patron:**  
HRH The Princess Royal

**2022-02652**

**Early pregnancy anxiety during the COVID-19 pandemic: preliminary findings from the UCSF ASPIRE study.** Morris JR, Jaswa EG, Kaing A, et al (2022), BMC Pregnancy and Childbirth vol 22, no 272, 31 March 2022

#### Background

Antenatal anxiety has been linked to adverse obstetric outcomes, including miscarriage and preterm birth. However, most studies investigating anxiety during pregnancy, particularly during the COVID-19 pandemic, have focused on symptoms during the second and third trimester. This study aims to describe the prevalence of anxiety symptoms early in pregnancy and identify predictors of early pregnancy anxiety during the COVID-19 pandemic.

#### Methods

We assessed baseline moderate-to-severe anxiety symptoms after enrollment in the UCSF ASPIRE (Assessing the Safety of Pregnancy in the Coronavirus Pandemic) Prospective Cohort from May 2020 through February 2021. Pregnant persons < 10 weeks' gestation completed questions regarding sociodemographic characteristics, obstetric/medical history, and pandemic-related experiences. Univariate and multivariate hierarchical logistic regression analyses determined predictors of moderate or severe anxiety symptoms (Generalized Anxiety Disorder-7 questionnaire score  $\geq 10$ ). All analyses performed with Statistical Analysis Software (SAS<sup>®</sup>) version 9.4.

#### Results

A total of 4,303 persons completed the questionnaire. The mean age of this nationwide sample was 33 years of age and 25.7% of participants received care through a fertility clinic. Over twelve percent of pregnant persons reported moderate-to-severe anxiety symptoms. In univariate analysis, less than a college education ( $p < 0.0001$ ), a pre-existing history of anxiety ( $p < 0.0001$ ), and a history of prior miscarriage ( $p = 0.0143$ ) were strong predictors of moderate-to-severe anxiety symptoms. Conversely, having received care at a fertility center was protective (26.6% vs. 25.7%,  $p = 0.0009$ ). COVID-19 related stressors including job loss, reduced work hours during the pandemic, inability to pay rent, very or extreme worry about COVID-19, and perceived stress were strongly predictive of anxiety in pregnancy ( $p < 0.0001$ ). In the hierarchical logistic regression model, pre-existing history of anxiety remained associated with anxiety during pregnancy, while the significance of the effect of education was attenuated.

#### Conclusion(s)

Pre-existing history of anxiety and socioeconomic factors likely exacerbated the impact of pandemic-related stressors on early pregnancy anxiety symptoms during the COVID-19 pandemic. Despite on-going limitations for in-person prenatal care administration, continued emotional health support should remain an important focus for providers, particularly when caring for less privileged pregnant persons or those with a pre-existing history of anxiety. (Author)

**Full URL:** <https://doi.org/10.1186/s12884-022-04595-1>

---

MIDIRS is part of RCM Information Services Limited which is a company incorporated in England and Wales under company no.11914882 with registered office at 10-18 Union Street, London SE1 1SZ  
RCM Information Services Limited is a subsidiary of The Royal College of Midwives



**The Royal College of Midwives Trust**  
10-18 Union Street  
London  
SE1 1SZ



**Open 24 hours a day, 7 days a week**  
T: 0300 303 0444  
F: +44 20 7312 3536  
E: [info@rcm.org.uk](mailto:info@rcm.org.uk)  
W: [www.rcm.org.uk](http://www.rcm.org.uk)



**Chief Executive:**  
Gill Walton,  
MA, PGDip, BSc Hons, RM  
**President:**  
Rebecca Davies, RM  
**Patron:**  
HRH The Princess Royal

**2022-02650**

**Coping with Covid-19: stress, control and coping among pregnant women in Ireland during the Covid-19 pandemic.** Crowe S, Sarma KVR (2022), BMC Pregnancy and Childbirth vol 22, no 274, 1 April 2022

#### Background

The aim of the current study is to investigate the relationship between perceived control, coping and psychological distress among pregnant women in Ireland during the Covid-19 pandemic. It is hypothesised that lower levels of perceived control, greater use of avoidant coping and greater Covid-19 related pregnancy concern will be associated with psychological distress. In addition, it is hypothesised that the relationship between Covid-19 related pregnancy concern and psychological distress will be moderated by perceived control and avoidant coping.

#### Method

The study is cross-sectional, utilizing an online questionnaire, which was completed by 761 women in January 2021. The questionnaire includes measures of perceived control, coping style, perceived stress, anxiety and depression.

#### Results

Correlation analyses found that lower levels of perceived control were associated with higher levels of avoidant coping and psychological distress. There was also a significant positive relationship between avoidant coping and psychological distress. Using multiple regression, perceived control, avoidant coping and Covid-19 related pregnancy concern were found to predict 51% of the variance in psychological distress. However, in the moderation analysis, perceived control and avoidant coping were not found to moderate the relationship between Covid-19 related pregnancy concern and psychological distress.

#### Conclusion

The results from this study suggest that pregnant women in Ireland are experiencing increased levels of psychological distress during the Covid-19 pandemic. The findings also suggest that perceptions of control and avoidant coping are associated with psychological distress in this group and could be used as intervention targets. (Author)

**Full URL:** <https://doi.org/10.1186/s12884-022-04579-1>

---

MIDIRS is part of RCM Information Services Limited which is a company incorporated in England and Wales under company no.11914882 with registered office at 10-18 Union Street, London SE1 1SZ  
RCM Information Services Limited is a subsidiary of The Royal College of Midwives



**The Royal College of Midwives Trust**  
10-18 Union Street  
London  
SE1 1SZ



**Open 24 hours a day, 7 days a week**  
T: 0300 303 0444  
F: +44 20 7312 3536  
E: [info@rcm.org.uk](mailto:info@rcm.org.uk)  
W: [www.rcm.org.uk](http://www.rcm.org.uk)



**Chief Executive:**  
Gill Walton,  
MA, PGDip, BSc Hons, RM  
**President:**  
Rebecca Davies, RM  
**Patron:**  
HRH The Princess Royal

**2022-02647**

**The 'new normal' includes online prenatal exercise: exploring pregnant women's experiences during the pandemic and the role of virtual group fitness on maternal mental health.** Silva-Jose C, Nagpal TS, Coterón J, et al (2022), BMC Pregnancy and Childbirth vol 22, no 251, 25 March 2022

#### Background

Prenatal anxiety and depressive symptoms have significantly increased since the onset of the coronavirus (COVID-19) pandemic. In addition, home confinement regulations have caused a drastic increase in time spent sedentary. Online group fitness classes may be an effective strategy that can increase maternal physical activity levels and improve mental health outcomes by providing an opportunity for social connectedness. The present study explores the experiences of pregnant women who participated in an online group exercise program during the pandemic and identifies relationships with maternal mental health and well-being. In addition, we present person-informed recommendations on how to improve the delivery of future online prenatal exercise programs.

#### Methods

Semi-structured interviews were conducted with pregnant women (8-39 weeks of pregnancy) who participated in an online group exercise program, from March to October 2020 in Spain. A phenomenological approach was taken, and open-ended questions were asked to understand women's experiences throughout the pandemic and the role the online exercise classes may have had on their physical activity levels, mental health, and other health behaviours such as diet. A thematic analysis was performed to evaluate data. In addition, women completed the State-Trait Anxiety Inventory and these data supplemented qualitative findings.

#### Results

Twenty-four women were interviewed, and the anxiety scores were on average  $32.23 \pm 9.31$ , ranging from low to moderate levels. Thematic analysis revealed that women felt safe exercising from home, an increased availability of time to schedule a structured exercise class, and consequently an improvement in their adherence to the program and other behaviours (i.e., healthier diet). Women emphasized feeling connected to other pregnant women when they exercised online together, and overall, this had a positive effect on their mental well-being. Women suggested that future online exercise programs should include flexible options, detailed instructions and facilitation by a qualified exercise professional.

#### Conclusion

Pregnant women are receptive to online group exercise classes and expressed that they are an accessible option to accommodating physical activity during the pandemic. In addition, the online group environment provides an important sense of connectivity among pregnant women exercising together and this may mitigate the detrimental effect of COVID-19 on maternal mental health. (Author)

**Full URL:** <https://doi.org/10.1186/s12884-022-04587-1>

---

MIDIRS is part of RCM Information Services Limited which is a company incorporated in England and Wales under company no.11914882 with registered office at 10-18 Union Street, London SE1 1SZ  
RCM Information Services Limited is a subsidiary of The Royal College of Midwives



**The Royal College of Midwives Trust**  
10-18 Union Street  
London  
SE1 1SZ



**Open 24 hours a day, 7 days a week**  
T: 0300 303 0444  
F: +44 20 7312 3536  
E: [info@rcm.org.uk](mailto:info@rcm.org.uk)  
W: [www.rcm.org.uk](http://www.rcm.org.uk)



**Chief Executive:**  
Gill Walton,  
MA, PGDip, BSc Hons, RM  
**President:**  
Rebecca Davies, RM  
**Patron:**  
HRH The Princess Royal

## 2022-02630

### Self-reported mental health status of pregnant women in Sweden during the COVID-19 pandemic: a cross-sectional survey.

Ho-Fung C, Andersson E, Hsuan-Ying H, et al (2022), BMC Pregnancy and Childbirth vol 22, no 260, 28 March 2022

#### Background

The COVID-19 pandemic has contributed to unprecedented worries and challenges for pregnant women due to social restrictions and changes in maternity care provision. We aimed to investigate the mental health impact of COVID-19 pandemic on pregnant women in Sweden and explore factors associated with poor perinatal mental health in this specific context.

#### Method

This was a nation-wide cross-sectional survey of pregnant women living in Sweden. Validated questionnaires were distributed through non-profit organizations' websites and social media channels from May 2020 to February 2021. Perinatal depression, anxiety, and acute stress reaction were assessed using the Edinburgh Postnatal Depression Scale (EPDS), Generalized Anxiety Disorder-7 (GAD-7) and Impact Event Scale (Revised) (IES-R), respectively. Sociodemographic characteristics and self-perceived mental well-being were also obtained. Factors associated with mental health outcomes were analyzed using multivariate logistic regression model.

#### Results

Among a total of 470 participants, 43.2% (n = 203) reported depression (EPDS  $\geq 13$ ), 25.7% (n = 121) moderate to severe anxiety (GAD-7 score  $\geq 10$ ), and 23.7% (n = 110) moderate to severe acute stress reaction (IES-R  $\geq 33$ ). 27.4% participants (n = 129) expressed concerns regarding their mental well-being during the pandemic. Pregnant mothers who had sick family members reported poorer mental health outcomes than those who did not (median [Interquartile range (IQR)] EPDS scores: 14.0 [8.75–18.0] vs 11.0 [6.25–15.0],  $p < .001$ ; median (IQR) GAD7 scores: 7.0 [4.0–12.25] vs 6.0 [3.0–9.0],  $p = .003$ ; median (IQR) IES-R scores: 20.0 [9.0–38.0] vs 15.0 [7.0–30.0],  $p = .048$ ). Logistic regression analyses revealed that risk factors for poor mental health outcomes were having a sick family member with any illness, unemployment, and experiencing a substantially stressful life event. Having a higher educational level and a younger age during the pandemic were protective.

#### Conclusion

Depression and anxiety were highly prevalent among pregnant women in Sweden during the COVID-19 pandemic, indicating a need for professional mental health support for this vulnerable group of population. Unemployment was an associated risk factor whereas younger age and higher educational level were protective suggesting an important role of socio-economic factors in modulating the impact of COVID-19 pandemic on perinatal mental health. (Author)

**Full URL:** <https://doi.org/10.1186/s12884-022-04553-x>

---

MIDIRS is part of RCM Information Services Limited which is a company incorporated in England and Wales under company no.11914882 with registered office at 10-18 Union Street, London SE1 1SZ  
RCM Information Services Limited is a subsidiary of The Royal College of Midwives



**The Royal College of Midwives Trust**  
10-18 Union Street  
London  
SE1 1SZ



**Open 24 hours a day, 7 days a week**  
T: 0300 303 0444  
F: +44 20 7312 3536  
E: [info@rcm.org.uk](mailto:info@rcm.org.uk)  
W: [www.rcm.org.uk](http://www.rcm.org.uk)



**Chief Executive:**  
Gill Walton,  
MA, PGDip, BSc Hons, RM  
**President:**  
Rebecca Davies, RM  
**Patron:**  
HRH The Princess Royal

**2022-02617**

**Estimation of sleep problems among pregnant women during COVID-19 pandemic: a systematic review and meta-analysis.**

Alimoradi Z, Abdi F, Gozal D, et al (2022), *BMJ Open* vol 12, no 4, April 2022, e056044

**Objective** To estimate the sleep problems among pregnant women during the COVID-19 pandemic.

**Eligibility criteria** English, peer-reviewed, observational studies published between December 2019 and July 2021 which assessed and reported sleep problem prevalence using a valid and reliable measure were included.

**Information sources** Scopus, Medline/PubMed Central, ProQuest, ISI Web of Knowledge and Embase.

**Risk of bias assessment tool** The Newcastle-Ottawa Scale checklist.

**Synthesis of results** Prevalence of sleep problems was synthesised using STATA software V.14 using a random effects model. To assess moderator analysis, meta-regression was carried out. Funnel plot and Egger's test were used to assess publication bias. Meta-trim was used to correct probable publication bias. The jackknife method was used for sensitivity analysis.

**Included studies** A total of seven cross-sectional studies with 2808 participants from four countries were included.

**Synthesis of results** The pooled estimated prevalence of sleep problems was 56% (95% CI 23% to 88%,  $I^2=99.81%$ ,  $\text{Tau}^2=0.19$ ). Due to the probability of publication bias, the fill-and-trim method was used to correct the estimated pooled measure, which imputed four studies. The corrected results based on this method showed that pooled prevalence of sleep problems was 13% (95% CI 0% to 45%;  $p<0.001$ ). Based on meta-regression, age was the only significant predictor of prevalence of sleep problems among pregnant women.

**Limitations of evidence** All studies were cross-sectional absence of assessment of sleep problems prior to COVID-19, and the outcomes of the pregnancies among those with and without sleep problems in a consistent manner are among the limitation of the current review.

**Interpretation** Pregnant women have experienced significant declines in sleep quality when faced with the COVID-19 pandemic. The short-term and long-term implications of such alterations in sleep on gestational and offspring outcomes are unclear and warrant further studies.

PROSPERO registration number CRD42020181644. (Author)

**Full URL:** <http://dx.doi.org/10.1136/bmjopen-2021-056044>

---

MIDIRS is part of RCM Information Services Limited which is a company incorporated in England and Wales under company no.11914882 with registered office at 10-18 Union Street, London SE1 1SZ  
RCM Information Services Limited is a subsidiary of The Royal College of Midwives



**The Royal College of Midwives Trust**  
10-18 Union Street  
London  
SE1 1SZ



**Open 24 hours a day, 7 days a week**  
T: 0300 303 0444  
F: +44 20 7312 3536  
E: [info@rcm.org.uk](mailto:info@rcm.org.uk)  
W: [www.rcm.org.uk](http://www.rcm.org.uk)



**Chief Executive:**  
Gill Walton,  
MA, PGDip, BSc Hons, RM  
**President:**  
Rebecca Davies, RM  
**Patron:**  
HRH The Princess Royal



**2022-02611**

**Knowledge and attitudes among Lebanese pregnant women and women seeking fertility treatment during the COVID-19 outbreak: a cross-sectional survey.** El Taha L, Beyrouthy C, Tamim H, et al (2022), *BMJ Open* vol 12, no 3, March 2022, e057873

Objectives COVID-19 has been recognised as a global health emergency necessitating collaborative efforts to halt further disease spread. The success of public health interventions and vaccination campaigns is contingent on the knowledge and awareness level of the public. We aim to assess COVID-19 knowledge and attitudes among Lebanese pregnant women and women seeking fertility treatment.

Design Cross-sectional study using telehealth administered survey.

Setting University-affiliated tertiary care centre.

Participants The data of 402-Lebanese women pregnant or seeking fertility treatment aged 20–45 years were analysed.

Outcome measures Extent of COVID-19 general knowledge, pregnancy-specific knowledge and attitudes toward COVID-19 practices.

Results All participants reported being knowledgeable about COVID-19, 70% of which rated their knowledge as 7 or more on a numerical scale of 0–10. The mean general COVID-19 knowledge was 22.15 (SD 2.44, range 14–27) indicating a high level of knowledge. The mean pregnancy-specific COVID-19 knowledge 6.84 (SD 2.061, range 0–10) indicated poorer pregnancy-specific knowledge compared with general COVID-19 knowledge. A trend towards higher knowledge was noted with higher income status. Reproductive age women with higher pregnancy-specific knowledge had more positive attitudes toward COVID-19 pregnancy practices.

Conclusion Our findings suggest a deficiency in pregnancy-specific COVID-19 knowledge stressing the necessity for targeted public health education interventions. It highlights the need for enhancing COVID-19 pregnancy-specific awareness which can serve as a stepping stone in the success of COVID-19 vaccination campaigns and in halting further disease spread. (Author)

**Full URL:** <http://dx.doi.org/10.1136/bmjopen-2021-057873>

---

MIDIRS is part of RCM Information Services Limited which is a company incorporated in England and Wales under company no.11914882 with registered office at 10-18 Union Street, London SE1 1SZ  
RCM Information Services Limited is a subsidiary of The Royal College of Midwives



**The Royal College of Midwives Trust**  
10-18 Union Street  
London  
SE1 1SZ



**Open 24 hours a day, 7 days a week**  
T: 0300 303 0444  
F: +44 20 7312 3536  
E: [info@rcm.org.uk](mailto:info@rcm.org.uk)  
W: [www.rcm.org.uk](http://www.rcm.org.uk)



**Chief Executive:**  
Gill Walton,  
MA, PGDip, BSc Hons, RM  
**President:**  
Rebecca Davies, RM  
**Patron:**  
HRH The Princess Royal

2022-02599

**Preeclampsia and Severe Maternal Morbidity During the COVID-19 Pandemic: A Population-Based Cohort Study in Ontario, Canada.** Snelgrove JW, Simpson AN, Sutradhar R, et al (2022), JOGC [Journal of Obstetrics and Gynaecology Canada] 4 April 2022, online

#### Objective

Significant changes to the delivery obstetrical care that occurred with the onset of the COVID-19 pandemic may be associated with higher risks of adverse maternal outcomes. We evaluated preeclampsia/HELLP (hemolysis, elevated liver enzymes and low platelets) syndrome and composite severe maternal morbidity (SMM) among pregnant people who gave birth during the COVID-19 pandemic and compared these data with those of people who gave birth before the pandemic in Ontario, Canada.

#### Methods

This was a population-based, retrospective cohort study using linked administrative data sets from ICES. Data on pregnant people at  $\geq 20$  weeks gestation who gave birth between March 15, 2020, and September 30, 2021, were compared with those of pregnant people who gave birth within the same date range for the years 2015–2019. We used multivariable logistic regression to assess the effect of the pandemic period on the odds of preeclampsia/HELLP syndrome and composite SMM, adjusting for maternal baseline characteristics and comorbidities.

#### Results

There were no differences between the study periods in the adjusted odds ratios (aORs) for preeclampsia/HELLP syndrome among primiparous (aOR 1.00; 95% CI 0.91–1.11) and multiparous (aOR 0.94; 95% CI 0.81–1.09) patients and no differences for composite SMM (primiparous, aOR 1.00; 95% CI 0.95–1.05; multiparous, aOR 1.01; 95% CI 0.95–1.08).

#### Conclusion

Adverse maternal outcomes were not higher among pregnant people who gave birth during the first 18 months of the COVID-19 pandemic in Ontario, Canada, when compared with those who gave birth before the pandemic. (Author)

**Full URL:** <https://doi.org/10.1016/j.jogc.2022.03.008>

---

MIDIRS is part of RCM Information Services Limited which is a company incorporated in England and Wales under company no.11914882 with registered office at 10-18 Union Street, London SE1 1SZ  
RCM Information Services Limited is a subsidiary of The Royal College of Midwives



**The Royal College of Midwives Trust**  
10-18 Union Street  
London  
SE1 1SZ



**Open 24 hours a day, 7 days a week**  
T: 0300 303 0444  
F: +44 20 7312 3536  
E: [info@rcm.org.uk](mailto:info@rcm.org.uk)  
W: [www.rcm.org.uk](http://www.rcm.org.uk)



**Chief Executive:**  
Gill Walton,  
MA, PGDip, BSc Hons, RM  
**President:**  
Rebecca Davies, RM  
**Patron:**  
HRH The Princess Royal

**2022-02132**

**Evaluating depression and anxiety throughout pregnancy and after birth: impact of the COVID-19 pandemic.** Zhang CXW, Okeke JC, Levitan RD, et al (2022), American Journal of Obstetrics & Gynecology MFM vol 4, no 3, May 2022, 100605

#### BACKGROUND

The COVID-19 pandemic presents unique social, economic, and psychological challenges for individuals globally. Thus, women who are pregnant face unprecedented mental health challenges.

#### OBJECTIVE

We sought to determine the impact of the pandemic on perinatal depression and anxiety in a longitudinal pregnancy cohort. We hypothesized increased depression and anxiety scores in women during pregnancy and after birth in the pandemic at all time points.

#### STUDY DESIGN

Participants were enrolled in the Ontario Birth Study, a pregnancy cohort embedded in clinical care at Mount Sinai Hospital, Toronto, Canada. Perinatal depression and anxiety were assessed using the 2-Item Patient Health Questionnaire and 2-Item Generalized Anxiety Disorder Questionnaire in early pregnancy, whereas the Edinburgh Postnatal Depression Scale and 2-Item Generalized Anxiety Disorder Questionnaire were used in late pregnancy and after birth. Logistic regression models were created to examine the association of the pandemic with clinically elevated mental health scores in the prepandemic group vs pandemic group while adjusting for covariates.

#### RESULTS

A total of 1159 survey responses from 649 participants between March 1, 2019, and February 28, 2021, were used to conduct this study. Participants were assessed in early pregnancy (n=416), in late pregnancy (n=373), and after birth (n=370). Responses received on or before February 29, 2020, were considered the “prepandemic” responses, whereas responses after the aforementioned date were considered the “pandemic” responses. Mean rank scores of depression and anxiety were significantly higher in the pandemic group (P=.02 and P=.003, respectively) in the postpartum period. There was no significant association between pandemic time and antenatal scores. However, postnatally, mothers were 2.6 times more likely to score  $\geq 13$  on the Edinburgh Postnatal Depression Scale during the pandemic than before the pandemic (95% confidence interval, 1.2–5.7; P=.02). Adjustment for ethnicity and income strengthened this association as the odds ratio increased to 3.3 (95% confidence interval, 1.4–8.0; P=.007).

#### CONCLUSION

Pandemic-associated increases in depression and anxiety scores were confined to the postpartum period, highlighting a need for increased screening and interventions for perinatal mood and anxiety disorders postnatally as this pandemic continues. (Author)

**Full URL:** <https://doi.org/10.1016/j.ajogmf.2022.100605>

---

MIDIRS is part of RCM Information Services Limited which is a company incorporated in England and Wales under company no.11914882 with registered office at 10-18 Union Street, London SE1 1SZ  
RCM Information Services Limited is a subsidiary of The Royal College of Midwives



**The Royal College of Midwives Trust**  
10-18 Union Street  
London  
SE1 1SZ



**Open 24 hours a day, 7 days a week**  
T: 0300 303 0444  
F: +44 20 7312 3536  
E: [info@rcm.org.uk](mailto:info@rcm.org.uk)  
W: [www.rcm.org.uk](http://www.rcm.org.uk)



**Chief Executive:**  
Gill Walton,  
MA, PGDip, BSc Hons, RM  
**President:**  
Rebecca Davies, RM  
**Patron:**  
HRH The Princess Royal

## 2022-02123

**The impact of the coronavirus pandemic curfew on the psychosocial lives of pregnant women in Jordan.** Qudsieh S, Mahfouz IA, Qudsieh H, et al (2022), *Midwifery* vol 109, June 2022, 103317

### Objectives

Worldwide the COVID-19 pandemic has negatively affected the health and psychosocial lives of people. International guidelines recommend special attention to pregnant women during pandemics and national emergencies. This study aimed to report the impact of the COVID-19 pandemic curfew on the psychosocial lives of pregnant women in Jordan.

### Design

A cross-sectional study was conducted and included women who were pregnant during the COVID-19 curfew in Jordan, which took place between mid-March and mid-June of 2020.

### Settings

A web-based survey that was posted on various social media platforms.

### Participants

Women who at the time of the study were 18 years of age or more, were living in Jordan, and were pregnant during the curfew.

### Measurements and findings

Data collected included women's characteristics, the impact of the curfew on the pregnancy, physical activity, and psychosocial lives and the barriers to seeking healthcare, in addition to pregnancy and delivery details, and changes in nutrition and supplements intake.

A total of 877 women responded to the survey. The results showed that 21.1% of the respondents did not receive any antenatal care (ANC) during the curfew. The respondents also reported that the main barriers for seeking ANC included healthcare facilities being closed (85.2%), the need for travel permits (76.8%), financial difficulties (63.9%), and fear of catching the COVID-19 virus (60.1%). Furthermore, 93.3% reported that they had psychological stress, and 29.9% reported that they had at least one form of domestic violence. Statistically significant associations existed between various women's characteristics, obstetric, psychosocial factors, and the level of psychological stress.

### Key conclusions

The COVID-19 pandemic curfew, which was applied in Jordan, resulted in a negative impact on the psychosocial lives of pregnant women. As a result, pregnant women did not receive optimal antenatal care and experienced higher degrees of psychological stress and domestic violence.

### Implications for practice

The findings of our study may encourage national healthcare policymakers to ensure the provision of appropriate psychosocial support of pregnant women during large scale emergencies. (Author)

**Full URL:** <https://doi.org/10.1016/j.midw.2022.103317>

---

MIDIRS is part of RCM Information Services Limited which is a company incorporated in England and Wales under company no.11914882 with registered office at 10-18 Union Street, London SE1 1SZ  
RCM Information Services Limited is a subsidiary of The Royal College of Midwives



**The Royal College of Midwives Trust**  
10-18 Union Street  
London  
SE1 1SZ



**Open 24 hours a day, 7 days a week**  
T: 0300 303 0444  
F: +44 20 7312 3536  
E: [info@rcm.org.uk](mailto:info@rcm.org.uk)  
W: [www.rcm.org.uk](http://www.rcm.org.uk)



**Chief Executive:**  
Gill Walton,  
MA, PGDip, BSc Hons, RM  
**President:**  
Rebecca Davies, RM  
**Patron:**  
HRH The Princess Royal

**2022-02118**

**“It’s always hard being a mom, but the pandemic has made everything harder”: A qualitative exploration of the experiences of perinatal women during the COVID-19 pandemic.** Kinser P, Jallo N, Moyer S, et al (2022), Midwifery vol 109, June 2022, 103313

#### Background

Understanding the psychosocial impacts of the COVID-19 pandemic in vulnerable groups, such as pregnant and parenting women, is a critical research and clinical imperative. Although many survey-based perinatal health studies have contributed important information about mental health, few have given full voice about the experiences of pregnant and postpartum women during the prolonged worldwide pandemic using a qualitative approach.

#### Objective

The purpose of this study is to explore the lived experience of pregnant and postpartum women in the United States during the ongoing COVID-19 pandemic.

#### Design

Qualitative phenomenological study.

#### Setting

This study was conducted in the community, by recruiting women throughout the U.S.

#### Participants

Fifty-four pregnant and postpartum women participated in qualitative interviews.

#### Methods

Data from one-on-one semi-structured interviews were analyzed using a team-based phenomenological qualitative approach.

#### Results

Two key themes were apparent: the pandemic has shined a light on the many typical struggles of motherhood; and, there is a lack of consistent, community-based or healthcare system resources available to address the complex needs of pregnant and postpartum women, both in general and during the pandemic.

#### Conclusions

Going forward, as the world continues to deal with the current pandemic and possible future global health crises, health care systems and providers are encouraged to consider the suggestions provided by these participants: talk early and often to women about mental health; help pregnant and postpartum women create and institute a personal plan for early support of their mental health needs and create an easily accessible mental health network; conceptualize practice methods that enhance coping and resilience; practice in community-based and interdisciplinary teams (e.g., midwives, doulas, perinatal social workers/ psychotherapists) to ensure continuity of care and to foster relationships between providers and pregnant/postpartum women; and consider learning from other countries’ successful perinatal healthcare practices.

#### Registration

Number (& date of first recruitment): not applicable.

#### Tweetable abstract

Pregnant and postpartum women insist that mental health care must be overhauled, stating the pandemic has highlighted inherent cracks in the system. (Author)

MIDIRS is part of RCM Information Services Limited which is a company incorporated in England and Wales under company no.11914882 with registered office at 10-18 Union Street, London SE1 1SZ  
RCM Information Services Limited is a subsidiary of The Royal College of Midwives



**The Royal College of Midwives Trust**  
10-18 Union Street  
London  
SE1 1SZ



**Open 24 hours a day, 7 days a week**  
T: 0300 303 0444  
F: +44 20 7312 3536  
E: [info@rcm.org.uk](mailto:info@rcm.org.uk)  
W: [www.rcm.org.uk](http://www.rcm.org.uk)



**Chief Executive:**  
Gill Walton,  
MA, PGDip, BSc Hons, RM  
**President:**  
Rebecca Davies, RM  
**Patron:**  
HRH The Princess Royal

## 2022-02117

**Exploring lived experiences of informal caregivers for pregnant women seeking scheduled antenatal care during the COVID-19 lockdown in China: A phenomenological study.** Zuo Y, Luo B-R, Wang L-N, et al (2022), Midwifery vol 109, June 2022, 103316

### Objective

We aimed to explore the lived experiences of informal caregivers for pregnant women seeking scheduled antenatal care during the early stage of China's COVID-19 lockdown and potential measures to address the challenges.

### Design

This is a phenomenological qualitative study.

### Setting

The study was carried out in a leading teaching hospital in Southwest China.

### Participants

We recruited 15 informal caregivers for healthy pregnant women on routine antenatal visits about six months after China launched the city-wide lockdown and other control measures for COVID-19, including 10 males and 5 females with diverse demographic backgrounds.

### Measures and findings

The research team developed a demographic form and an interview outline with key questions, conducted semi-structured interviews with the informal caregivers, and analyzed the data using the Colaizzi's method. Five themes of lived experiences were revealed, i.e., increased caregiving burdens, disruption of routines in family life, lack of accurate information and knowledge, active role adjustment, and positive attitudes and coping in a difficult time. Some caregivers reacted positively to the lockdown experience and saw it as an opportunity to rethink their lives and improve family relations.

### Key conclusions

The informal caregivers experienced increased physical and psychological burdens. Strategies such as adoption of a less frequent prenatal visit schedule, use of tele-medicine technologies, and provision of accurate information and knowledge may help to ease the increased informal caregiving burdens. Psychological counseling, community services and disaster response policies specially targeting pregnant women and their informal caregivers may also be valuable resources.

### Implications for practice

Attention should be drawn to the group of informal caregivers for pregnant women during a COVID-19 lockdown, including professional assistance delivered by nursing and other related professionals. Measures are called for to minimize exposure opportunities such as adoption of a new prenatal care schedule and tele-medicine technologies. Patient education with reliable information should be provided, preferably by nursing staff and physicians. Social support efforts including professional mental counseling may added and work with other resources such as community services and policy makers. (Author)



**2022-02095**

**Influence of the COVID-19 outbreak on transportation of pregnant women in an emergency medical service system:**

**Population-based, ORION registry.** Ota K, Nishioka D, Katayama Y, et al (2022), International Journal of Gynecology & Obstetrics vol 157, no 2, May 2022, pp 366-374

**Background**

The coronavirus disease 2019 (COVID-19), caused by Severe Acute Respiratory Syndrome Coronavirus 2, has spread rapidly across the world.

**Objective**

To assess the influence of the COVID-19 pandemic on the emergency medical service (EMS) for transportation of pregnant women by ambulance.

**Methods**

This study was a retrospective, descriptive study using the Osaka Emergency Information Research Intelligent Operation Network system, and included pregnant women transported by ambulance in Osaka Prefecture between January 1, 2018 and December 31, 2020. The main outcome of the study was difficulty in obtaining hospital acceptance for transfer of patients (difficult-to-transfer cases). We calculated the rates of difficult-to-transfer cases using univariate and multivariate analyses.

**Results**

Of the 1 346 457 total patients transported to hospitals by ambulance in Osaka Prefecture during the study period, pregnant women accounted for 2586 (909, 943, and 734, in 2018, 2019, and 2020, respectively). Logistic regression analysis revealed that pregnant women were negatively associated with difficult-to-transfer cases (adjusted OR 0.36, 95% CI 0.26–0.50). Compared with 2018, 2020 was significantly associated with difficult-to-transfer cases (adjusted OR 1.27, 95% CI 1.24–1.30).

**Conclusion**

Pregnant women were consistently associated with reduced odds for being difficult-to-transfer cases. The COVID-19 pandemic might have influenced difficult-to-transfer cases in 2020. (Author)

---

MIDIRS is part of RCM Information Services Limited which is a company incorporated in England and Wales under company no.11914882 with registered office at 10-18 Union Street, London SE1 1SZ  
RCM Information Services Limited is a subsidiary of The Royal College of Midwives



**The Royal College of Midwives Trust**  
10-18 Union Street  
London  
SE1 1SZ



**Open 24 hours a day, 7 days a week**  
T: 0300 303 0444  
F: +44 20 7312 3536  
E: [info@rcm.org.uk](mailto:info@rcm.org.uk)  
W: [www.rcm.org.uk](http://www.rcm.org.uk)



**Chief Executive:**  
Gill Walton,  
MA, PGDip, BSc Hons, RM  
**President:**  
Rebecca Davies, RM  
**Patron:**  
HRH The Princess Royal

## 2022-02092

**Establishing communication with relatives of admitted obstetrical patients with COVID-19 infection during COVID-19 pandemic: A quality improvement initiative.** Rana A, Sharma KA, Kulshrestha S, et al (2022), International Journal of Gynecology & Obstetrics 6 February 2022, online

### Objectives

To establish communication with relatives of obstetrical patients with coronavirus disease 2019 (COVID-19) admitted to an isolation ward by systematic use of quality improvement tools during the COVID-19 pandemic as there were many challenges in communicating with relatives.

### Methods

The study was conducted in the Department of Obstetrics and Gynecology at a tertiary-care teaching hospital based on four systematic steps of Point of Care Improvement methodology. After identifying the problem, a quality improvement team was constituted, which formed a specific aim. After root-cause analysis with fishbone tool, three Plan-Do-Study-Act (PDSA) cycles with various interventions were planned.

### Results

The outcome was measured as percentage of relatives of obstetrical patients admitted to the hospital with COVID-19 who were counseled about vital patient-related information. The baseline percentage of counseling of relatives of COVID-19-positive obstetrical patients admitted to the hospital was 14% per day. After three PDSA cycles, the target of 66.5% was achieved.

### Conclusion

Communication with the relatives of COVID-19-positive obstetrical patients admitted to isolation wards in the hospital could be easily streamlined without any additional resources using the principles of quality improvement during the COVID-19 pandemic.

### Synopsis

SynopsisThe contents of this page will be used as part of issue TOC only. It will not be published as part of main article.

This quality improvement project establishes that communication with the relatives of isolated COVID-19-positive obstetrical patients could be easily streamlined without any additional resources. (Author)

---

## 2022-02087

**Teenage motherhood in Africa: The epidemic in the COVID-19 pandemic.** Molek K, Bellizzi S (2022), International Journal of Gynecology & Obstetrics 15 February 2022, online

Efforts are urgently needed to increase access to sexual education and contraception, which has been proven to be effective in other contexts during the COVID-19 pandemic. (Author)

---

## 2022-02081

**Maternal and neonatal outcomes of COVID-19 co-infection in pregnant women with chronic hepatitis B virus infection: A prospective cohort study.** Rajan M, Sachan S, Abhinay A, et al (2022), International Journal of Gynecology & Obstetrics 25 February 2022, online

Co-infection with COVID-19 in pregnant women with pre-existing HBV infection led to a higher proportion of preterm deliveries and lower mean birth weight. (Author)

---

MIDIRS is part of RCM Information Services Limited which is a company incorporated in England and Wales under company no.11914882 with registered office at 10-18 Union Street, London SE1 1SZ  
RCM Information Services Limited is a subsidiary of The Royal College of Midwives



**The Royal College of Midwives Trust**  
10-18 Union Street  
London  
SE1 1SZ



**Open 24 hours a day, 7 days a week**  
T: 0300 303 0444  
F: +44 20 7312 3536  
E: [info@rcm.org.uk](mailto:info@rcm.org.uk)  
W: [www.rcm.org.uk](http://www.rcm.org.uk)



**Chief Executive:**  
Gill Walton,  
MA, PGDip, BSc Hons, RM  
**President:**  
Rebecca Davies, RM  
**Patron:**  
HRH The Princess Royal



**2022-02002**

**Telehealth multidisciplinary prenatal consultation during the COVID-19 pandemic: enhancing patient care coordination while maintaining high provider satisfaction.** Hargis-Villanueva A, Lai K, van Leeuwen KM, et al (2022), Journal of Maternal-Fetal and Neonatal Medicine 25 March 2022, online

#### Objective

Comprehensive fetal care centers address congenital anomalies by developing pre- and post-natal care plans in a multidisciplinary format. To reduce exposure during the Coronavirus Infectious Disease-2019 (COVID-19) pandemic, the Centers for Medicare & Medicaid Services (CMS) broadened access to telehealth services. We assessed provider satisfaction with the rapid transition from in-person prenatal visits to multidisciplinary consultations via telehealth as an adaptive response to the pandemic.

#### Methods

Patients referred to an urban academic fetal care center during the first 6 weeks of the COVID-19 pandemic underwent advanced imaging including fetal MRI, focused ultrasound, and fetal echocardiography. Subsequently, multidisciplinary telehealth consultations occurred with all providers attending virtually. Patients were given the option of attending the multidisciplinary telehealth consultation in a conference room in the hospital or from home. During these meetings, relevant images were reviewed with all participants via screen sharing through a secure video platform. Provider satisfaction with the telehealth paradigm was assessed using an electronic survey.

#### Results

Twenty-two surveys were administered with a response rate of 82%. 89% of providers were highly satisfied with the telehealth format. 72% of providers would prefer the multidisciplinary telehealth format to an in-person visit for future visits after COVID-19 restrictions are lifted. 22% of providers would leave the choice to the patient's family. One provider preferred in-person visits. Some providers noted that virtual conferences limited the ability to draw pictures, show educational materials, and provide emotional support.

#### Conclusion

Providers were overwhelmingly supportive of continuing multidisciplinary telehealth conferences for complex prenatal consultations, even after restrictions are lifted, which has led to the continuation of this model for the duration of the pandemic. Providers highlighted the convenience and improved care coordination across specialties. Further studies to examine the patient experience with virtual consultations are warranted. (Author)

---

MIDIRS is part of RCM Information Services Limited which is a company incorporated in England and Wales under company no.11914882 with registered office at 10-18 Union Street, London SE1 1SZ  
RCM Information Services Limited is a subsidiary of The Royal College of Midwives



**The Royal College of Midwives Trust**  
10-18 Union Street  
London  
SE1 1SZ



**Open 24 hours a day, 7 days a week**  
T: 0300 303 0444  
F: +44 20 7312 3536  
E: [info@rcm.org.uk](mailto:info@rcm.org.uk)  
W: [www.rcm.org.uk](http://www.rcm.org.uk)



**Chief Executive:**  
Gill Walton,  
MA, PGDip, BSc Hons, RM  
**President:**  
Rebecca Davies, RM  
**Patron:**  
HRH The Princess Royal

**2022-01974**

**Health disparities, COVID-19, and maternal and childbirth outcomes: a meta-epidemiological study of equity reporting in systematic reviews.** Hartwell M, Lin V, Gatewood AC, et al (2022), Journal of Maternal-Fetal and Neonatal Medicine 13 March 2022, online

#### Background

Pregnant women with COVID-19 are at increased risk for adverse maternal and pregnancy outcomes, and birth complications. Given the health outcome disparities among pregnant women of racial and ethnic minorities and the reliance of medical practice on systematic reviews and meta-analyses (SRMAs)—as they are the apical component in the hierarchy of evidence in medical research—the primary objective of the study is to examine the inclusion of the equity reporting in SRMAs focused on pregnancy outcomes and COVID-19 using PROGRESS-Plus equity framework. PROGRESS represents equity measures of Place, Race, Occupation, Gender, Religion, Education, Social capital, and Socio-economic status.

#### Methods

We conducted a systematic search of three databases to identify SRMAs related to maternal and pregnancy outcomes related to COVID-19. We extracted whether SRMAs reported or analyzed PROGRESS-Plus components among other study characteristics.

#### Results

Nearly 85% of SRMAs did not include any equity items to account for racial or geographic disparities. Reporting of race was absent from 95% of the studies. Place was the most common PROGRESS item and maternal age was the most common PROGRESS-Plus item reported overall.

#### Conclusion

When research is performed and reported in a way that fails to address disparities, the downstream repercussions may include medical care in the form of new protocol-driven hospital management, pharmacologic interventions, and other treatment options that mirror this absence in reporting. The absence of adequate reporting widens gaps in health outcomes among at-risk groups, such as pregnant women of racial and ethnic minorities. (Author)

---

MIDIRS is part of RCM Information Services Limited which is a company incorporated in England and Wales under company no.11914882 with registered office at 10-18 Union Street, London SE1 1SZ  
RCM Information Services Limited is a subsidiary of The Royal College of Midwives



**The Royal College of Midwives Trust**  
10-18 Union Street  
London  
SE1 1SZ



**Open 24 hours a day, 7 days a week**  
T: 0300 303 0444  
F: +44 20 7312 3536  
E: [info@rcm.org.uk](mailto:info@rcm.org.uk)  
W: [www.rcm.org.uk](http://www.rcm.org.uk)



**Chief Executive:**  
Gill Walton,  
MA, PGDip, BSc Hons, RM  
**President:**  
Rebecca Davies, RM  
**Patron:**  
HRH The Princess Royal

## 2022-01964

**Obstetric and perinatal outcomes of pregnancies with COVID 19: a systematic review and meta-analysis.** Pérez-López FR, Savirón-Cornudella R, Chedraui P, et al (2022), Journal of Maternal-Fetal and Neonatal Medicine 13 March 2022, online

### Objective

This meta-analysis aimed at comparing obstetric and perinatal outcomes in laboratory-tested pregnant women for severe acute respiratory syndrome coronavirus 2 (SARS-CoV-2) infection before delivering.

### Method

We performed a comprehensive systematic review of electronic databases for studies reporting pregnant women with and without SARS-CoV-2 infection, as determined by polymerase chain reaction (PCR) before delivery, during the pandemic period published up to June 25, 2021. Results are reported as mean difference (MD) or odds ratio (OR) and their 95% confidence interval (CI).

### Results

Seventeen observational studies with low to moderate risk of bias, reported on 2,769 pregnant women with a positive SARS-CoV-2 PCR test and 13,807 with a negative test. Pregnant women with a positive PCR test delivered at an earlier gestational age (MD -0.19; 95% CI -0.36 to -0.02 weeks), smoked less (OR 0.75; 95% CI 0.61–0.94) and were associated with higher odds for preeclampsia (OR 1.30; 95% CI 1.09–1.54), NICU admissions (OR 2.37; 95% CI 1.18–4.76), stillbirths (OR 2.70; 95% CI, 1.38–5.29), and perinatal mortality (OR 3.23; 95% CI 1.23–8.52). There were no significant differences between positive and negative tested women in terms of nulliparity, multiple pregnancies, gestational diabetes, route of delivery, labor induction, preterm birth, infant birth weight, 5 min Apgar scores < 7, small-for-gestational-age infants and fetal malformations. Eleven studies included neonatal PCR SARS-CoV-2 testing which was performed on 129 infants, of which 20 were positive.

### Conclusion

Positive SARS-CoV-2 tested pregnant women had higher odds for preeclampsia/hypertensive disorders of pregnancy, NICU admissions, stillbirths and perinatal mortality. (Author)

---

## 2022-01961

**Antenatal classes [written answer].** Scottish Parliament (2022), Official Report Written question S6W-07397, 17 March 2022

Maree Todd responds to a written question asked by Monica Lennon to the Scottish Government regarding which NHS boards are currently providing in-person antenatal classes. This question is asked further to answer S6W-06381 by Hamza Yousaf on 2 March 2022. (LDO)

**Full URL:** <https://archive2021.parliament.scot/parliamentarybusiness/28877.aspx?SearchType=Advance&ReferenceNumbers=S6W-07397>

---

## 2022-01937

**Antenatal classes [written answer].** Scottish Parliament (2022), Official Report Written question S6W-07219, 10 March 2022

Humza Yousaf responds to a written question asked by Monica Lennon to the Scottish Government regarding the date by which face-to-face antenatal classes that are run by NHS Scotland will resume, in light of the reported resumption of private classes. (MB)

**Full URL:** <https://archive2021.parliament.scot/parliamentarybusiness/28877.aspx?SearchType=Advance&ReferenceNumbers=S6W-07219>

---

MIDIRS is part of RCM Information Services Limited which is a company incorporated in England and Wales under company no.11914882 with registered office at 10-18 Union Street, London SE1 1SZ  
RCM Information Services Limited is a subsidiary of The Royal College of Midwives



**The Royal College of Midwives Trust**  
10-18 Union Street  
London  
SE1 1SZ



**Open 24 hours a day, 7 days a week**  
T: 0300 303 0444  
F: +44 20 7312 3536  
E: [info@rcm.org.uk](mailto:info@rcm.org.uk)  
W: [www.rcm.org.uk](http://www.rcm.org.uk)



**Chief Executive:**  
Gill Walton,  
MA, PGDip, BSc Hons, RM  
**President:**  
Rebecca Davies, RM  
**Patron:**  
HRH The Princess Royal

## 2022-01936

**Antenatal classes [written answer].** Scottish Parliament (2022), Official Report Written question S6W-07397, 17 March 2022  
Maree Todd responds to a written question asked by Monica Lennon to the Scottish Government further to the answer to question S6W-06381 by Humza Yousaf on 2 March 2022, regarding which NHS boards are currently providing in-person antenatal classes. (MB)

**Full URL:** <https://archive2021.parliament.scot/parliamentarybusiness/28877.aspx?SearchType=Advance&ReferenceNumbers=S6W-07397>

---

## 2022-01925

**SARS-CoV-2 Placentitis and Intraparenchymal Thrombohematomas Among COVID-19 Infections in Pregnancy.** Huynh A, Sehn JK, Goldfarb IT, et al (2022), JAMA Network Open vol 5, no 3, March 2022, e225345

This cases series examines SARS-CoV-2 placentitis and intraparenchymal thrombohematomas among COVID-19 infections during pregnancy. (Author)

**Full URL:** <https://doi.org/10.1001/jamanetworkopen.2022.5345>

---

## 2022-01923

**COVID-19 Cases and Disease Severity in Pregnancy and Neonatal Positivity Associated With Delta (B.1.617.2) and Omicron (B.1.1.529) Variant Predominance.** Adhikari EH, Macdonald L, SoRelle JA, et al (2022), JAMA (Journal of the American Medical Association) vol 327, no 15, 19 April 2022, pp 1500-1502

This study examines infections, illness severity, vaccinations, and early neonatal infections among obstetric patients during the pre-Delta, Delta, and Omicron periods of the COVID-19 pandemic. (Author)

**Full URL:** <https://doi.org/10.1001/jama.2022.4356>

---

MIDIRS is part of RCM Information Services Limited which is a company incorporated in England and Wales under company no.11914882 with registered office at 10-18 Union Street, London SE1 1SZ  
RCM Information Services Limited is a subsidiary of The Royal College of Midwives



**The Royal College of Midwives Trust**  
10-18 Union Street  
London  
SE1 1SZ



**Open 24 hours a day, 7 days a week**  
T: 0300 303 0444  
F: +44 20 7312 3536  
E: [info@rcm.org.uk](mailto:info@rcm.org.uk)  
W: [www.rcm.org.uk](http://www.rcm.org.uk)



**Chief Executive:**  
Gill Walton,  
MA, PGDip, BSc Hons, RM  
**President:**  
Rebecca Davies, RM  
**Patron:**  
HRH The Princess Royal

**2022-01705**

**Knowledge, attitude and practices of pregnant women related to COVID-19 infection: A cross-sectional survey in seven countries from the Global Network for Women's and Children's Health.** Naqvi F, Naqvi S, Billah SM, et al (2022), BJOG: An International Journal of Obstetrics and Gynaecology 14 February 2022, online

**Objective**

We sought to understand knowledge, attitudes and practices (KAP) regarding COVID-19 in pregnant women in seven low and middle-income countries (LMIC).

**Design**

Population-based prospective, observational study.

**Settings**

Study sites in DRC, Kenya, Zambia, Bangladesh, India (two sites), Pakistan and Guatemala.

**Population and sample**

Pregnant women in the Global Network's Maternal and Neonatal Health Registry (MNHR).

**Methods**

A KAP questionnaire was administered in face-to-face interviews with pregnant women from September 2020 through October 2021 in the MNHR.

**Main outcome measures**

KAP regarding COVID-19 during pregnancy.

**Results**

In all, 25 260 women completed the survey. Overall, 56.8% of women named  $\geq 3$  COVID-19 symptoms, 34.3% knew  $\geq 2$  transmission modes, 51.3% knew  $\geq 3$  preventive measures and 79.7% named at least one high-risk condition. Due to COVID-19 exposure concerns, 23.8% had avoided prenatal care and 7.5% planned to avoid hospital delivery. Over half the women in the Guatemalan site and 40% in the Pakistan site reduced care seeking due to COVID-19 exposure concerns. Of the women, 24.0% were afraid of getting COVID-19 from healthcare providers. Overall, 63.3% reported wearing a mask and 29.1% planned to stay at home to reduce COVID-19 exposure risk.

**Conclusions**

We found a decrease in planned antenatal and delivery care use due to COVID-19 concerns. The clinical implications of potential decreases in care are unclear, but decline in essential healthcare utilisation during pregnancy and delivery could pose challenges for maternal and newborn health. More research is needed to address the impact of COVID-19 on routine pregnancy and delivery care. (Author)

**Full URL:** <https://doi.org/10.1111/1471-0528.17122>

---

MIDIRS is part of RCM Information Services Limited which is a company incorporated in England and Wales under company no.11914882 with registered office at 10-18 Union Street, London SE1 1SZ  
RCM Information Services Limited is a subsidiary of The Royal College of Midwives



**The Royal College of Midwives Trust**  
10-18 Union Street  
London  
SE1 1SZ



**Open 24 hours a day, 7 days a week**  
T: 0300 303 0444  
F: +44 20 7312 3536  
E: [info@rcm.org.uk](mailto:info@rcm.org.uk)  
W: [www.rcm.org.uk](http://www.rcm.org.uk)



**Chief Executive:**  
Gill Walton,  
MA, PGDip, BSc Hons, RM  
**President:**  
Rebecca Davies, RM  
**Patron:**  
HRH The Princess Royal

**2022-01578**

**Management and implications of severe COVID-19 in pregnancy in the UK: data from the UK Obstetric Surveillance System national cohort.** Vousden N, Ramakrishnan R, Bunch K, et al (2022), *Acta Obstetrica et Gynecologica Scandinavica* vol 101, no 4, April 2022, pp 461-470

#### Introduction

There is a lack of population level data on risk factors and impact of severe COVID-19 in pregnancy. The aims of this study were to determine the characteristics, and maternal and perinatal outcomes associated with severe COVID-19 in pregnancy compared with those with mild and moderate COVID-19 and to explore the impact of timing of birth.

#### Material and methods

This was a secondary analysis of a national, prospective cohort study. All pregnant women admitted to hospital in the UK with symptomatic SARS-CoV-2 from March 1, 2020 to October 31, 2021 were included. The severity of maternal infection (need for high flow or invasive ventilation, intensive care admission or died), pregnancy and perinatal outcomes, and the impact of timing of birth were analyzed using multivariable logistic regression.

#### Results

Of 4436 pregnant women, 13.9% (n = 616) had severe infection. Women with severe infection were more likely to be aged  $\geq 30$  years (adjusted odds ratio [aOR] aged 30–39 1.48, 95% confidence interval [CI] 1.20–1.83), be overweight or obese (aOR 1.73, 95% CI 1.34–2.25 and aOR 2.52 95% CI 1.97–3.23, respectively), be of mixed ethnicity (aOR 1.93, 95% CI 1.17–3.21) or have gestational diabetes (aOR 1.43, 95% CI 1.09–1.87) compared with those with mild or moderate infection. Women with severe infection were more likely to have a pre-labor cesarean birth (aOR 8.84, 95% CI 6.61–11.83), a very or extreme preterm birth (28–31+ weeks' gestation, aOR 18.97, 95% CI 7.78–14.85; <28 weeks' gestation, aOR 12.35, 95% CI 6.34–24.05) and their babies were more likely to be stillborn (aOR 2.51, 95% CI 1.35–4.66) or admitted to a neonatal unit (aOR 11.61, 95% CI 9.28–14.52). Of 112 women with severe infection who were discharged and gave birth at a later admission, the majority gave birth  $\geq 36$  weeks (85.7%), noting that three women in this group (2.7%) had a stillbirth.

#### Conclusions

Severe COVID-19 in pregnancy increases the risk of adverse outcomes. Information to promote uptake of vaccination should specifically target those at greatest risk of severe outcomes. Decisions about timing of birth should be informed by multidisciplinary team discussion; however, our data suggest that women with severe infection who do not require early delivery have mostly good outcomes but that those with severe infection at term may warrant rapid delivery. (Author)

**Full URL:** <https://doi.org/10.1111/aogs.14329>

**2022-01480**

**Adverse Effects of COVID-19 on Perinatal Outcomes Globally.** Callister LC (2022), *MCN - American Journal of Maternal/Child Nursing* vol 47, no 2, March/April 2022, p 110

The COVID-19 pandemic has adversely affected outcomes for childbearing women and their babies around the globe, especially in low resource, low-to-middle income countries. Our global health and nursing expert, Dr. Callister, reviews some of the most recently published articles on the impact of COVID-19 on this vulnerable population. (Author)

MIDIRS is part of RCM Information Services Limited which is a company incorporated in England and Wales under company no.11914882 with registered office at 10-18 Union Street, London SE1 1SZ  
RCM Information Services Limited is a subsidiary of The Royal College of Midwives



**The Royal College of Midwives Trust**  
10-18 Union Street  
London  
SE1 1SZ



**Open 24 hours a day, 7 days a week**  
T: 0300 303 0444  
F: +44 20 7312 3536  
E: [info@rcm.org.uk](mailto:info@rcm.org.uk)  
W: [www.rcm.org.uk](http://www.rcm.org.uk)



**Chief Executive:**  
Gill Walton,  
MA, PGDip, BSc Hons, RM  
**President:**  
Rebecca Davies, RM  
**Patron:**  
HRH The Princess Royal

## 2022-01460

**Changes in rates of adverse pregnancy outcomes during the COVID-19 pandemic: a cross-sectional study in the United States, 2019–2020.** Simeone RM, Downing KF, Wallace B, et al (2022), *Journal of Perinatology* vol 42, no 5, May 2022, pp 617-623

### Objective

Our objective was to assess differences in pregnancy outcomes during the COVID-19 pandemic compared to the previous year.

### Study design

In a cross-sectional study of delivery hospitalizations in the Premier Healthcare Database Special COVID-19 Release, we assessed differences in selected maternal and pregnancy outcomes occurring April–December in 2019 and 2020 in the United States.

### Result

Among 663,620 deliveries occurring in 2019 and 614,093 deliveries occurring in 2020, we observed an increase in in-hospital maternal death from 2019 to 2020, which was no longer statistically significant after excluding deliveries with a COVID-19 diagnosis. Intensive care unit admission and preterm birth decreased from 2019 to 2020. There was no difference in the prevalence of most other outcomes examined.

### Conclusion

The full impact of the COVID-19 pandemic on maternal and pregnancy outcomes remains to be understood. Most outcomes investigated experienced minimal change from 2019 to 2020. (Author)

Full URL: <https://doi.org/10.1038/s41372-022-01327-3>

---

## 2022-01385

**A Mixed-Methods Study of Experiences During Pregnancy Among Black Women During the COVID-19 Pandemic.**

Dove-Medows E, Davis J, McCracken L, et al (2022), *The Journal of Perinatal and Neonatal Nursing* vol 36, no 2, April/June 2022, pp 161-172

Pregnant women experienced disruptions in their prenatal care during the coronavirus disease-2019 (COVID-19) pandemic. While there is emerging research about the impact of COVID-19 on experiences of pregnancy, the majority of studies that have reported on prenatal care and birth during COVID-19 have not incorporated the first-person accounts of Black women. The purpose of this mixed-methods study was to explore the perspectives of Black women on prenatal care, labor, and birth during the pandemic. A total of 33 participants completed questionnaires. Fourteen of these 33 women and an additional 2 participated in qualitative interviews. Descriptive statistics and a mixed-methods analysis were employed. Participants expressed disappointment about disruptions in their experiences of pregnancy including the way their prenatal care was experienced, cancellation of planned “rites of passage,” and visitor policy restrictions during and after the birth. Forty-five percent of participants reported being worried about getting COVID-19 and (61%) about their infant getting COVID-19. Many participants experienced a sense of loss that may permeate through other aspects of their lives. Providing extra support and points of contact can help lessen feelings of isolation during the pandemic and can also offer more explanation for rapidly changing policies and procedures. (Author)

---

MIDIRS is part of RCM Information Services Limited which is a company incorporated in England and Wales under company no.11914882 with registered office at 10-18 Union Street, London SE1 1SZ  
RCM Information Services Limited is a subsidiary of The Royal College of Midwives



**The Royal College of Midwives Trust**  
10-18 Union Street  
London  
SE1 1SZ



**Open 24 hours a day, 7 days a week**  
T: 0300 303 0444  
F: +44 20 7312 3536  
E: [info@rcm.org.uk](mailto:info@rcm.org.uk)  
W: [www.rcm.org.uk](http://www.rcm.org.uk)



**Chief Executive:**  
Gill Walton,  
MA, PGDip, BSc Hons, RM  
**President:**  
Rebecca Davies, RM  
**Patron:**  
HRH The Princess Royal

**2022-01352**

**Tailoring the response to COVID-19: experiences of an inner city maternity unit with a virtual patient surveillance approach.**

Elsmore A, Redjepova O, Wright J, et al (2022), Journal of Obstetrics and Gynaecology 18 February 2022, online

This study outlines the characteristics and outcomes of pregnant women with severe acute respiratory syndrome coronavirus 2 (SARS-CoV-2). We present the success of our 'COVID Surveillance Team' – a dedicated team of midwives and medics that regularly contact patients, identifying early any need for escalation of care. Data were collected prospectively from March to September 2020. Patients are followed up by our team for 14 days following diagnosis, via telephone. Maternal and neonatal outcomes were studied. Fifty-five women were diagnosed with COVID-19. 39/55 (70.9%) were of BAME background. 10/55 (18%) had medical comorbidities. 35/55 (63.6%) were diagnosed in the 3rd trimester, 7/55 (12.7%) were postnatal. Three women (5.4%) required critical care unit admission. One was transferred to a tertiary centre for extra-corporeal membrane oxygenation (ECMO), one patient died. Of the 43 deliveries, 28 (65.1%) underwent caesarean section. 11/45 (24.4%) of babies were born preterm. Eight babies were tested for COVID-19, one was positive. Our study demonstrates most pregnant women suffer mild illness, with no adverse outcomes for mother or neonate. We highlight the success of our COVID surveillance team, that should be considered best practice and consideration should be given for adoption by other maternity units to enhance patient safety.

**Impact Statement**

What is already known on this subject? Many studies present maternal and neonatal characteristics and outcomes of COVID-19 infection in the pregnant population, demonstrating most patients suffer mild disease with minimal adverse outcomes.

What do the results of this study add? We highlight the important work of our COVID surveillance team, and the positive impact it has had on the wellbeing and safety of our women. We believe we are the first maternity unit in the UK to adopt and report on a virtual patient surveillance approach supporting our patients and leading to increased patient safety.

What are the implications of these findings for clinical practice and/or further research? Our surveillance team has proved very successful and has been described as exemplar by NHS England and promoted as best practice. This approach could be adopted by other units, within the UK and further afield, for the benefit of women's health and safety. The team has shared their guidance and standard operating procedure with maternity units across the UK. (Author)

---

MIDIRS is part of RCM Information Services Limited which is a company incorporated in England and Wales under company no.11914882 with registered office at 10-18 Union Street, London SE1 1SZ  
RCM Information Services Limited is a subsidiary of The Royal College of Midwives



**The Royal College of Midwives Trust**  
10-18 Union Street  
London  
SE1 1SZ



**Open 24 hours a day, 7 days a week**  
T: 0300 303 0444  
F: +44 20 7312 3536  
E: [info@rcm.org.uk](mailto:info@rcm.org.uk)  
W: [www.rcm.org.uk](http://www.rcm.org.uk)



**Chief Executive:**  
Gill Walton,  
MA, PGDip, BSc Hons, RM  
**President:**  
Rebecca Davies, RM  
**Patron:**  
HRH The Princess Royal



**2022-01298**

**Indonesian midwives' perspectives on changes in the provision of maternity care during the COVID-19 pandemic: A qualitative study.** Hazfiarini A, Zahroh RI, Akter S, et al (2022), *Midwifery* vol 108, May 2022, 103291

Objective

To explore how COVID-19 influenced the provision of high-quality maternity care in Indonesia.

Design and methods

A qualitative descriptive study using in-depth interviews was undertaken. Thematic analysis was used to analyse data, and behaviour change frameworks (Theoretical Domain Framework (TDF) and Capability, Opportunity, and Motivation (COM-B)) were used to identify and map facilitators and barriers influencing maternity care provision during the COVID-19 pandemic.

Setting and participants

Fifteen midwives working in community maternity care facilities in Surabaya and Mataram, Indonesia were included. Surabaya is in western Indonesia, with around 56,000 births per year and a population of around 3 million. Mataram is in eastern Indonesia, with around 7,000 births per year and a population of around 500,000.

Findings

The main changes to maternity care provision during the COVID-19 pandemic were reduced frequency of antenatal and postpartum care visits, reduced support for women, including unavailability of maternity care and reduced number of antenatal care and labour companions, changes in location of provision of care, and public health changes related to COVID-19. The main factors influencing the provision of high-quality maternity care during the COVID-19 pandemic were behavioural regulation, professional role and identity, and environmental context and resources.

Key conclusions and implications for practice

Maternity care provision underwent substantial changes during the COVID-19 pandemic in Indonesia. Findings from this study can contribute to better understanding of how maternity care provision changed during the pandemic, and how positive changes can be reinforced, and negative changes can be addressed. (Author)

Full URL: <https://doi.org/10.1016/j.midw.2022.103291>

---

**2022-01284**

**Increasing oxygen requirements and disease severity in pregnant individuals with the SARS-CoV-2 Delta variant.** Eid J, Abdelwahab M, Caplan M, et al (2022), *American Journal of Obstetrics & Gynecology MFM* vol 4, no 3, May 2022, 100612

Correspondence piece aiming to determine whether the Delta variant of SARS-CoV-2 is associated with increased oxygen requirement and disease severity among pregnant women. Results suggest that patients with the Delta variant were more likely to require oxygen supplementation, have more severe disease and require admission to the intensive care unit. (LDO)

Full URL: <https://doi.org/10.1016/j.ajogmf.2022.100612>

---

**2022-01276**

**Pregnancy: Employment [written answer].** House of Commons (2022), Hansard Written question 117055, 2 February 2022

Maria Caulfield responds to a written question from Annelise Dodds to the Secretary of State for Health and Social Care, regarding whether his Department intends to publish guidance for (a) employers and (b) pregnant workers on new and expectant mothers in the workplace during covid-19. (MB)

Full URL: <https://questions-statements.parliament.uk/written-questions/detail/2022-02-02/117055>

---

MIDIRS is part of RCM Information Services Limited which is a company incorporated in England and Wales under company no.11914882 with registered office at 10-18 Union Street, London SE1 1SZ  
RCM Information Services Limited is a subsidiary of The Royal College of Midwives



**The Royal College of Midwives Trust**  
10-18 Union Street  
London  
SE1 1SZ



**Open 24 hours a day, 7 days a week**  
T: 0300 303 0444  
F: +44 20 7312 3536  
E: [info@rcm.org.uk](mailto:info@rcm.org.uk)  
W: [www.rcm.org.uk](http://www.rcm.org.uk)



**Chief Executive:**  
Gill Walton,  
MA, PGDip, BSc Hons, RM  
**President:**  
Rebecca Davies, RM  
**Patron:**  
HRH The Princess Royal

**2022-01259**

**Influence of the COVID-19 pandemic on abortions and births in Sweden: a mixed-methods study.** Rydelius J, Edalat M, Nyman V, et al (2022), *BMJ Open* vol 12, no 2, February 2022, e054076

Introduction Although considered an essential service by the WHO, there are indications that access to induced abortion care has been restricted during the COVID-19 pandemic.

Objectives To investigate if the number of induced abortions and ongoing pregnancies changed during the first pandemic wave of COVID-19 in 2020 compared with recent years prior to the pandemic and explore possible reasons for the findings.

Design Convergent parallel mixed-methods design. Collection of quantitative data from the Swedish National Board of Health and Welfare and the Swedish Pregnancy Register, and qualitative data from interviews.

Setting and time period National data on abortions (January 2018–June 2020) and births (January 2018–March 2021). Interviews performed at the main abortion clinic, Gothenburg, Sweden, in June 2020.

Participants All women aged 15–44 years living in Sweden 2018–2020, approximately 1.9 million. 15 women who sought abortion were interviewed.

Primary and secondary outcome measures Number of abortions and births/1000 women aged 15–44 years. Themes and subthemes identified from interviews.

Results The number of abortions and ongoing pregnancies did not change significantly during the study period compared with before the pandemic started. Interview themes identified were the following: meeting with abortion care during the COVID-19 pandemic (availability, and fear of being infected and infecting others); and the impact of the COVID-19 pandemic on the abortion decision (to catch COVID-19 during pregnancy, feelings of loneliness and isolation, and social aspects).

Conclusions This study shows that the number of abortions and ongoing pregnancies remained unchanged during the first wave of the COVID-19 pandemic in 2020 in Sweden compared with before the start of the pandemic. Abortion-seeking women did not hesitate to proceed with the abortion. The women expressed a number of fears concerning both availability of care and their health, which could have been properly addressed by the authorities. (Author)

**Full URL:** <http://dx.doi.org/10.1136/bmjopen-2021-054076>

---

MIDIRS is part of RCM Information Services Limited which is a company incorporated in England and Wales under company no.11914882 with registered office at 10-18 Union Street, London SE1 1SZ  
RCM Information Services Limited is a subsidiary of The Royal College of Midwives



**The Royal College of Midwives Trust**  
10-18 Union Street  
London  
SE1 1SZ



**Open 24 hours a day, 7 days a week**  
T: 0300 303 0444  
F: +44 20 7312 3536  
E: [info@rcm.org.uk](mailto:info@rcm.org.uk)  
W: [www.rcm.org.uk](http://www.rcm.org.uk)



**Chief Executive:**  
Gill Walton,  
MA, PGDip, BSc Hons, RM  
**President:**  
Rebecca Davies, RM  
**Patron:**  
HRH The Princess Royal

## 2022-01142

**Impacts and effects of COVID-19 infection in pregnancy.** Sunder A, Varghese B, Darwish B, et al (2022), Saudi Medical Journal vol 43, no 1, January 2022, pp 67-74

Objectives: To explore the trimester wise significance of the primary outcome in pregnant women during coronavirus disease-19 (COVID-19) pandemic.

Methods: Retrospective observational study of pregnant women who were infected with COVID-19 from April 2020 until March 2021 at Bahrain Defense Force Hospital, Riffa, Bahrain. The study focused on the effects in relation to gestational age (GA), association with variables, severity, and treatment. A p-value of  $\leq 0.05$  was considered significant.

Results: During the study period, 74 COVID-19 cases were identified from the recorded 2944 pregnant women. The mean GA at diagnosis was  $33.5 \pm 12.2$  weeks, and the mean GA at birth was  $38.4 \pm 1.8$  weeks. Analysis of the obstetric complications revealed fetal growth restriction (FGR) had a p-value of  $< 0.001$ . According to the trimester wise analysis, between the gestational period at diagnosis and the outcome of pregnancy, significant p-value of  $< 0.01$  was found in miscarriage. There were no significant associations found in GA at diagnosis and delivery, complications in relation to maternal age and body mass index, and no maternal morbidities or mortalities.

Conclusion: In our study, FGR and miscarriage were the identified complications. However, the maternal and neonatal end result of COVID-19 was satisfactory. (Author)

Full URL: <https://doi.org/10.15537/smj.2022.43.1.20210694>

---

## 2022-01110

**Home use of both pills for early medical abortion (EMA) up to 10 weeks gestation: summary of consultation responses.**

Department of Health and Social Care (2022), 10 March 2022

Presents the outcome of a consultation to ascertain opinion on whether to make permanent a temporary approval put in place by the Government during the coronavirus pandemic, to return to the pre-pandemic arrangements, or to extend the temporary approval by 12 months. The temporary approval allowed women in England to take both pills for an early medical abortion (EMA) in their own homes, up to 10 weeks' gestation, following a consultation with a clinician via the telephone or online, without the need to first attend a hospital or clinic. 70% of respondents indicated that they wish the temporary approval to end immediately, however some Health professional organisations, abortion providers and some other organisations supported the temporary measure becoming permanent. (JSM)

Full URL: <https://www.gov.uk/government/consultations/home-use-of-both-pills-for-early-medical-abortion/outcome/home-use-of-both-pills-for-early-medical-abortion-ema-up-to-10-weeks-gestation-summary-of-consultation-responses>

---

MIDIRS is part of RCM Information Services Limited which is a company incorporated in England and Wales under company no.11914882 with registered office at 10-18 Union Street, London SE1 1SZ  
RCM Information Services Limited is a subsidiary of The Royal College of Midwives



**The Royal College of Midwives Trust**  
10-18 Union Street  
London  
SE1 1SZ



**Open 24 hours a day, 7 days a week**  
T: 0300 303 0444  
F: +44 20 7312 3536  
E: [info@rcm.org.uk](mailto:info@rcm.org.uk)  
W: [www.rcm.org.uk](http://www.rcm.org.uk)



**Chief Executive:**  
Gill Walton,  
MA, PGDip, BSc Hons, RM  
**President:**  
Rebecca Davies, RM  
**Patron:**  
HRH The Princess Royal

**2022-01104**

**COVID-19 in pregnancy—what study designs can we use to assess the risk of congenital anomalies in relation to COVID-19 disease, treatment and vaccination?**. Dolk H, Damase-Michel C, Morris JK, et al (2022), Paediatric and Perinatal Epidemiology 2 March 2022, online

Background

The COVID-19 pandemic has accelerated pregnancy outcome research, but little attention has been given specifically to the risk of congenital anomalies (CA) and first trimester exposures.

Objectives

We reviewed the main data sources and study designs used internationally, particularly in Europe, for CA research, and their strengths and limitations for investigating COVID-19 disease, medications and vaccines.

Population

We classify research designs based on four data sources: a) spontaneous adverse event reporting, where study subjects are positive for both exposure and outcome, b) pregnancy exposure registries, where study subjects are positive for exposure, c) congenital anomaly registries, where study subjects are positive for outcome and d) population healthcare data where the entire population of births is included, irrespective of exposure and outcome.

Study Design

Each data source allows different study designs, including case series, exposed pregnancy cohorts (with external comparator), ecological studies, case-control studies and population cohort studies (with internal comparator).

Methods

The quality of data sources for CA studies is reviewed in relation to criteria including diagnostic accuracy of CA data, size of study population, inclusion of terminations of pregnancy for foetal anomaly, inclusion of first trimester COVID-19-related exposures and use of an internal comparator group. Multinational collaboration models are reviewed.

Results

Pregnancy exposure registries have been the main design for COVID-19 pregnancy studies, but lack detail regarding first trimester exposures relevant to CA, or a suitable comparator group. CA registries present opportunities for improving diagnostic accuracy in COVID-19 research, especially when linked to other data sources. Availability of inpatient hospital medication use in population healthcare data is limited. More use of ongoing mother-baby linkage systems would improve research efficiency. Multinational collaboration delivers statistical power.

Conclusions

Challenges and opportunities exist to improve research on CA in relation to the COVID-19 pandemic and future pandemics.  
(Author)

Full URL: <https://doi.org/10.1111/ppe.12840>

---

MIDIRS is part of RCM Information Services Limited which is a company incorporated in England and Wales under company no.11914882 with registered office at 10-18 Union Street, London SE1 1SZ  
RCM Information Services Limited is a subsidiary of The Royal College of Midwives



**The Royal College of Midwives Trust**  
10-18 Union Street  
London  
SE1 1SZ



**Open 24 hours a day, 7 days a week**  
T: 0300 303 0444  
F: +44 20 7312 3536  
E: [info@rcm.org.uk](mailto:info@rcm.org.uk)  
W: [www.rcm.org.uk](http://www.rcm.org.uk)



**Chief Executive:**  
Gill Walton,  
MA, PGDip, BSc Hons, RM  
**President:**  
Rebecca Davies, RM  
**Patron:**  
HRH The Princess Royal

**2022-01010**

**The impact of Covid-19 restrictions on depressive symptoms in low-risk and high-risk pregnant women: a cross-sectional study before and during pandemic.** Smorti M, Gemignani A, Bonassi L, et al (2022), BMC Pregnancy and Childbirth vol 22, no 191, 8 March 2022

#### Background

The COVID-19 social restrictions have increased the risk for depression compared to the previous period in Italian women with Low-Risk Pregnancy (LRP). Less is known about the impact of COVID-19 restrictions on High-Risk Pregnancy (HRP). This study aimed: 1) to explore levels of depression in women who become pregnant before and during COVID-19 pandemic, distinguishing between LRP and HRP; 2) to analyze the impact of COVID-19 restrictions on pregnancy experience in LRP and HRP.

#### Methods

A before-during COVID-19 pandemic cross-sectional study was carried out on 155 pregnant women (Mean age = 34.18), between 23 and 32 weeks of gestation. 77 women were recruited before COVID-19 pandemic (51.9% LRP; 48.1% HRP) and 78 women were recruited during COVID-19 pandemic (51.3% LRP; 48.7% HRP). HRP group was enrolled during hospitalization for high-risk pregnancy. Participants filled out Edinburgh Postnatal Depression Scale. Moreover, only COVID-19 group answered an open-ended question about the impact of restriction on pregnancy experience.

#### Results

HRP women reported higher levels of depressive symptoms than LRP. No difference emerged for COVID (before/during) but an interaction effect between COVID-19 and obstetric condition was found. The qualitative results showed the impact of restrictions on emotions and concerns.

#### Conclusion

Respect to the previous period, LRP women during COVID-19 presented an increased risk for depressive symptoms than HRP. The HRP women during COVID-19 seemed to use hospitalization as a resource to find a social support network with other pregnant women and to be reassured on the clinical ongoing of pregnancy. (Author)

**Full URL:** <https://doi.org/10.1186/s12884-022-04515-3>

---

MIDIRS is part of RCM Information Services Limited which is a company incorporated in England and Wales under company no.11914882 with registered office at 10-18 Union Street, London SE1 1SZ  
RCM Information Services Limited is a subsidiary of The Royal College of Midwives



**The Royal College of Midwives Trust**  
10-18 Union Street  
London  
SE1 1SZ



**Open 24 hours a day, 7 days a week**  
T: 0300 303 0444  
F: +44 20 7312 3536  
E: [info@rcm.org.uk](mailto:info@rcm.org.uk)  
W: [www.rcm.org.uk](http://www.rcm.org.uk)



**Chief Executive:**  
Gill Walton,  
MA, PGDip, BSc Hons, RM  
**President:**  
Rebecca Davies, RM  
**Patron:**  
HRH The Princess Royal

**2022-00975**

**Adverse perinatal outcomes in a large United States birth cohort during the COVID-19 pandemic.** Litman EA, Yin Y, Nelson SJ, et al (2022), American Journal of Obstetrics & Gynecology MFM vol 4, no 3, May 2022, 100577

**BACKGROUND**

The impact of coronavirus disease 2019 (COVID-19) on adverse perinatal outcomes remains unclear.

**OBJECTIVE**

This study aimed to investigate whether COVID-19 is associated with adverse perinatal outcomes in a large national dataset and to examine the rates of adverse outcomes during the pandemic compared with the rates of adverse outcomes during the prepandemic period.

**STUDY DESIGN**

This observational cohort study included 683,905 patients, between the ages of 12 and 50, hospitalized for childbirth and abortion between January 1, 2019, and May 31, 2021. During the prepandemic period, 271,444 women were hospitalized for childbirth. During the pandemic, 308,532 women were hospitalized for childbirth, and 2708 women had COVID-19. The associations between COVID-19 and in-hospital adverse perinatal outcomes were examined using propensity score-adjusted logistic regression.

**RESULTS**

Women with COVID-19 were more likely to experience both early and late preterm birth (adjusted odds ratios, 1.38 [95% confidence interval, 1.1–1.7] and 1.62 [95% confidence interval, 1.3–1.7], respectively), preeclampsia (adjusted odds ratio, 1.2 [95% confidence interval, 1.0–1.4]), disseminated intravascular coagulopathy (adjusted odds ratio, 1.57 [95% confidence interval, 1.1–2.2]), pulmonary edema (adjusted odds ratio, 2.7 [95% confidence interval, 1.1–6.3]), and need for mechanical ventilation (adjusted odds ratio, 8.1 [95% confidence interval, 3.8–17.3]) than women without COVID-19. There was no significant difference in the prevalence of stillbirth among women with COVID-19 (16/2708) and women without COVID-19 (174/39,562) ( $P=.257$ ). There was no difference in adverse outcomes among women who delivered during the pandemic vs prepandemic period. Combined in-hospital mortality was significantly higher for women with COVID-19 (147 [95% confidence interval, 3.0–292.0] vs 2.5 [95% confidence interval, 0.0–7.5] deaths per 100,000 women). Women diagnosed with COVID-19 within 30 days before hospitalization were more likely to experience early preterm birth, placental abruption, and mechanical ventilation than women diagnosed with COVID-19 >30 days before hospitalization for childbirth (4.0% vs 2.4% for early preterm birth [adjusted odds ratio, 1.7; 95% confidence interval, 1.1–2.7]; 2.2% vs 1.2% for placental abruption [adjusted odds ratio, 1.86; 95% confidence interval, 1.0–3.4]; and 0.9% vs 0.1% for mechanical ventilation [adjusted odds ratio, 13.7; 95% confidence interval, 1.8–107.2]).

**CONCLUSION**

Women with COVID-19 had a higher prevalence of adverse perinatal outcomes and increased in-hospital mortality, with the highest risk occurring when the diagnosis was within 30 days of hospitalization, raising the possibility of a high-risk period. (Author)

**Full URL:** <https://doi.org/10.1016/j.ajogmf.2022.100577>

---

**2022-00954**

**Domestic violence: an invisible pandemic.** Dey T, Thakar R (2022), The Obstetrician and Gynaecologist vol 24, no 2, April 2022, pp 90-92

Commentary on the increase in domestic violence during the COVID-19 pandemic and the ways in which obstetricians, midwives and allied health professionals can help with early recognition and referral. (LDO)

**Full URL:** <https://doi.org/10.1111/tog.12798>

---

MIDIRS is part of RCM Information Services Limited which is a company incorporated in England and Wales under company no.11914882 with registered office at 10-18 Union Street, London SE1 1SZ  
RCM Information Services Limited is a subsidiary of The Royal College of Midwives



**The Royal College of Midwives Trust**  
10-18 Union Street  
London  
SE1 1SZ



**Open 24 hours a day, 7 days a week**  
T: 0300 303 0444  
F: +44 20 7312 3536  
E: [info@rcm.org.uk](mailto:info@rcm.org.uk)  
W: [www.rcm.org.uk](http://www.rcm.org.uk)



**Chief Executive:**  
Gill Walton,  
MA, PGDip, BSc Hons, RM  
**President:**  
Rebecca Davies, RM  
**Patron:**  
HRH The Princess Royal

**2022-00868**

**The role of perceived social support on pregnant women's mental health during the COVID-19 pandemic.** Corno G, Villani D, de Montigny F, et al (2022), Journal of Reproductive and Infant Psychology 23 February 2022, online

**Purpose**

The present study aimed at investigating which sources of social support best account for pregnant women's levels of psychological distress and mental well-being during the COVID-19 pandemic.

**Methods**

274 Italian and Canadian expectant mothers completed an online-based survey including measures of perceived social support (from family, significant other and friends), state anxiety, depressive symptoms, and satisfaction with life. Correlation analyses and multivariate analysis of covariance were performed to explore how social support from different sources was related to depressive symptoms, state anxiety and satisfaction with life.

**Results**

Different sources of social support contributed to explaining women's psychological distress and mental well-being. Social support both from family and friends was significantly related to women's state anxiety and depressive symptoms. Social support from friends was specifically related to women's satisfaction with life.

**Conclusion**

Our findings endorse the crucial role of perceived social support as a protective factor for pregnant women's mental health. In the context of the COVID-19 pandemic, our results suggest that support from family seems important in preventing psychological distress, whereas support from friends is also associated with mental well-being. These results may help designing future interventions aimed at improving women's perinatal mental health in life-threatening conditions. (Author)

---

MIDIRS is part of RCM Information Services Limited which is a company incorporated in England and Wales under company no.11914882 with registered office at 10-18 Union Street, London SE1 1SZ  
RCM Information Services Limited is a subsidiary of The Royal College of Midwives



**The Royal College of Midwives Trust**  
10-18 Union Street  
London  
SE1 1SZ



**Open 24 hours a day, 7 days a week**  
T: 0300 303 0444  
F: +44 20 7312 3536  
E: [info@rcm.org.uk](mailto:info@rcm.org.uk)  
W: [www.rcm.org.uk](http://www.rcm.org.uk)



**Chief Executive:**  
Gill Walton,  
MA, PGDip, BSc Hons, RM  
**President:**  
Rebecca Davies, RM  
**Patron:**  
HRH The Princess Royal

**2022-00849**

**Rural Residents' Perinatal Experiences During the Initial Months of the COVID-19 Pandemic: A Qualitative Study in British Columbia.** Sullivan E, Cameron A, Kornelsen J (2022), Journal of Midwifery & Women's Health 4 March 2022, online

#### Introduction

Many studies have explored the impact of the coronavirus disease 2019 (COVID-19) pandemic on perinatal health, but few have examined the effects of the pandemic on birthing families through a rural lens. Given that the COVID-19 pandemic has reinforced long-standing disparities between urban and rural communities, it is important that the significance of place on the health and wellness of rural populations is made visible.

#### Methods

In-depth interviews and focus groups with 16 participants from rural communities in British Columbia, Canada, were performed. Participants included those who had been pregnant or given birth after March 11, 2020. Data from the interviews and focus groups were analyzed using the principles of thematic analysis to understand the perinatal experiences of rural families during the initial months of the COVID-19 pandemic.

#### Results

Analysis of the data revealed 4 major themes: perceived risk of infection, navigating uncertainty, experience of care received, and resilience and silver linings. In general, participants conceptualized rural communities as safer bubbles. Exceptions included specific vectors of risk such as tourism travel and border communities. Challenges experienced by rural families including anxiety around changing health guidelines, reduced social support, and potential loss of their partners' support at births. Additional concerns specific to rural experiences added to this burden, including fear of traveling to referral centers for care and increased difficulties accessing resources.

#### Discussion

Participants reported positive, compassionate care experiences that helped to mitigate some of the added stressors of the pandemic. These findings highlight the importance of perinatal care provision that integrates physiologic and mental health supports. This study provides a foundation for a comprehensive inquiry into the experiences of rural perinatal services during COVID-19. (Author)

**Full URL:** <https://doi.org/10.1111/jmwh.13338>

---

**2022-00724**

**Coronavirus: Screening [written answer].** House of Commons (2022), Hansard Written question 131322, 28 February 2022

Maggie Throup responds to a written question from Daisy Cooper to the Secretary of State for Health and Social Care, regarding what assessment his Department has made of the impact of ending free-of-charge covid-19 lateral flow tests on pregnant women. (JSM)

**Full URL:** <https://questions-statements.parliament.uk/written-questions/detail/2022-02-28/131322>

---

MIDIRS is part of RCM Information Services Limited which is a company incorporated in England and Wales under company no.11914882 with registered office at 10-18 Union Street, London SE1 1SZ  
RCM Information Services Limited is a subsidiary of The Royal College of Midwives



**The Royal College of Midwives Trust**  
10-18 Union Street  
London  
SE1 1SZ



**Open 24 hours a day, 7 days a week**  
T: 0300 303 0444  
F: +44 20 7312 3536  
E: [info@rcm.org.uk](mailto:info@rcm.org.uk)  
W: [www.rcm.org.uk](http://www.rcm.org.uk)



**Chief Executive:**  
Gill Walton,  
MA, PGDip, BSc Hons, RM  
**President:**  
Rebecca Davies, RM  
**Patron:**  
HRH The Princess Royal



**2022-00702**

**Feeding during therapeutic hypothermia is safe and may improve outcomes in newborns with perinatal asphyxia.** Wang X, Chen X, Zhang K, et al (2022), Journal of Maternal-Fetal and Neonatal Medicine 13 February 2022, online

**Background:** The Coronavirus disease 2019 (COVID-19) pandemic has become worldwide, posing particularly severe challenges. Pregnancy brings changes that might make individuals more vulnerable to this viral infection. To date, the impact of COVID-19 infection on pregnancy outcomes remains controversial.

**Method:** We performed a meta-analysis to address the impact of COVID-19 infection on pregnancy outcomes. We searched the PubMed and China National Knowledge infrastructure (CNKI) databases for related articles. The odds ratio (OR) corresponding to the 95% confidence interval (95% CI) was used to define the impact of INFECTION and severity of COVID-19 on pregnancy outcomes. The statistical heterogeneity among studies was batched with the Q-test and I2 statistics.

**Results:** We collected 38 studies including 127,805 pregnancy women. Our meta-analysis revealed that pregnant women with COVID-19 have been linked to an increased risk of premature birth (OR = 1.66, 95% CI = 1.41–1.96), stillbirth (OR = 1.98, 95% CI = 1.22–3.21), pre-eclampsia (OR = 1.46, 95% CI = 1.18–1.80), and PROM (OR = 1.39, 95% CI = 1.07–1.81).

**Conclusions:** Our meta-analysis showed that infection with COVID-19 increases the risk of preterm birth, stillbirth, pre-eclampsia, and PROM. Screening and early care for pregnant women to intervene with COVID-19 is important, given the increased risk of adverse pregnancy outcomes. (Author)

---

MIDIRS is part of RCM Information Services Limited which is a company incorporated in England and Wales under company no.11914882 with registered office at 10-18 Union Street, London SE1 1SZ  
RCM Information Services Limited is a subsidiary of The Royal College of Midwives



**The Royal College of Midwives Trust**  
10-18 Union Street  
London  
SE1 1SZ



**Open 24 hours a day, 7 days a week**  
T: 0300 303 0444  
F: +44 20 7312 3536  
E: [info@rcm.org.uk](mailto:info@rcm.org.uk)  
W: [www.rcm.org.uk](http://www.rcm.org.uk)



**Chief Executive:**  
Gill Walton,  
MA, PGDip, BSc Hons, RM  
**President:**  
Rebecca Davies, RM  
**Patron:**  
HRH The Princess Royal

**2022-00699**

**Remdesivir use in pregnancy during the SARS-CoV-2 pandemic.** Gutierrez R, Mendez-Figueroa H, Biebighauser JG, et al (2022), Journal of Maternal-Fetal and Neonatal Medicine 15 February 2022, online

#### Objective

To ascertain the composite maternal and neonatal outcomes in pregnant individuals with moderate, severe, or critical coronavirus disease 2019 (COVID-19) treated with remdesivir.

#### Materials and methods

This is a secondary analysis of the COVID in Pregnancy Registry in Houston, Texas. Women were included if they met the criteria of moderate, severe or critical COVID-19 illness. Composite adverse maternal outcome was defined as any of the following outcomes: placental abruption, pregnancy-related hypertension, chorioamnionitis, stroke, delivery with estimated blood loss >1000 mL, diagnosis of pulmonary embolism or deep venous thromboembolism, or maternal death. Composite adverse neonatal outcome was defined as any of the following: Apgar score  $\leq 3$  at 5 min, arterial cord pH <7.0, positive SAR-CoV-2 test, intraventricular hemorrhage, periventricular leukomalacia, stillbirth, or neonatal death. Comparative analyses between participants receiving remdesivir versus those not exposed were performed.

#### Results

A total of 994 patients were diagnosed with COVID-19 infection. Of these, 95 (9.6%) met criteria for moderate, severe, or critical disease. Forty-one percent of these patients (n = 39) received remdesivir. Baseline demographic characteristics were not different between groups. No patients reported an allergic reaction with the administration of remdesivir; however, 16.7% of the patients had the medication discontinued due to transaminitis. Patients receiving the drug were more likely to have a longer illness duration on admission, more likely to require oxygen support on arrival and have a longer hospital stay.

#### Conclusions

Remdesivir appears to be safe, well tolerated within our cohort with no cases of recorded adverse reaction. (Author)

---

MIDIRS is part of RCM Information Services Limited which is a company incorporated in England and Wales under company no.11914882 with registered office at 10-18 Union Street, London SE1 1SZ  
RCM Information Services Limited is a subsidiary of The Royal College of Midwives



**The Royal College of Midwives Trust**  
10-18 Union Street  
London  
SE1 1SZ



**Open 24 hours a day, 7 days a week**  
T: 0300 303 0444  
F: +44 20 7312 3536  
E: [info@rcm.org.uk](mailto:info@rcm.org.uk)  
W: [www.rcm.org.uk](http://www.rcm.org.uk)



**Chief Executive:**  
Gill Walton,  
MA, PGDip, BSc Hons, RM  
**President:**  
Rebecca Davies, RM  
**Patron:**  
HRH The Princess Royal

**2022-00698**

**Symptomatic versus asymptomatic COVID-19: does it impact placental vasculopathy?**. Ramey-Collier K, Craig AM, Hall A, et al (2022), Journal of Maternal-Fetal and Neonatal Medicine 16 February 2022, online

This study sought to assess the impact of COVID-19 on placental vasculature in the context of maternal symptomatology – comparing asymptomatic to symptomatic pregnant patients – and disease severity – comparing pregnant patients with mild, moderate, severe, and critical COVID-19 infection. PCR-confirmed COVID-19 positive pregnant patients in a single health system who delivered between 3/2020-5/2021 included. All patients had positive COVID test and delivered during the study period. Primary outcome was incidence of any vascular malperfusion on placental pathology. Secondary outcomes were FVM and MVM on placental pathology. Placental pathology compared between symptomatic (sCOVID) and asymptomatic (aCOVID) patients. Secondary analysis of symptomatic patients, comparing placental pathology between mild disease(mCOVID) and worse disease(moderate, severe, or critical-defined by 2020 NIH guidelines) (dCOVID), also performed. Of 112 patients, 53 (47%) had symptoms. Twenty-seven (24.1%) patients had evidence of vascular malperfusion; 26 (23.2%) had MVM. When comparing aCOVID and sCOVID patients, no difference in rate of vascular malperfusion identified, nor any differences in rates of FVM or MVM. Among sCOVID patients (n = 53), 39 (74%) had mCOVID and 14 (26%) had dCOVID (moderate n = 4, severe n = 9, critical n = 1). Patients with dCOVID had earlier median delivery GA (37.4wks vs 39.2wks, p = .03). No difference in latency from diagnosis to delivery seen between mCOVID and dCOVID groups (4.4 vs 3.0wks, p = .96). Twelve (30.8%) patients had vascular malperfusion on pathology, all had mCOVID (p = .02). Eleven (28.2%) mCOVID patients had MVM; no dCOVID patients had evidence of vascular malperfusion (p = .03). No difference in FVM was found between cohorts. Symptomatic COVID-19 infection did not impact placental vasculature differently than asymptomatic infection, even when stratifying by trimester of infection. Among pregnant patients with symptomatic COVID-19, mild disease was associated with placental vascular changes on the maternal side while severe disease was not. Further studies are needed to understand the implications of these findings. (Author)

---

**2022-00692**

**Maternity leave: Coronavirus [written answer]**. Northern Ireland Assembly (2022), Hansard Written question AQW 29593/17-22, 10 February 2022

The Minister for the Economy responds to a written question asked by Mr Harry Harvey, regarding whether he will outline the COVID-19 risk assessment policy in relation to maternity leave guidelines. (LDO)

**Full URL:** <http://aims.niassembly.gov.uk/questions/printquestionssummary.aspx?docid=367579>

---

**2022-00689**

**Antenatal classes [written answer]**. Scottish Parliament (2022), Official Report Written question S6W-06381, 9 February 2022

Humza Yousaf responds to a written question asked by Monica Lennon to the Scottish Government regarding the date by which NHS Scotland will resume face-to-face antenatal classes, in light of the reported resumption of private classes. (LDO)

**Full URL:** <https://archive2021.parliament.scot/parliamentarybusiness/28877.aspx?SearchType=Advance&ReferenceNumbers=S6W-06381>

---

**2022-00686**

**Comparison of Sexually Transmitted Infections and Adverse Perinatal Outcomes in Underserved Pregnant Patients Before vs During the COVID-19 Pandemic in Texas.** Stafford IA, Coselli JO, Wilson DF, et al (2022), JAMA Network Open vol 5, no 2, February 2022, e220568

This cohort study assesses the rates of sexually transmitted infections (STI) and associated adverse perinatal outcomes among underserved pregnant patients during the COVID-19 pandemic in a public health system in Southeastern Texas. (Author)

**Full URL:** <https://doi.org/10.1001/jamanetworkopen.2022.0568>

---

MIDIRS is part of RCM Information Services Limited which is a company incorporated in England and Wales under company no.11914882 with registered office at 10-18 Union Street, London SE1 1SZ  
RCM Information Services Limited is a subsidiary of The Royal College of Midwives



**The Royal College of Midwives Trust**  
10-18 Union Street  
London  
SE1 1SZ



**Open 24 hours a day, 7 days a week**  
T: 0300 303 0444  
F: +44 20 7312 3536  
E: [info@rcm.org.uk](mailto:info@rcm.org.uk)  
W: [www.rcm.org.uk](http://www.rcm.org.uk)



**Chief Executive:**  
Gill Walton,  
MA, PGDip, BSc Hons, RM  
**President:**  
Rebecca Davies, RM  
**Patron:**  
HRH The Princess Royal

**2022-00657**

**Severity of maternal infection and perinatal outcomes during periods of SARS-CoV-2 wildtype, alpha, and delta variant dominance in the UK: prospective cohort study.** Vousden N, Ramakrishnan R, Bunch K, et al (2022), *BMJ Medicine* 28 February 2022, online

**Objective** To compare the severity of maternal infection and perinatal outcomes during periods in which wildtype, alpha variant, and delta variant of SARS-CoV-2 were dominant in the UK.

**Design** Prospective cohort study.

**Setting** 194 obstetric units across the UK, during the following periods: between 1 March and 30 November 2020 (wildtype dominance), between 1 December 2020 and 15 May 2021 (alpha variant dominance), and between 16 May and 31 October 2021 (delta variant dominance).

**Participants** 4436 pregnant women admitted to hospital with covid-19 related symptoms.

**Main outcome measures** Moderate to severe maternal SARS-CoV-2 infection (indicated by any of the following: oxygen saturation <95% on admission, need for oxygen treatment, evidence of pneumonia on imaging, admission to intensive care, or maternal death), and pregnancy and perinatal outcomes (including mode and gestation of birth, stillbirth, live birth, admission to neonatal intensive care, and neonatal death).

**Results** 1387, 1613, and 1436 pregnant women were admitted to hospital with covid-19 related symptoms during the wildtype, alpha, and delta dominance periods, respectively; of these women, 340, 585, and 614 had moderate to severe infection, respectively. The proportion of pregnant women admitted with moderate to severe infection increased during the subsequent alpha and delta dominance periods, compared with the wildtype dominance period (wildtype 24.5% v alpha 36.2% (adjusted odds ratio 1.98, 95% confidence interval 1.66% to 2.37%); wildtype 24.5% v delta 42.8% (2.66, 2.21 to 3.20)). Compared with the wildtype dominance period, women admitted during the alpha dominance period were significantly more likely to have pneumonia, require respiratory support, and be admitted to intensive care; these three risks were even greater during the delta dominance period (wildtype v delta: pneumonia, adjusted odds ratio 2.52, 95% confidence interval 2.06 to 3.09; respiratory support, 1.90, 1.52 to 2.37; and intensive care, 2.71, 2.06 to 3.56). Of 1761 women whose vaccination status was known, 38 (2.2%) had one dose and 16 (1%) had two doses before their diagnosis (of whom 14 (88%) had mild infection). The proportion of women receiving drug treatment for SARS-CoV-2 management was low, but did increase between the wildtype dominance period and the alpha and delta dominance periods (10.4% wildtype v 14.9% alpha (2.74, 2.08 to 3.60); 10.4% wildtype v 13.6% delta (2.54, 1.90 to 3.38)).

**Conclusions** While limited by the absence of variant sequencing data, these findings suggest that during the periods when the alpha and delta variants of SARS-CoV-2 were dominant, covid-19 was associated with more severe maternal infection and worse pregnancy outcomes than during the wildtype dominance period. Most women admitted with SARS-CoV-2 related symptoms were unvaccinated. Urgent action to prioritise vaccine uptake in pregnancy is essential.

**Study registration** ISRCTN40092247. (Author)

**Full URL:** <https://bmjmedicine.bmj.com/content/1/1/e000053>

---

MIDIRS is part of RCM Information Services Limited which is a company incorporated in England and Wales under company no.11914882 with registered office at 10-18 Union Street, London SE1 1SZ  
RCM Information Services Limited is a subsidiary of The Royal College of Midwives



**The Royal College of Midwives Trust**  
10-18 Union Street  
London  
SE1 1SZ



**Open 24 hours a day, 7 days a week**  
T: 0300 303 0444  
F: +44 20 7312 3536  
E: [info@rcm.org.uk](mailto:info@rcm.org.uk)  
W: [www.rcm.org.uk](http://www.rcm.org.uk)



**Chief Executive:**  
Gill Walton,  
MA, PGDip, BSc Hons, RM  
**President:**  
Rebecca Davies, RM  
**Patron:**  
HRH The Princess Royal

**2022-00654**

**Covid-19: Severe infection in pregnancy significantly increases risks, study shows.** Iacobucci G (2022), BMJ vol 376, 24 February 2022, o480

Severe covid-19 infection in pregnant women significantly increases the risk of harmful outcomes for mothers and babies, a study (1) has found.

1. Vousden N et al. Acta Obstetrica et Gynecologica Scandinavica, 25 February 2022, online.

<https://doi.org/10.1111/aogs.14329>.

(Author, edited)

**Full URL:** <https://doi.org/10.1136/bmj.o480>

---

**2022-00641**

**COVID-19 and Pregnancy.** Walter K (2022), JAMA (Journal of the American Medical Association) vol 327, no 8, 22 February 2022, p 790

This JAMA Patient Page describes characteristics of COVID-19 among pregnant people and vaccination recommendations for people who are pregnant or breastfeeding. (Author)

---

MIDIRS is part of RCM Information Services Limited which is a company incorporated in England and Wales under company no.11914882 with registered office at 10-18 Union Street, London SE1 1SZ  
RCM Information Services Limited is a subsidiary of The Royal College of Midwives



**The Royal College of Midwives Trust**  
10-18 Union Street  
London  
SE1 1SZ



**Open 24 hours a day, 7 days a week**  
T: 0300 303 0444  
F: +44 20 7312 3536  
E: [info@rcm.org.uk](mailto:info@rcm.org.uk)  
W: [www.rcm.org.uk](http://www.rcm.org.uk)



**Chief Executive:**  
Gill Walton,  
MA, PGDip, BSc Hons, RM  
**President:**  
Rebecca Davies, RM  
**Patron:**  
HRH The Princess Royal

## 2022-00640

**Association of SARS-CoV-2 Infection With Serious Maternal Morbidity and Mortality From Obstetric Complications.** Metz TD, Clifton RG, Hughes BL, et al (2022), JAMA (Journal of the American Medical Association) vol 327, no 8, 22 February 2022, pp 748-759

**Importance** It remains unknown whether SARS-CoV-2 infection specifically increases the risk of serious obstetric morbidity.

**Objective** To evaluate the association of SARS-CoV-2 infection with serious maternal morbidity or mortality from common obstetric complications.

**Design, Setting, and Participants** Retrospective cohort study of 14 104 pregnant and postpartum patients delivered between March 1, 2020, and December 31, 2020 (with final follow-up to February 11, 2021), at 17 US hospitals participating in the Eunice Kennedy Shriver National Institute of Child Health and Human Development's Gestational Research Assessments of COVID-19 (GRAVID) Study. All patients with SARS-CoV-2 were included and compared with those without a positive SARS-CoV-2 test result who delivered on randomly selected dates over the same period.

**Exposures** SARS-CoV-2 infection was based on a positive nucleic acid or antigen test result. Secondary analyses further stratified those with SARS-CoV-2 infection by disease severity.

**Main Outcomes and Measures** The primary outcome was a composite of maternal death or serious morbidity related to hypertensive disorders of pregnancy, postpartum hemorrhage, or infection other than SARS-CoV-2. The main secondary outcome was cesarean birth.

**Results** Of the 14 104 included patients (mean age, 29.7 years), 2352 patients had SARS-CoV-2 infection and 11 752 did not have a positive SARS-CoV-2 test result. Compared with those without a positive SARS-CoV-2 test result, SARS-CoV-2 infection was significantly associated with the primary outcome (13.4% vs 9.2%; difference, 4.2% [95% CI, 2.8%-5.6%]; adjusted relative risk [aRR], 1.41 [95% CI, 1.23-1.61]). All 5 maternal deaths were in the SARS-CoV-2 group. SARS-CoV-2 infection was not significantly associated with cesarean birth (34.7% vs 32.4%; aRR, 1.05 [95% CI, 0.99-1.11]). Compared with those without a positive SARS-CoV-2 test result, moderate or higher COVID-19 severity (n = 586) was significantly associated with the primary outcome (26.1% vs 9.2%; difference, 16.9% [95% CI, 13.3%-20.4%]; aRR, 2.06 [95% CI, 1.73-2.46]) and the major secondary outcome of cesarean birth (45.4% vs 32.4%; difference, 12.8% [95% CI, 8.7%-16.8%]; aRR, 1.17 [95% CI, 1.07-1.28]), but mild or asymptomatic infection (n = 1766) was not significantly associated with the primary outcome (9.2% vs 9.2%; difference, 0% [95% CI, -1.4% to 1.4%]; aRR, 1.11 [95% CI, 0.94-1.32]) or cesarean birth (31.2% vs 32.4%; difference, -1.4% [95% CI, -3.6% to 0.8%]; aRR, 1.00 [95% CI, 0.93-1.07]).

**Conclusions and Relevance** Among pregnant and postpartum individuals at 17 US hospitals, SARS-CoV-2 infection was associated with an increased risk for a composite outcome of maternal mortality or serious morbidity from obstetric complications. (Author)

---

## 2022-00598

**Maternity Services: Coronavirus [written answer].** House of Commons (2022), Hansard Written question 99094, 6 January 2022

Maria Caulfield responds to a written question asked by Olivia Blake to the Secretary of State for Health and Social Care, with reference to NHS guidance, Supporting pregnant women using maternity services during the coronavirus pandemic: Actions for NHS providers, published on 14 December 2020, what steps he is taking to ensure that NHS Trusts adhere to that guidance; and what steps his department is taking to support NHS trusts to follow that guidance. (LDO)

**Full URL:** <https://questions-statements.parliament.uk/written-questions/detail/2022-01-06/99094>

---

MIDIRS is part of RCM Information Services Limited which is a company incorporated in England and Wales under company no.11914882 with registered office at 10-18 Union Street, London SE1 1SZ  
RCM Information Services Limited is a subsidiary of The Royal College of Midwives



**The Royal College of Midwives Trust**  
10-18 Union Street  
London  
SE1 1SZ



**Open 24 hours a day, 7 days a week**  
T: 0300 303 0444  
F: +44 20 7312 3536  
E: [info@rcm.org.uk](mailto:info@rcm.org.uk)  
W: [www.rcm.org.uk](http://www.rcm.org.uk)



**Chief Executive:**  
Gill Walton,  
MA, PGDip, BSc Hons, RM  
**President:**  
Rebecca Davies, RM  
**Patron:**  
HRH The Princess Royal

**2022-00583**

**Use of single-dose tocilizumab for treatment of severe COVID-19 in pregnancy: implications for the timing of live infant vaccines.** Burkhardt I, Whittaker E (2022), Archives of Disease in Childhood vol 107, no 5, May 2022, p 517

Correspondence piece discussing the safety of live vaccines administered to infants after prenatal exposure to tocilizumab for the treatment of COVID-19 in the mother. The authors recommend delaying live vaccines until six months of age for infants who have been exposed. (LDO)

Full URL: <http://dx.doi.org/10.1136/archdischild-2021-323628>

---

**2022-00525**

**Impact of the COVID-19 pandemic on the incidence of prematurity: critical role of gestational age and environment.**

Weinberger B, Divers J, Campbell D, et al (2022), American Journal of Obstetrics & Gynecology (AJOG) vol 226, no 6, June 2022, pp 864-866

Research letter comparing rates of extreme prematurity with and without pre-eclampsia in New York during the COVID-19 pandemic. Results demonstrate an overall decrease in the number of extreme preterm births despite an increased rate of pre-eclampsia. (LDO)

Full URL: <https://doi.org/10.1016/j.ajog.2022.02.028>

---

**2022-00443**

**Industrial Health and Safety: Pregnancy [written answer].** House of Commons (2022), Hansard Written question 125552, 18 February 2022

Chloe Smith responds to a written question asked by Zarah Sultana to the Secretary of State for Health and Social Care, regarding what assessment her department has made of the implications for its policies of the recommendations of the December 2021 Maternity Action report, Unsafe and Unsupported, on workplace health and safety for pregnant women during the COVID-19 outbreak. (LDO)

Full URL: <https://questions-statements.parliament.uk/written-questions/detail/2022-02-18/125552>

---

**2022-00436**

**Severe Covid-19 infection linked to increase in risk during pregnancy.** Baines E (2022), Nursing Times 24 February 2022

Pregnant women who are admitted to hospital with severe Covid-19 are at increased risk of serious birth complications, a UK study has shown. (Author)

---

MIDIRS is part of RCM Information Services Limited which is a company incorporated in England and Wales under company no.11914882 with registered office at 10-18 Union Street, London SE1 1SZ  
RCM Information Services Limited is a subsidiary of The Royal College of Midwives



**The Royal College of Midwives Trust**  
10-18 Union Street  
London  
SE1 1SZ



**Open 24 hours a day, 7 days a week**  
T: 0300 303 0444  
F: +44 20 7312 3536  
E: [info@rcm.org.uk](mailto:info@rcm.org.uk)  
W: [www.rcm.org.uk](http://www.rcm.org.uk)



**Chief Executive:**  
Gill Walton,  
MA, PGDip, BSc Hons, RM  
**President:**  
Rebecca Davies, RM  
**Patron:**  
HRH The Princess Royal

## 2022-00362

**Obstetric outcomes in pregnant COVID-19 women: the imbalance of von Willebrand factor and ADAMTS13 axis.** Grandone E, Vimercati A, Sorrentino F, et al (2022), BMC Pregnancy and Childbirth vol 22, no 142, 21 February 2022

### Background

Thrombotic microangiopathy has been invoked as one of the most important mechanisms of damage in COVID-19 patients. Protease ADAMTS13 is a marker of microangiopathy responsible for controlling von Willebrand multimers size. Von Willebrand factor/ADAMTS13 ratio has been found impaired in COVID-19 patients outside pregnancy.

### Methods

We prospectively investigated 90 pregnant women admitted to two tertiary academic hospitals in Italy with a laboratory-confirmed diagnosis of SARS-CoV-2 infection. Demographic, clinical information and routine laboratory data were collected at the hospital admission and until discharge. We investigated whether vonWillebrand /ADAMTS13 axis imbalance is a predictor of adverse outcomes. Logistic regression analysis, which controlled for potential confounders, was performed to evaluate the association between laboratory parameters and clinical outcomes.

### Results

Most women (55.6%) were parae, with median gestational age at admission of 39 weeks. At hospital admission, 63.3% were asymptomatic for COVID-19 and 24.4% showed more than one sign or symptom of infection. Nulliparae with group O showed Willebrand / ADA MTS-13 ratios significantly lower than non-O, whereas in multiparae this difference was not observed. Logistic regression showed that ratio von Willebrand to ADAMTS13 was significantly and independently associated with preterm delivery (OR 1.9, 95%CI 1.1–3.5).

### Conclusion

This study shows an imbalance of vonWillebrand /ADAMTS13 axis in pregnant women with COVID-19, leading to a significantly higher and independent risk of preterm delivery. Monitoring these biomarkers might support decision making process to manage and follow-up pregnancies in this setting. (Author)

**Full URL:** <https://doi.org/10.1186/s12884-022-04405-8>

---

## 2022-00293

**Potential role of neurofilament in COVID-19 and preeclampsia.** Samara A, Herlenius E, O' Brien P, et al (2022), Cell Reports. Medicine vol 3, no 1, January 2022, e100490

Neurofilament light (NFL) is a promising circulating biomarker in preeclampsia and COVID-19, even without evident neurological complications. Several pathways might contribute to the elevated serum NFL levels seen in both pathologies. Future studies will determine whether NFL is a long COVID marker and delineate NFL's role in COVID-19-associated preeclampsia. (Author)

**Full URL:** <https://doi.org/10.1016/j.xcrm.2021.100490>

---

MIDIRS is part of RCM Information Services Limited which is a company incorporated in England and Wales under company no.11914882 with registered office at 10-18 Union Street, London SE1 1SZ  
RCM Information Services Limited is a subsidiary of The Royal College of Midwives



**The Royal College of Midwives Trust**  
10-18 Union Street  
London  
SE1 1SZ



**Open 24 hours a day, 7 days a week**  
T: 0300 303 0444  
F: +44 20 7312 3536  
E: [info@rcm.org.uk](mailto:info@rcm.org.uk)  
W: [www.rcm.org.uk](http://www.rcm.org.uk)



**Chief Executive:**  
Gill Walton,  
MA, PGDip, BSc Hons, RM  
**President:**  
Rebecca Davies, RM  
**Patron:**  
HRH The Princess Royal



## 2022-00157

**Pregnancy and risk of COVID-19: a Norwegian registry-linkage study.** Magnus MC, Oakley L, Gjessing HK, et al (2022), BJOG: An International Journal of Obstetrics and Gynaecology vol 129, no 1, January 2022, pp 101-109

### Objective

To compare the risk of acute respiratory syndrome coronavirus 2 (SARS-CoV-2) infection and contact with specialist healthcare services for coronavirus disease 2019 (COVID-19) between pregnant and non-pregnant women.

### Population or sample

All women ages 15–45 living in Norway on 1 March 2020 (n = 1 033 699).

### Methods

We linked information from the national birth, patient, communicable diseases and education databases using unique national identifiers.

### Main outcome measure

We estimated hazard ratios (HR) among pregnant compared to non-pregnant women of having a positive test for SARS-CoV-2, a diagnosis of COVID-19 in specialist healthcare, or hospitalisation with COVID-19 using Cox regression. Multivariable analyses adjusted for age, marital status, education, income, country of birth and underlying medical conditions.

### Results

Pregnant women were not more likely to be tested for or to have a positive SARS-CoV-2 test (adjusted HR 0.99; 95% CI 0.92–1.07). Pregnant women had higher risk of hospitalisation with COVID-19 (HR 4.70, 95% CI 3.51–6.30) and any type of specialist care for COVID-19 (HR 3.46, 95% CI 2.89–4.14). Pregnant women born outside Scandinavia were less likely to be tested, and at higher risk of a positive test (HR 2.37, 95% CI 2.51–8.87). Compared with pregnant Scandinavian-born women, pregnant women with minority background had a higher risk of hospitalisation with COVID-19 (HR 4.72, 95% CI 2.51–8.87).

### Conclusion

Pregnant women were not more likely to be infected with SARS-CoV-2. Still, pregnant women with COVID-19, especially those born outside of Scandinavia, were more likely to be hospitalised. (Author)

**Full URL:** <https://doi.org/10.1111/1471-0528.16969>

---

## 2022-00142

**Maternal Outcomes After Severe Acute Respiratory Syndrome Coronavirus 2 (SARS-CoV-2) Infection in Vaccinated Compared With Unvaccinated Pregnant Patients.** Morgan JA, Biggio JR, Martin JK, et al (2022), Obstetrics & Gynecology vol 139, no 1, January 2022, pp 107-109

Severe acute respiratory syndrome coronavirus 2 (SARS-CoV-2) vaccination is associated with lower odds of severe or critical coronavirus disease 2019 (COVID-19) or COVID-19 of any severity in pregnant patients during the Delta-predominant fourth SARS-CoV-2 surge. (Author)

**Full URL:** [https://journals.lww.com/greenjournal/fulltext/9900/maternal\\_outcomes\\_after\\_severe\\_acute\\_respiratory.320.aspx](https://journals.lww.com/greenjournal/fulltext/9900/maternal_outcomes_after_severe_acute_respiratory.320.aspx)

---

MIDIRS is part of RCM Information Services Limited which is a company incorporated in England and Wales under company no.11914882 with registered office at 10-18 Union Street, London SE1 1SZ  
RCM Information Services Limited is a subsidiary of The Royal College of Midwives



**The Royal College of Midwives Trust**  
10-18 Union Street  
London  
SE1 1SZ



**Open 24 hours a day, 7 days a week**  
T: 0300 303 0444  
F: +44 20 7312 3536  
E: [info@rcm.org.uk](mailto:info@rcm.org.uk)  
W: [www.rcm.org.uk](http://www.rcm.org.uk)



**Chief Executive:**  
Gill Walton,  
MA, PGDip, BSc Hons, RM  
**President:**  
Rebecca Davies, RM  
**Patron:**  
HRH The Princess Royal

**2022-00121**

**SARS-CoV-2 infection in pregnancy during the first wave of COVID-19 in the Netherlands: a prospective nationwide population-based cohort study (NethOSS).** Overtoom EM, Rosman AN, Zwart JJ, et al (2022), BJOG: An International Journal of Obstetrics and Gynaecology vol 129, no 1, January 2022, pp 91-100

**Objective**

To describe characteristics, risk factors and maternal, obstetric and neonatal outcomes of pregnant women infected with severe acute respiratory syndrome coronavirus 2 (SARS-CoV-2).

**Design**

Multi-centre prospective population-based cohort study.

**Setting**

Nationwide study in the Netherlands.

**Population**

Pregnant women with confirmed SARS-CoV-2 infection admitted to hospital or in home-isolation: 1 March 2020 to 31 August 2020.

**Methods**

Pregnant women with positive polymerase chain reaction or antibody tests were registered using the Netherlands Obstetrics Surveillance System (NethOSS). (Selective) testing occurred according to national guidelines. Data from the national birth registry (pregnant pre-coronavirus disease 2019 [COVID-19] cohort) and an age-matched cohort of COVID-19-positive women (National Institute for Public Health and the Environment; fertile age COVID-19 cohort) were used as reference.

**Main outcome measures**

Incidence of SARS-CoV-2 infection in pregnant women. Maternal, obstetric and neonatal outcomes including hospital and intensive care admission.

**Results**

Of 376 registered pregnant women with confirmed SARS-CoV-2 infection, 20% (74/376) were admitted to hospital, of whom 84% (62/74) were due to SARS-CoV-2; 10% (6/62) were admitted to intensive care and 15% (9/62) to obstetric high-care units. Risk factors for admission were non-European country of origin (odds ratio [OR] 1.73, 95% CI 1.01–2.96) and being overweight/obese (OR 1.86, 95% CI 1.51–3.20). No maternal or perinatal deaths occurred. Caesarean section after labour-onset was increased (OR 1.58, 95% CI 1.09–2.28). Hospital and intensive care admission were higher compared with the fertile age COVID-19 cohort (OR 6.75, 95% CI 5.18–8.81 and OR 2.52, 95% CI 1.11–5.77, respectively).

**Conclusions**

Non-European country of origin and being overweight/obese are risk factors for severe course of SARS-CoV-2 infection in pregnancy, risk of caesarean section and hospital and intensive care unit admission are increased. (Author)

**Full URL:** <https://doi.org/10.1111/1471-0528.16903>

---

MIDIRS is part of RCM Information Services Limited which is a company incorporated in England and Wales under company no.11914882 with registered office at 10-18 Union Street, London SE1 1SZ  
RCM Information Services Limited is a subsidiary of The Royal College of Midwives



**The Royal College of Midwives Trust**  
10-18 Union Street  
London  
SE1 1SZ



**Open 24 hours a day, 7 days a week**  
T: 0300 303 0444  
F: +44 20 7312 3536  
E: [info@rcm.org.uk](mailto:info@rcm.org.uk)  
W: [www.rcm.org.uk](http://www.rcm.org.uk)



**Chief Executive:**  
Gill Walton,  
MA, PGDip, BSc Hons, RM  
**President:**  
Rebecca Davies, RM  
**Patron:**  
HRH The Princess Royal

## 2022-00070

**[Withdrawn] Coronavirus (COVID-19): advice for pregnant employees [Withdrawn: 1 April 2022].** Department of Health and Social Care, Health and Safety Executive (2020), London: DHSC 23 December 2020

Advice for pregnant employees on risk assessments in the workplace and occupational health during the coronavirus (COVID-19) pandemic. NB: This guidance was withdrawn on 1 April 2022 (Author)

Full URL: <https://www.gov.uk/government/publications/coronavirus-covid-19-advice-for-pregnant-employees>

---

## 2022-00037

**Performance of diagnostic coding and laboratory testing results to measure COVID-19 during pregnancy and associations with pregnancy outcomes.** Regan AK, Arah OA, Sullivan SG (2022), Paediatric and Perinatal Epidemiology 25 January 2022, online

### Background

Large-scale evaluation of COVID-19 is likely to rely on the quality of ICD coding. However, little is known about the validity of ICD-coded COVID-19 diagnoses.

### Objectives

To evaluate the performance of diagnostic codes in detecting COVID-19 during pregnancy.

### Methods

We used data from a national cohort of 78,283 individuals with a pregnancy ending between 11 March 2020 and 31 January 2021 in the OptumLabs® Data Warehouse (OLDW). OLDW is a longitudinal, real-world data asset with de-identified administrative claims and electronic health record data. We identified all services with an ICD-10-CM diagnostic code of U07.1 and all laboratory claims records for COVID-19 diagnostic testing. We compared ICD-coded diagnoses to testing results to estimate positive and negative predictive values (PPV and NPV). To evaluate impact on risk estimation, we estimated risk of adverse pregnancy outcomes by source of exposure information.

### Results

Of 78,283 pregnancies, 5644 had a laboratory test result for COVID-19. Testing was most common among older individuals, Hispanic individuals, those with higher socioeconomic status and those with a diagnosed medical condition or pregnancy complication; 52% of COVID-19 cases was identified through ICD-coded diagnosis alone, 19% from laboratory test results alone and 29% from both sources. Agreement between ICD-coded diagnosis and laboratory testing records was high 91% (95% confidence interval [CI] 90, 92). However, the PPV of ICD-code diagnosis was low (36%; 95% CI 33, 39). We observed up to a 50% difference in risk estimates of adverse pregnancy outcomes when exposure was based on laboratory testing results or diagnostic coding alone.

### Conclusions

More than one-in-five COVID-19 cases would be missed by using ICD-coded diagnoses alone to identify COVID-19 during pregnancy. Epidemiological studies exclusively relying on diagnostic coding or laboratory testing results are likely to be affected by exposure misclassification. Research and surveillance should draw upon multiple sources of COVID-19 diagnostic information. (Author)

Full URL: <https://doi.org/10.1111/ppe.12863>

---

MIDIRS is part of RCM Information Services Limited which is a company incorporated in England and Wales under company no.11914882 with registered office at 10-18 Union Street, London SE1 1SZ  
RCM Information Services Limited is a subsidiary of The Royal College of Midwives



**The Royal College of Midwives Trust**  
10-18 Union Street  
London  
SE1 1SZ



**Open 24 hours a day, 7 days a week**  
T: 0300 303 0444  
F: +44 20 7312 3536  
E: [info@rcm.org.uk](mailto:info@rcm.org.uk)  
W: [www.rcm.org.uk](http://www.rcm.org.uk)



**Chief Executive:**  
Gill Walton,  
MA, PGDip, BSc Hons, RM  
**President:**  
Rebecca Davies, RM  
**Patron:**  
HRH The Princess Royal

## 2022-00033

**A review of COVID-19 therapeutics in pregnancy and lactation.** Jorgensen SCJ, Tabbara N, Burry L (2022), *Obstetric Medicine* 12 January 2022, online

Pregnant people have an elevated risk of severe COVID-19-related complications compared to their non-pregnant counterparts, underscoring the need for safe and effective therapies. In this review, we summarize published data on COVID-19 therapeutics in pregnancy and lactation to help inform clinical decision-making about their use in this population. Although no serious safety signals have been raised for many agents, data clearly have serious limitations and there are many important knowledge gaps about the safety and efficacy of key therapeutics used for COVID-19. Moving forward, diligent follow-up and documentation of outcomes in pregnant people treated with these agents will be essential to advance our understanding. Greater regulatory push and incentives are needed to ensure studies to obtain pregnancy data are expedited. (Author)

---

## 2022-00012

**Anti-Spike Monoclonal Antibody Therapy in Pregnant Women With Mild-to-Moderate Coronavirus Disease 2019 (COVID-19).**

Thilagar BP, Ghosh AK, Nguyen J, et al (2022), *Obstetrics & Gynecology* vol 139, no 4, April 2022, pp 616-618

Anti-spike monoclonal antibody therapy may be effective for pregnant women with coronavirus disease 2019 (COVID-19). (Author)

Full URL: <https://doi.org/10.1097/AOG.0000000000004700>

---

## 2022-00006

**Neutralizing Monoclonal Antibodies for Coronavirus Disease 2019 (COVID-19) in Pregnancy: A Case Series.** Richley M, Rao RR, Afshar Y, et al (2022), *Obstetrics & Gynecology* vol 139, no 3, March 2022, pp 368-372

### OBJECTIVE:

To describe outcomes associated with monoclonal antibody use in pregnant persons with mild-to-moderate coronavirus disease 2019 (COVID-19).

### METHODS:

We present a retrospective case series of pregnant patients who received anti-severe acute respiratory syndrome coronavirus 2 (SARS-CoV-2) monoclonal antibody infusions at a single center from April 1, 2021, through October 16, 2021. Pregnant patients who had a positive SARS-CoV-2 polymerase chain reaction (PCR) test result and mild-to-moderate COVID-19 symptoms were eligible for monoclonal antibody infusion. Exclusion criteria for administration included need for supplemental oxygen, hospitalization due to COVID-19, and positive SARS-CoV-2 PCR test result more than 7 days before screening. All patients received either bamlanivimab plus etesevimab or casirivimab plus imdevimab based on availability and dosing instructions of the product and emerging resistance patterns in the community.

### RESULTS:

During the study period, monoclonal antibody infusions were administered to 450 individuals at our institution, of whom 15 were pregnant. Of the 15 pregnant persons receiving monoclonal antibody, six (40%) had full-vaccination status at the time of infusion. Two individuals (13%, CI 0–31%) experienced systemic reactions during the infusion, both resulting in temporary changes in the fetal heart rate tracing that recovered with maternal and intrauterine resuscitative efforts. One patient delivered after infusion for worsening maternal and fetal status; the remainder of the patients did not require admission for COVID-19.

### CONCLUSION:

In this case series, pregnant persons who received anti-SARS-CoV-2 monoclonal antibody infusions had generally favorable outcomes. (Author)

Full URL: <https://doi.org/10.1097/AOG.000000000000468>

---

MIDIRS is part of RCM Information Services Limited which is a company incorporated in England and Wales under company no.11914882 with registered office at 10-18 Union Street, London SE1 1SZ  
RCM Information Services Limited is a subsidiary of The Royal College of Midwives



**The Royal College of Midwives Trust**  
10-18 Union Street  
London  
SE1 1SZ



**Open 24 hours a day, 7 days a week**  
T: 0300 303 0444  
F: +44 20 7312 3536  
E: [info@rcm.org.uk](mailto:info@rcm.org.uk)  
W: [www.rcm.org.uk](http://www.rcm.org.uk)



**Chief Executive:**  
Gill Walton,  
MA, PGDip, BSc Hons, RM  
**President:**  
Rebecca Davies, RM  
**Patron:**  
HRH The Princess Royal

**2022-00003**

**Early Administration of Remdesivir and Intensive Care Unit Admission in Hospitalized Pregnant Individuals With Coronavirus Disease 2019 (COVID-19).** Eid J, Abdelwahab M, Colburn N, et al (2022), *Obstetrics & Gynecology* vol 139, no 4, April 2022, pp 619-621

Remdesivir has been shown to shorten the time to recovery in hospitalized patients with coronavirus disease 2019 (COVID-19). Data on its use in pregnancy are limited. In this single-center retrospective cohort study, our objective was to determine whether early remdesivir use in pregnant individuals is associated with decreased risk of admission to the intensive care unit (ICU). Forty-one pregnant patients were included in this study, and outcomes were compared between those who received remdesivir less than 7 days (early group) and 7 or more days (late group) from onset of patient-reported symptoms. Early remdesivir administration was associated with improved clinical outcomes, including lower rates of ICU admission, decreased length of hospitalization, and decreased progression to critical disease in pregnant individuals hospitalized with COVID-19. (Author)

**Full URL:** <https://doi.org/10.1097/AOG.0000000000004734>

---

**2021-14558**

**Adapting obstetric and neonatal services during the COVID-19 pandemic: a scoping review.** Gold S, Clarfield L, Johnstone J, et al (2022), *BMC Pregnancy and Childbirth* vol 22, no 119, 11 February 2022

**Background**

The provision of care to pregnant persons and neonates must continue through pandemics. To maintain quality of care, while minimizing physical contact during the Severe Acute Respiratory Syndrome-related Coronavirus-2 (SARS-CoV2) pandemic, hospitals and international organizations issued recommendations on maternity and neonatal care delivery and restructuring of clinical and academic services. Early in the pandemic, recommendations relied on expert opinion, and offered a one-size-fits-all set of guidelines. Our aim was to examine these recommendations and provide the rationale and context to guide clinicians, administrators, educators, and researchers, on how to adapt maternity and neonatal services during the pandemic, regardless of jurisdiction.

**Method**

Our initial database search used Medical subject headings and free-text search terms related to coronavirus infections, pregnancy and neonatology, and summarized relevant recommendations from international society guidelines. Subsequent targeted searches to December 30, 2020, included relevant publications in general medical and obstetric journals, and updated society recommendations.

**Results**

We identified 846 titles and abstracts, of which 105 English-language publications fulfilled eligibility criteria and were included in our study. A multidisciplinary team representing clinicians from various disciplines, academics, administrators and training program directors critically appraised the literature to collate recommendations by multiple jurisdictions, including a quaternary care Canadian hospital, to provide context and rationale for viable options.

**Interpretation**

There are different schools of thought regarding effective practices in obstetric and neonatal services. Our critical review presents the rationale to effectively modify services, based on the phase of the pandemic, the prevalence of infection in the population, and resource availability. (Author)

**Full URL:** <https://doi.org/10.1186/s12884-022-04409-4>

---

MIDIRS is part of RCM Information Services Limited which is a company incorporated in England and Wales under company no.11914882 with registered office at 10-18 Union Street, London SE1 1SZ  
RCM Information Services Limited is a subsidiary of The Royal College of Midwives



**The Royal College of Midwives Trust**  
10-18 Union Street  
London  
SE1 1SZ



**Open 24 hours a day, 7 days a week**  
T: 0300 303 0444  
F: +44 20 7312 3536  
E: [info@rcm.org.uk](mailto:info@rcm.org.uk)  
W: [www.rcm.org.uk](http://www.rcm.org.uk)



**Chief Executive:**  
Gill Walton,  
MA, PGDip, BSc Hons, RM  
**President:**  
Rebecca Davies, RM  
**Patron:**  
HRH The Princess Royal

## 2021-14555

**Comorbidity, poverty and social vulnerability as risk factors for mortality in pregnant women with confirmed SARS-CoV-2 infection: analysis of 13 062 positive pregnancies including 176 maternal deaths in Mexico.** Torres-Torres J, Martinez-Portilla RJ, Espino-y-Sosa S, et al (2022), *Ultrasound in Obstetrics and Gynecology* vol 59, no 1, January 2022, pp 76-82

### Objective

Mortality in pregnancy due to coronavirus disease 2019 (COVID-19) is a current health priority in developing countries. Identification of clinical and sociodemographic risk factors related to mortality in pregnant women with COVID-19 could guide public policy and encourage such women to accept vaccination. We aimed to evaluate the association of comorbidities and socioeconomic determinants with COVID-19-related mortality and severe disease in pregnant women in Mexico.

### Methods

This is an ongoing nationwide prospective cohort study that includes all pregnant women with a positive reverse-transcription quantitative polymerase chain reaction result for severe acute respiratory syndrome coronavirus 2 (SARS-CoV-2) from the Mexican National Registry of Coronavirus. The primary outcome was maternal death due to COVID-19. The association of comorbidities and socioeconomic characteristics with maternal death was explored using a log-binomial regression model adjusted for possible confounders.

### Results

There were 176 (1.35%) maternal deaths due to COVID-19 among 13 062 consecutive SARS-CoV-2-positive pregnant women. Maternal age, as a continuous (adjusted relative risk (aRR), 1.08 (95% CI, 1.05–1.10)) or categorical variable, was associated with maternal death due to COVID-19; women aged 35–39 years (aRR, 3.16 (95% CI, 2.34–4.26)) or 40 years or older (aRR, 4.07 (95% CI, 2.65–6.25)) had a higher risk for mortality, as compared with those aged < 35 years. Other clinical risk factors associated with maternal mortality were pre-existing diabetes (aRR, 2.66 (95% CI, 1.65–4.27)), chronic hypertension (aRR, 1.75 (95% CI, 1.02–3.00)) and obesity (aRR, 2.15 (95% CI, 1.46–3.17)). Very high social vulnerability (aRR, 1.88 (95% CI, 1.26–2.80)) and high social vulnerability (aRR, 1.49 (95% CI, 1.04–2.13)) were associated with an increased risk of maternal mortality, while very low social vulnerability was associated with a reduced risk (aRR, 0.47 (95% CI, 0.30–0.73)). Being poor or extremely poor were also risk factors for maternal mortality (aRR, 1.53 (95% CI, 1.09–2.15) and aRR, 1.83 (95% CI, 1.32–2.53), respectively).

### Conclusion

This study, which comprises the largest prospective consecutive cohort of pregnant women with COVID-19 to date, has confirmed that advanced maternal age, pre-existing diabetes, chronic hypertension, obesity, high social vulnerability and low socioeconomic status are risk factors for COVID-19-related maternal mortality. © 2021 International Society of Ultrasound in Obstetrics and Gynecology. (Author)

Full URL: <https://doi.org/10.1002/uog.24797>

---

## 2021-14507

**Examining the impact of the COVID-19 pandemic on maternal mental health during pregnancy and the postnatal period.**

McIntosh GC (2022), *MIDIRS Midwifery Digest* vol 32, no 1, March 2022, pp 67-73

By exploring physiological aspects of COVID-19 and its adaptations to pregnancy, this paper will examine its prevalence and physical effects, discussing the ramifications for mental health during pregnancy and the postpartum period. (Author, edited)

---

MIDIRS is part of RCM Information Services Limited which is a company incorporated in England and Wales under company no.11914882 with registered office at 10-18 Union Street, London SE1 1SZ  
RCM Information Services Limited is a subsidiary of The Royal College of Midwives



**The Royal College of Midwives Trust**  
10-18 Union Street  
London  
SE1 1SZ



**Open 24 hours a day, 7 days a week**  
T: 0300 303 0444  
F: +44 20 7312 3536  
E: [info@rcm.org.uk](mailto:info@rcm.org.uk)  
W: [www.rcm.org.uk](http://www.rcm.org.uk)



**Chief Executive:**  
Gill Walton,  
MA, PGDip, BSc Hons, RM  
**President:**  
Rebecca Davies, RM  
**Patron:**  
HRH The Princess Royal

## 2021-14505

**A clearer view to COVID-19 domestic violence and abuse — gaining insight by using a visionary post-feminist lens.** Richards J (2022), MIDIRS Midwifery Digest vol 32, no 1, March 2022, pp 74-80

According to the Office for National Statistics (ONS), domestic abuse will affect 1:4 women and 1:6 men in their lifetime (ONS 2020a) (1). This paper looks at the facts — from a post-feminist perspective — and examines some of the issues, particularly in relation to routine enquiry in midwifery.

As a matter of child protection, issues highlighted in the smooth transition between hospital cot and nursery are discussed, facilitated by the midwife's pivotal role in discharge planning. The long-term repercussions of domestic violence and abuse, jeopardising the developmental health and wellbeing of the child beyond the uterus, are highlighted.

1. Office for National Statistics (ONS) (2020a). Domestic abuse victim characteristics, England and Wales: year ending March 2020. London: ONS.

<https://www.ons.gov.uk/peoplepopulationandcommunity/crimeandjustice/articles/domesticabusevictimcharacteristicsenglandandwales/yearendingmarch2020>.

(Author, edited)

---

## 2021-14499

**Hispanic race is a risk factor for COVID-19 during pregnancy: data from an urban New York City hospital.** Elkafrawi D, Sisti G, Mercado F, et al (2022), Journal of Obstetrics and Gynaecology 12 January 2022, online

There are limited studies on predisposing factors for COVID-19 positivity in asymptomatic pregnant women. The literature published to date on asymptomatic COVID-19 pregnant carriers does not focus on pregnancy or pre-pregnancy comorbidities. We wanted to identify risk factors for COVID-19 in asymptomatic pregnant women. We performed a retrospective chart review of 263 asymptomatic pregnant women admitted to labour and delivery at New York City Health + Hospitals/Lincoln.

We analysed the association between race, body mass index (BMI), smoking, indication for admission, gravidity, parity, pre-pregnancy comorbidity, pregnancy comorbidity via uni- and multivariate statistical tests. Only Hispanic race was significant in the univariate analysis ( $p = .049$ ). At the post-hoc analysis, Hispanics had a higher proportion of COVID-19 cases compared to non-Hispanic Blacks ( $p = .019$ ). No variables were significantly associated with COVID-19 positivity in the multivariate analysis.

Hispanic race appears to be a risk factor for asymptomatic COVID-19 infection during pregnancy. We speculate that the cultural and socioeconomic reality of Hispanic women living in our community leads to more exposure opportunities and therefore, a higher infection rate.

### Impact statement

What is already known on this subject? Little is known on the role of comorbidities and risk factors that can favour COVID-19 infection during pregnancy.

What do the results of this study add? We found that Hispanic pregnant asymptomatic women had a higher rate of COVID-19 in comparison to non-Hispanic Black women. Pre-pregnancy comorbidities such as pregestational diabetes, hypertension and asthma were not associated with COVID-19 positivity.

What are the implications of these findings for clinical practice and/or further research? The reasons why the Hispanic race is more affected by COVID-19 during pregnancy is unclear. The social environment of Hispanic women living in our community, such as their tendency to live in multigenerational and multi-family households, might contribute to a higher infection rate. More resources might be dedicated in the future to Hispanic-dense neighbourhoods. (Author)

---

MIDIRS is part of RCM Information Services Limited which is a company incorporated in England and Wales under company no.11914882 with registered office at 10-18 Union Street, London SE1 1SZ  
RCM Information Services Limited is a subsidiary of The Royal College of Midwives



**The Royal College of Midwives Trust**  
10-18 Union Street  
London  
SE1 1SZ



**Open 24 hours a day, 7 days a week**  
T: 0300 303 0444  
F: +44 20 7312 3536  
E: [info@rcm.org.uk](mailto:info@rcm.org.uk)  
W: [www.rcm.org.uk](http://www.rcm.org.uk)



**Chief Executive:**  
Gill Walton,  
MA, PGDip, BSc Hons, RM  
**President:**  
Rebecca Davies, RM  
**Patron:**  
HRH The Princess Royal

**2021-14492**

**Acute Respiratory Failure and Mechanical Ventilation in Women With COVID-19 During Pregnancy Best Clinical Practices.**

Troiano NH, Richter A, King C (2022), The Journal of Perinatal and Neonatal Nursing vol 36, no 1, January/March 2022, pp 27-36

Symptomatic pregnant women with coronavirus disease-2019 (COVID-19) are at increased risk of severe disease and death compared with symptomatic nonpregnant females of reproductive age. Among those who become critically ill, profound acute hypoxemic respiratory failure is the dominant finding. Significant morbidity and mortality from COVID-19 are largely due to acute viral pneumonia that evolves to acute respiratory distress syndrome. Admission of these patients with critical disease to an intensive care unit and initiation of invasive mechanical ventilation may be indicated. Effective ventilatory support can be challenging in the COVID-19 patient population, even more so when the need occurs in a woman during pregnancy. Key respiratory changes during pregnancy are reviewed. Principles related to maternal-fetal oxygen transport, assessment of ventilation and oxygenation status, and oxygenation goals are also reviewed. Selected concepts related to mechanical ventilatory support for the woman with COVID-19 and acute respiratory failure during pregnancy are presented including indications for ventilatory support, noninvasive support, and invasive ventilator management. Challenges in providing care to this patient population are identified as well as strategies to address them going forward. (Author)

---

**2021-14489**

**Mothers' Experiences of Pregnancy, Labor and Birth, and Postpartum During COVID-19 in the United States: Preliminary Results of a Mixed-Methods Study.**

Saleh L, Canclini S, Greer K, et al (2022), The Journal of Perinatal and Neonatal Nursing vol 36, no 1, January/March 2022, pp 55-67

The purpose of this study was to gain insight into the experiences of women who gave birth in the United States during coronavirus disease-2019 (COVID-19). A mixed-methods study was performed using online surveys and interviews. Data were analyzed using descriptive statistics and interview transcripts were analyzed by thematic analysis resulting in major themes. Participants (n = 32) were women who had given birth on or after March 13, 2020. Of the participants, 34% experienced depression, 46% experienced mild to moderate anxiety, and 28% experienced severe anxiety symptoms. Four major themes emerged: expectations versus reality, early versus late COVID-19 experience, mental distress versus mental health, and healthcare policy versus COVID-19 confusion. Experiences varied based upon geographical location, parity, and proximity to support. Short and long-term effects of COVID-19 on participants and their families were recognized. It is important to acknowledge the confusion experienced in many aspects of the birthing experience due to developing or conflicting pandemic or popular media information. Aligning expectations through providing clear, up-to-date information is helpful in decreasing mental distress. Finally, the impact of COVID-19 highlighted the critical need for professional and focused familial support and follow-up for women experiencing pregnancy-related mental health symptoms. (Author)

**Full URL:** <https://doi.org/10.1097/JPN.0000000000000624>

---

**2021-14486**

**Pandemics Past, Present, and Future: What History Can Teach Us.**

Kriebs JM (2022), The Journal of Perinatal and Neonatal Nursing vol 36, no 1, January/March 2022, pp 7-10

One of the lessons of the current pandemic is that Americans have lost trust in the public health system in the United States (US) and in the health recommendations of the Federal government. History tells us that each pandemic brings new challenges and new lessons. Looking back at the history of pandemics, and at the present experience, nurses and midwives can craft responses to patient concerns and contribute to future planning that better addresses the needs of maternal-child health practices. (Author)

---

MIDIRS is part of RCM Information Services Limited which is a company incorporated in England and Wales under company no.11914882 with registered office at 10-18 Union Street, London SE1 1SZ  
RCM Information Services Limited is a subsidiary of The Royal College of Midwives



**The Royal College of Midwives Trust**  
10-18 Union Street  
London  
SE1 1SZ



**Open 24 hours a day, 7 days a week**  
T: 0300 303 0444  
F: +44 20 7312 3536  
E: [info@rcm.org.uk](mailto:info@rcm.org.uk)  
W: [www.rcm.org.uk](http://www.rcm.org.uk)



**Chief Executive:**  
Gill Walton,  
MA, PGDip, BSc Hons, RM  
**President:**  
Rebecca Davies, RM  
**Patron:**  
HRH The Princess Royal



2021-14290

## Obstetric interventions and pregnancy outcomes during the COVID-19 pandemic in England: A nationwide cohort study.

Gurol-Urganci I, Waite L, Webster K, et al (2022), PLoS Medicine vol 19, no 1, January 2022, e1003884

### Background

The COVID-19 pandemic has disrupted maternity services worldwide and imposed restrictions on societal behaviours. This national study aimed to compare obstetric intervention and pregnancy outcome rates in England during the pandemic and corresponding pre-pandemic calendar periods, and to assess whether differences in these rates varied according to ethnic and socioeconomic background.

### Methods and findings

We conducted a national study of singleton births in English National Health Service hospitals. We compared births during the COVID-19 pandemic period (23 March 2020 to 22 February 2021) with births during the corresponding calendar period 1 year earlier. The Hospital Episode Statistics database provided administrative hospital data about maternal characteristics, obstetric interventions (induction of labour, elective or emergency cesarean section, and instrumental birth), and outcomes (stillbirth, preterm birth, small for gestational age [SGA; birthweight < 10th centile], prolonged maternal length of stay ( $\geq 3$  days), and maternal 42-day readmission). Multi-level logistic regression models were used to compare intervention and outcome rates between the corresponding pre-pandemic and pandemic calendar periods and to test for interactions between pandemic period and ethnic and socioeconomic background. All models were adjusted for maternal characteristics including age, obstetric history, comorbidities, and COVID-19 status at birth. The study included 948,020 singleton births (maternal characteristics: median age 30 years, 41.6% primiparous, 8.3% with gestational diabetes, 2.4% with preeclampsia, and 1.6% with pre-existing diabetes or hypertension); 451,727 births occurred during the defined pandemic period. Maternal characteristics were similar in the pre-pandemic and pandemic periods. Compared to the pre-pandemic period, stillbirth rates remained similar (0.36% pandemic versus 0.37% pre-pandemic,  $p = 0.16$ ). Preterm birth and SGA birth rates were slightly lower during the pandemic (6.0% versus 6.1% for preterm births, adjusted odds ratio [aOR] 0.96, 95% CI 0.94–0.97; 5.6% versus 5.8% for SGA births, aOR 0.95, 95% CI 0.93–0.96; both  $p < 0.001$ ). Slightly higher rates of obstetric intervention were observed during the pandemic (40.4% versus 39.1% for induction of labour, aOR 1.04, 95% CI 1.03–1.05; 13.9% versus 12.9% for elective cesarean section, aOR 1.13, 95% CI 1.11–1.14; 18.4% versus 17.0% for emergency cesarean section, aOR 1.07, 95% CI 1.06–1.08; all  $p < 0.001$ ). Lower rates of prolonged maternal length of stay (16.7% versus 20.2%, aOR 0.77, 95% CI 0.76–0.78,  $p < 0.001$ ) and maternal readmission (3.0% versus 3.3%, aOR 0.88, 95% CI 0.86–0.90,  $p < 0.001$ ) were observed during the pandemic period. There was some evidence that differences in the rates of preterm birth, emergency cesarean section, and unassisted vaginal birth varied according to the mother's ethnic background but not according to her socioeconomic background. A key limitation is that multiple comparisons were made, increasing the chance of false-positive results.

### Conclusions

In this study, we found very small decreases in preterm birth and SGA birth rates and very small increases in induction of labour and elective and emergency cesarean section during the COVID-19 pandemic, with some evidence of a slightly different pattern of results in women from ethnic minority backgrounds. These changes in obstetric intervention rates and pregnancy outcomes may be linked to women's behaviour, environmental exposure, changes in maternity practice, or reduced staffing levels.

(Author)

Full URL: <https://doi.org/10.1371/journal.pmed.1003884>

MIDIRS is part of RCM Information Services Limited which is a company incorporated in England and Wales under company no.11914882 with registered office at 10-18 Union Street, London SE1 1SZ  
RCM Information Services Limited is a subsidiary of The Royal College of Midwives



**The Royal College of Midwives Trust**  
10-18 Union Street  
London  
SE1 1SZ



**Open 24 hours a day, 7 days a week**  
T: 0300 303 0444  
F: +44 20 7312 3536  
E: [info@rcm.org.uk](mailto:info@rcm.org.uk)  
W: [www.rcm.org.uk](http://www.rcm.org.uk)



**Chief Executive:**  
Gill Walton,  
MA, PGDip, BSc Hons, RM  
**President:**  
Rebecca Davies, RM  
**Patron:**  
HRH The Princess Royal

## 2021-14144

**Coronavirus: Antenatal Appointments [written answer].** Scottish Parliament (2022), Official Report Written question S6W-05853, 24 January 2022

Maree Todd responds to a written question asked by Alexander Burnett to the Scottish Government, regarding when it expects COVID-19 restrictions to be eased to allow pregnant mothers to take their young children with them when attending midwife appointments, in order to alleviate the need to find alternative childcare. (MB)

**Full URL:** <https://archive2021.parliament.scot/parliamentarybusiness/28877.aspx?SearchType=Advance&ReferenceNumbers=S6W-05853>

---

## 2021-13995

**Mental well-being during stages of COVID-19 lockdown among pregnant women and new mothers.** Overbeck G, Rasmussen IS, Siersma V, et al (2022), BMC Pregnancy and Childbirth vol 22, no 92, 1 February 2022

### Background

Pregnancy and early motherhood are sensitive times where epidemic disease outbreaks can affect mental health negatively. Countries and health care systems handled the pandemic and lockdowns differently and knowledge about how the COVID-19 pandemic affected the mental well-being of pregnant women and new mothers is limited and points in different directions.

### Aim

To investigate symptoms of anxiety and depression in a population of pregnant women and new mothers in various stages of infection pressure and lockdown during the first 15 months of the COVID-19 pandemic in Denmark.

### Methods

The study population was nested an inception cohort of women recruited in their first trimester of pregnancy. Data about mental health of the woman were obtained in relation to pregnancy and child development (first trimester, 8 weeks postpartum and 5 months postpartum), and data were analysed cross-sectionally according to calendar time (periods defined by infection rate and lock-down during the COVID-19 pandemic).

### Results

No differences in reported levels of depressive symptoms between the six examined time periods of the pandemic were observed. Specifically, symptoms remained unchanged after the first lock-down. No major changes in anxiety symptoms were observed in relation to increased infection pressure or lockdowns, but a small increase was observed during the second lockdown in women 8 weeks postpartum.

### Conclusion

No clear change in mood among pregnant women was seen between during the stages of COVID-19 pandemic in Denmark. (Author)

**Full URL:** <https://doi.org/10.1186/s12884-021-04374-4>

---

MIDIRS is part of RCM Information Services Limited which is a company incorporated in England and Wales under company no.11914882 with registered office at 10-18 Union Street, London SE1 1SZ  
RCM Information Services Limited is a subsidiary of The Royal College of Midwives



**The Royal College of Midwives Trust**  
10-18 Union Street  
London  
SE1 1SZ



**Open 24 hours a day, 7 days a week**  
T: 0300 303 0444  
F: +44 20 7312 3536  
E: [info@rcm.org.uk](mailto:info@rcm.org.uk)  
W: [www.rcm.org.uk](http://www.rcm.org.uk)



**Chief Executive:**  
Gill Walton,  
MA, PGDip, BSc Hons, RM  
**President:**  
Rebecca Davies, RM  
**Patron:**  
HRH The Princess Royal

**2021-13988**

**The mental health crisis of expectant women in the UK: effects of the COVID-19 pandemic on prenatal mental health, antenatal attachment and social support.** Filippetti ML, Clarke ADF, Rigato S (2022), BMC Pregnancy and Childbirth vol 22, no 68, 26 January 2022

#### Background

Pregnancy has been shown to be times in a woman's life particularly prone to mental health issues, however a substantial percentage of mothers report subclinical perinatal mental health symptoms that go undetected. Experiences of prenatal trauma, such as the COVID-19 pandemic, may exacerbate vulnerability to negative health outcomes for pregnant women and their infants. We aimed to examine the role of: 1) anxiety, depression, and stress related to COVID-19 in predicting the quality of antenatal attachment; 2) perceived social support and COVID-19 appraisal in predicting maternal anxiety and depression.

#### Methods

A sample of 150 UK expectant women were surveyed during the COVID-19 pandemic. Questions included demographics, pregnancy details, and COVID-19 appraisal. Validated measures were used to collect self-reported maternal antenatal attachment (MAAS), symptoms of anxiety (STAI), depression (BDI-II), and stress related to the psychological impact of COVID-19 (IES-r).

#### Results

We found that the pandemic has affected UK expectant mothers' mental health by increasing prevalence of depression (47%), anxiety (60%) and stress related to the psychological impact of COVID-19 (40%). Women for whom COVID-19 had a higher psychological impact were more likely to suffer from depressive (95% HDPI = [0.04, 0.39]) and anxiety symptoms (95% HPDI = [0.40, 0.69]). High depressive symptoms were associated with reduced attachment to the unborn baby (95% HPDI [-0.46, -0.1]). Whilst women who appraised the impact of COVID-19 to be more negative showed higher levels of anxiety (HPDI = [0.15, 0.46]), higher social support acted as a protective factor and was associated with lower anxiety (95% HPDI = [-0.52, -0.21]).

#### Conclusions

The current findings demonstrate that direct experience of prenatal trauma, such as the one experienced during the COVID-19 pandemic, significantly amplifies mothers' vulnerability to mental health symptoms and impairs the formation of a positive relationship with their unborn baby. Health services should prioritise interventions strategies aimed at fostering support for pregnant women. (Author)

**Full URL:** <https://doi.org/10.1186/s12884-022-04387-7>

---

MIDIRS is part of RCM Information Services Limited which is a company incorporated in England and Wales under company no.11914882 with registered office at 10-18 Union Street, London SE1 1SZ  
RCM Information Services Limited is a subsidiary of The Royal College of Midwives



**The Royal College of Midwives Trust**  
10-18 Union Street  
London  
SE1 1SZ



**Open 24 hours a day, 7 days a week**  
T: 0300 303 0444  
F: +44 20 7312 3536  
E: [info@rcm.org.uk](mailto:info@rcm.org.uk)  
W: [www.rcm.org.uk](http://www.rcm.org.uk)



**Chief Executive:**  
Gill Walton,  
MA, PGDip, BSc Hons, RM  
**President:**  
Rebecca Davies, RM  
**Patron:**  
HRH The Princess Royal

**2021-13962**

**The experience of women with recent gestational diabetes during the COVID-19 lockdown: a qualitative study from Denmark.** Jensen NH, Nielsen KK, Dahl-Petersen IK, et al (2022), BMC Pregnancy and Childbirth vol 22, no 84, 29 January 2022

**Background**

Following COVID-19 and the lockdowns, maternity care and support for women after delivery have been temporary restructured. Studies show that COVID-19 adversely impacts pregnant and peripartum women in the general population, but experiences among women in the first year after delivery/in the wider postpartum period remain unexplored. Moreover, experiences among women with recent gestational diabetes mellitus (GDM) are lacking; though it is a group with a potential high need for support after delivery. The aim of our study was to investigate (i) how women with recent GDM experienced COVID-19 and the first lockdown in Denmark, and (ii) the women's risk perception and health literacy in terms of interaction with the healthcare system in relation to COVID-19.

**Methods**

We performed a qualitative study among 11 women with recent GDM (infants aged 2-11 months old). Semi-structured interviews were conducted in April-May 2020 by telephone or Skype for Business, when Denmark was under lockdown. We analysed data using a thematic qualitative content analysis.

**Results**

Three themes emerged: i) Everyday life and family well-being, ii) Worries about COVID-19 and iii) Health literacy: Health information and access to healthcare. The women were generally not worried about their own or their infant's risk of COVID-19. The lockdown had a negative impact on everyday life e.g. routines, loneliness, breastfeeding uncertainties and worries for the infant's social well-being; but better family dynamics were also described. It was challenging to maintain healthy behaviours and thus the women described worries for the risk of type 2 diabetes and GDM in subsequent pregnancies. The women missed peer support and face-to-face visits from health visitors and found it difficult to navigate the restructured care with online/telephone set-ups.

**Conclusions**

COVID-19 and the lockdown affected everyday life among women with recent GDM both positively and negatively. Our findings suggest a need for care that are responsive to psychological and social aspects of health throughout the COVID-19 pandemic and support to limit worries about adaptation to motherhood and the infant's social well-being. Communication focusing on the importance and relevance of contacting healthcare providers should also be strengthened. (Author)

**Full URL:** <https://doi.org/10.1186/s12884-022-04424-5>

---

MIDIRS is part of RCM Information Services Limited which is a company incorporated in England and Wales under company no.11914882 with registered office at 10-18 Union Street, London SE1 1SZ  
RCM Information Services Limited is a subsidiary of The Royal College of Midwives



**The Royal College of Midwives Trust**  
10-18 Union Street  
London  
SE1 1SZ



**Open 24 hours a day, 7 days a week**  
T: 0300 303 0444  
F: +44 20 7312 3536  
E: [info@rcm.org.uk](mailto:info@rcm.org.uk)  
W: [www.rcm.org.uk](http://www.rcm.org.uk)



**Chief Executive:**  
Gill Walton,  
MA, PGDip, BSc Hons, RM  
**President:**  
Rebecca Davies, RM  
**Patron:**  
HRH The Princess Royal

## 2021-13961

**Effects of the COVID-19 pandemic on perinatal outcomes: a retrospective cohort study from Turkey.** Yalçın SS, Boran P, Tezel B, et al (2022), BMC Pregnancy and Childbirth vol 22, no 51, 20 January 2022

### Background

Lockdowns, pregnant women's fear from hospitalization in addition to uncertainties about appropriate birthing practices at the beginning of the pandemic may have affected the health outcomes of mother-infant couples. We aimed to explore whether pregnancy outcomes including the rates of cesarean delivery (CS), preterm, and low birth weight (LBW) births have changed during the pandemic period compared with the pre-pandemic period.

### Methods

We applied a population-based retrospective cohort, before-after approach in 2020 vs. similar calendar months in 2019 for five periods [Jan-Feb (pre-pandemic); March–May (1st wave and lockdown); June–August; September–October; November–December (2nd wave and lockdown)]. The data was modelled through multiple logistic regressions using key outcomes; CS, preterm, and LBW births as the dependent variables, and adjustments were made for independent variables in SPSS software. We evaluated the modification of years by periods by adding interaction term (yearXperiod) to the model.

### Results

The rate of CS in hospital births increased from 57.7% in 2019 to 60.2% in 2020. CS rates were significantly increased during the 3rd and 4th periods. The overall preterm rate was 11%. When singleton pregnancies were considered, adjusted multivariable analyses showed a decrease in preterm proportions during all time periods with respect to the pre-pandemic period. The percentage of LBW was 7.7% during the pandemic period and was found to be significantly reduced compared to the pre-pandemic period. There was a significant reduction in LBW rates in all periods except the second lockdown period.

### Conclusions

Our findings suggested significant reductions in preterm and LBW births possibly due to the indirect effects of the pandemic. Moreover, strategies need to be considered to address the increased CS rates and shifting of maternity service utilization to private facilities. (Author)

**Full URL:** <https://doi.org/10.1186/s12884-021-04349-5>

---

MIDIRS is part of RCM Information Services Limited which is a company incorporated in England and Wales under company no.11914882 with registered office at 10-18 Union Street, London SE1 1SZ  
RCM Information Services Limited is a subsidiary of The Royal College of Midwives



**The Royal College of Midwives Trust**  
10-18 Union Street  
London  
SE1 1SZ



**Open 24 hours a day, 7 days a week**  
T: 0300 303 0444  
F: +44 20 7312 3536  
E: [info@rcm.org.uk](mailto:info@rcm.org.uk)  
W: [www.rcm.org.uk](http://www.rcm.org.uk)



**Chief Executive:**  
Gill Walton,  
MA, PGDip, BSc Hons, RM  
**President:**  
Rebecca Davies, RM  
**Patron:**  
HRH The Princess Royal

## 2021-13958

**Effect of initial COVID-19 outbreak during first trimester on pregnancy outcome in Wuxi, China.** Liu Y, Dai M, Tang S (2022), BMC Pregnancy and Childbirth vol 22, no 54, 21 January 2022

### Background

A hospital-based retrospective study was conducted to examine the effect of initial COVID-19 outbreak during first trimester on pregnancy outcome in Wuxi, China.

### Methods

Women who delivered children at our hospital during June 2020 to July 2020 (control group), and October 2020 to December 2020 (exposure group) were recruited in the present study. All of the participants were not infected with COVID-19. The last menstrual period (LMP) of the exposure group was between January 24th, 2020 and March 12th, 2020, whilst in the control group, the LMP was between May 12th and October 31st, 2019.

### Results

There were 1,456 women in the exposure group and 1,816 women in the control group. Women in the exposure group were more susceptible to hypertension during pregnancy (HDP,  $P = 0.004$ ,  $OR[95\%CI] = 1.90[1.22-2.95]$ ) and gestational diabetes mellitus (GDM,  $P = 0.008$ ,  $OR[95\%CI] = 1.31[1.08-1.60]$ ) compared to those in the control group. Mothers diagnosed with HDP were more likely to deliver premature infants, leading to a higher rate of low birth weight (all  $P < 0.05$ ). The other common outcomes of pregnancy showed no statistical differences between the two groups.

### Conclusions

The initial COVID-19 outbreak might increase the incidence rates of HDP and GDM among pregnant women whose first trimesters were during that period, resulting in higher percentages of premature delivery and low birth weight. These results should be confirmed by studies from other hospitals or cities. (Author)

**Full URL:** <https://doi.org/10.1186/s12884-022-04395-7>

---

MIDIRS is part of RCM Information Services Limited which is a company incorporated in England and Wales under company no.11914882 with registered office at 10-18 Union Street, London SE1 1SZ  
RCM Information Services Limited is a subsidiary of The Royal College of Midwives



**The Royal College of Midwives Trust**  
10-18 Union Street  
London  
SE1 1SZ



**Open 24 hours a day, 7 days a week**  
T: 0300 303 0444  
F: +44 20 7312 3536  
E: [info@rcm.org.uk](mailto:info@rcm.org.uk)  
W: [www.rcm.org.uk](http://www.rcm.org.uk)



**Chief Executive:**  
Gill Walton,  
MA, PGDip, BSc Hons, RM  
**President:**  
Rebecca Davies, RM  
**Patron:**  
HRH The Princess Royal

**2021-13857**

**Companionship for women/birthing people using antenatal and intrapartum care in England during COVID-19: a mixed-methods analysis of national and organisational responses and perspectives.** Thomson G, Balaam M-C, Harris RN, et al (2022), *BMJ Open* vol 12, no 1, January 2022, e051965

Objectives To explore stakeholders' and national organisational perspectives on companionship for women/birthing people using antenatal and intrapartum care in England during COVID-19, as part of the Achieving Safe and Personalised maternity care In Response to Epidemics (ASPIRE) COVID-19 UK study.

Setting Maternity care provision in England.

Participants Interviews were held with 26 national governmental, professional and service-user organisation leads (July–December 2020). Other data included public-facing outputs logged from 25 maternity Trusts (September/October 2020) and data extracted from 78 documents from eight key governmental, professional and service-user organisations that informed national maternity care guidance and policy (February–December 2020).

Results Six themes emerged: 'Postcode lottery of care' highlights variations in companionship and visiting practices between trusts/locations, 'Confusion and stress around 'rules'' relates to a lack of and variable information concerning companionship/visiting, 'Unintended consequences' concerns the negative impacts of restricted companionship or visiting on women/birthing people and staff, 'Need for flexibility' highlights concerns about applying companionship and visiting policies irrespective of need, 'Acceptable' time for support' highlights variations in when and if companionship was 'allowed' antenatally and intrapartum and 'Loss of human rights for gain in infection control' emphasises how a predominant focus on infection control was at a cost to psychological safety and human rights.

Conclusions Policies concerning companionship and visiting have been inconsistently applied within English maternity services during the COVID-19 pandemic. In some cases, policies were not justified by the level of risk, and were applied indiscriminately regardless of need. There is an urgent need to determine how to sensitively and flexibly balance risks and benefits and optimise outcomes during the current and future crisis situations. (Author)

Full URL: <http://dx.doi.org/10.1136/bmjopen-2021-051965>

---

**2021-13835**

**A real-world assessment of tolerability and treatment outcomes of COVID-19 monoclonal antibodies administered in pregnancy.** Chang MH, Cowman K, Guo Y, et al (2022), *American Journal of Obstetrics & Gynecology (AJOG)* vol 226, no 5, May 2022, pp 743-745

Research letter aiming to evaluate the tolerability of monoclonal antibodies for treatment of COVID-19 in pregnancy and to assess subjective improvement in symptoms. Results indicate that monoclonal antibodies are well tolerated and should be considered in pregnancy. (LDO)

Full URL: <https://doi.org/10.1016/j.ajog.2022.01.018>

---

**2021-13830**

**Changes and geographic variation in rates of preterm birth and stillbirth during the prepandemic period and COVID-19 pandemic, according to health insurance claims in the United States, April–June 2019 and April–June 2020.** Chen J, Ferre C, Ouyang L, et al (2022), *American Journal of Obstetrics & Gynecology MFM* vol 4, no 1, January 2022, 100508

Research letter comparing premature birth and stillbirth rates between the pre-pandemic period and COVID-19 lockdown period in the United States. Results show a 0.4% decrease in premature birth and no changes in stillbirth rates during the COVID-19 pandemic. (LDO)

Full URL: <https://doi.org/10.1016/j.ajogmf.2021.100508>

---

MIDIRS is part of RCM Information Services Limited which is a company incorporated in England and Wales under company no.11914882 with registered office at 10-18 Union Street, London SE1 1SZ  
RCM Information Services Limited is a subsidiary of The Royal College of Midwives



**The Royal College of Midwives Trust**  
10-18 Union Street  
London  
SE1 1SZ



**Open 24 hours a day, 7 days a week**  
T: 0300 303 0444  
F: +44 20 7312 3536  
E: [info@rcm.org.uk](mailto:info@rcm.org.uk)  
W: [www.rcm.org.uk](http://www.rcm.org.uk)



**Chief Executive:**  
Gill Walton,  
MA, PGDip, BSc Hons, RM  
**President:**  
Rebecca Davies, RM  
**Patron:**  
HRH The Princess Royal

## 2021-13814

**COVID-19 pandemic and population-level pregnancy and neonatal outcomes in general population: A living systematic review and meta-analysis (Update#2: November 20, 2021).** Yang J, D'Souza R, Kharrat A, et al (2022), *Acta Obstetrica et Gynecologica Scandinavica* vol 101, no 3, March 2022, pp 273-292

### Introduction

Conflicting reports of increases and decreases in rates of preterm birth (PTB) and stillbirth in the general population during the coronavirus disease 2019 (COVID-19) pandemic have surfaced. The objective of our study was to conduct a living systematic review and meta-analyses of studies reporting pregnancy and neonatal outcomes by comparing the pandemic and pre-pandemic periods.

### Material and methods

We searched the PubMed and Embase databases and reference lists of articles published up until November 20, 2021, and included English language studies that compared outcomes between the COVID-19 pandemic time period with pre-pandemic time periods. Risk of bias was assessed using the Newcastle-Ottawa scale. We conducted random-effects meta-analysis using the inverse variance method.

### Results

Fifty-two studies with low-to-moderate risk of bias, reporting on 2 372 521 pregnancies during the pandemic period and 28 518 300 pregnancies during the pre-pandemic period, were included. There was significant reduction in unadjusted estimates of PTB (43 studies, unadjusted odds ratio [uaOR] 0.95, 95% CI 0.93–0.98), but not in adjusted estimates (five studies, adjusted OR [aOR] 0.94, 95% CI 0.74–1.19). This reduction was noted in studies from single centers/health areas (29 studies, uaOR 0.90, 95% CI 0.85–0.94) but not in regional/national studies (14 studies, uaOR 0.99, 95% CI 0.99–1.01). There was reduction in spontaneous PTB (nine studies, uaOR 0.91, 95% CI 0.88–0.94) but not in induced PTB (eight studies, uaOR 0.90, 95% CI 0.79–1.01). There was no difference in the odds of stillbirth between the pandemic and pre-pandemic time periods (32 studies, uaOR 1.07, 95% CI 0.97–1.18 and three studies, aOR 1.18, 95% CI 0.86–1.63). There was an increase in mean birthweight during the pandemic period compared with the pre-pandemic period (nine studies, mean difference 21 g, 95% CI 13–30 g). The odds of maternal mortality were increased (five studies, uaOR 1.15, 95% CI 1.05–1.26); however, only unadjusted estimates were available, and the result was mostly influenced by one study from Mexico. There was significant publication bias for the outcome of PTB.

### Conclusions

The COVID-19 pandemic may be associated with a reduction in PTB; however, referral bias cannot be excluded. There was no statistically significant difference in stillbirths between pandemic and pre-pandemic periods. (Author)

Full URL: <https://doi.org/10.1111/aogs.14318>

## 2021-13805

**SARS-CoV-2 infection and a subsequent secondary atrophy/atresia of one of the umbilical arteries.** Ignatov PN, Neykova KN (2022), *The Journal of Maternal-Fetal and Neonatal Medicine* 23 January 2022, online

We conducted a retrospective study among patients who visited two tertiary clinical settings between April 2020 and July 2021. Diagnosis of a single umbilical artery (SUA) was made on four patients during the 18th-23rd week fetal anatomy scan after the previous 11th-13th week scan records proved affirmative for the presence of two umbilical arteries (UAs). (Author, edited)

MIDIRS is part of RCM Information Services Limited which is a company incorporated in England and Wales under company no.11914882 with registered office at 10-18 Union Street, London SE1 1SZ  
RCM Information Services Limited is a subsidiary of The Royal College of Midwives



**The Royal College of Midwives Trust**  
10-18 Union Street  
London  
SE1 1SZ



**Open 24 hours a day, 7 days a week**  
T: 0300 303 0444  
F: +44 20 7312 3536  
E: [info@rcm.org.uk](mailto:info@rcm.org.uk)  
W: [www.rcm.org.uk](http://www.rcm.org.uk)



**Chief Executive:**  
Gill Walton,  
MA, PGDip, BSc Hons, RM  
**President:**  
Rebecca Davies, RM  
**Patron:**  
HRH The Princess Royal



## 2021-13698

**Severe COVID-19 in pregnancy is almost exclusively limited to unvaccinated women – time for policies to change.** Engjom H, van den Akker T, Aabakke A, et al (2022), The Lancet Regional Health - Europe 26 January 2022, online  
Commentary piece analysing data from the International Network of Obstetric Survey Systems (INOSS) collected in the United Kingdom (UK), the Netherlands, Norway, Denmark, Finland and Italy between May and December 2021. Results show that at least 80% of pregnant women admitted to critical care with COVID-19 were unvaccinated across the six countries, including 98% of pregnant women in the UK. (LDO)

**Full URL:** <https://doi.org/10.1016/j.lanepe.2022.100313>

---

## 2021-13621

**Spontaneous Intracerebral Hemorrhage (ICH) associated with pregnancy and SARS-CoV-2 infection: a case report.** Dini P, Aminimoghaddam S, Mirzaasgari Z, et al (2022), BMC Pregnancy and Childbirth vol 22, no 14, 6 January 2022

### Background

Coronavirus Disease 2019 (COVID-19) is predominately known as a respiratory disease associated with pneumonia, acute respiratory distress syndrome and multiorgan failure. However, extra-pulmonary complications of severe acute respiratory syndrome coronavirus 2 (SARS-CoV-2) are increasingly being recognized. In this regard, some studies implied the hemostatic and vascular involvements in patients with SARS-CoV-2 infection.

### Case presentation

We describe a case of spontaneous Intracerebral Hemorrhage (ICH) in a pregnant patient with COVID-19 and history of cesarean section a week before the occurrence of ICH. The patient underwent emergent craniotomy with acceptable outcome. Hemorrhagic events, including ICH, may happen during COVID-19 infection with several possible mechanisms.

### Conclusion

COVID-19 patients, especially high-risk groups, are at a risk of intracranial hemorrhage. Therefore, close follow-up must be maintained and hemorrhagic events must be kept in mind in these cases. (Author)

**Full URL:** <https://doi.org/10.1186/s12884-021-04345-9>

---

## 2021-13581

**Mind the Gap: COVID-19 highlights the research void in pregnancy.** Dangel A, Yu V, Liang C, et al (2022), American Journal of Obstetrics & Gynecology MFM vol 4, no 3, May 2022, 100566

Discusses the lack of research involving pregnant women and suggests that the COVID-19 pandemic can be used as an opportunity to evaluate the research landscape. (LDO)

**Full URL:** <https://doi.org/10.1016/j.ajogmf.2022.100566>

---

MIDIRS is part of RCM Information Services Limited which is a company incorporated in England and Wales under company no.11914882 with registered office at 10-18 Union Street, London SE1 1SZ  
RCM Information Services Limited is a subsidiary of The Royal College of Midwives



**The Royal College of Midwives Trust**  
10-18 Union Street  
London  
SE1 1SZ



**Open 24 hours a day, 7 days a week**  
T: 0300 303 0444  
F: +44 20 7312 3536  
E: [info@rcm.org.uk](mailto:info@rcm.org.uk)  
W: [www.rcm.org.uk](http://www.rcm.org.uk)



**Chief Executive:**  
Gill Walton,  
MA, PGDip, BSc Hons, RM  
**President:**  
Rebecca Davies, RM  
**Patron:**  
HRH The Princess Royal

## 2021-13570

**Twin-to-twin transfusion syndrome and coronavirus disease 2019: impact on diagnosis, referral, eligibility for fetoscopic laser therapy, and outcomes.** López-Briones H, Villalobos-Gómez R, Chávez-González E, et al (2022), *AJOG Global Reports* vol 2, no 1, February 2022, 100040

### Background

Due to the progressive nature of twin-to-twin transfusion syndrome (TTTS), hindering on health care access during the coronavirus disease (COVID-19) pandemic may lead to delayed diagnosis and referral to fetal surgery centers, which may have repercussions on outcomes.

### Objective

To assess the clinical impact of the COVID-19 pandemic on pregnancies complicated with TTTS.

### Study Design

A retrospective cohort of consecutive monochorionic diamniotic twin pregnancies complicated with TTTS evaluated in our national referral fetal surgery center at Queretaro, Mexico for possible surgical fetoscopy was constructed. Maternal-fetal characteristics and perinatal outcomes of cases evaluated during the first year of the World Health Organization's COVID-19 pandemic declaration (11 March 2020- 10 March 2021) were retrospectively compared with cases evaluated during the same period in the previous year (11 March 2019 – 10 March 2020).

### Results

109 consecutive TTTS cases were evaluated during the 2-year study period, 54 during the COVID-19 pandemic and 55 in the previous year. In the former group, a higher proportion of cases with fetal surveillance interval longer than 2 weeks (70.4% vs. 47.3%,  $p=0.01$ ), TTTS complications precluding laser therapy such as intrauterine fetal demise, preterm rupture of membranes, or cervical dilatation with prolapsed amniotic membranes (18.5% vs. 1.8%,  $p<0.01$ ), advanced TTTS (53.7% vs. 36.4%,  $p=0.07$ ), preoperative short cervix (25.9% vs. 10.9%,  $p<0.05$ ), and lower overall perinatal survival (56.9% vs. 80.0%,  $p=0.01$  of at least one twin; and 39.2% vs. 56.4%,  $p=0.08$  of both twins, respectively) were observed. A significantly lower number of cases were selected for fetoscopic laser therapy during the pandemic period (75.9% vs. 92.7%,  $p=0.01$ ), with similar postoperative outcomes seen in both study periods.

### Conclusion

In pregnancies with TTTS, the COVID-19 pandemic has shown an adverse impact involving suboptimal fetal surveillance, advanced stages at diagnosis, poorer survival rates, and higher number of complications that preclude fetoscopic laser therapy. (Author)

**Full URL:** <https://doi.org/10.1016/j.xagr.2021.100040>

---

## 2021-13520

**Covid-19: Severe complications during pregnancy are more common in unvaccinated women, study finds.** Mahase E (2022), *BMJ* vol 376, no 8322, 17 January 2022, o117

Unvaccinated women accounted for 77% of SARS-CoV-2 infections that have occurred during pregnancy in Scotland and 98% of infections that led to a critical care admission, a study has found (1).

1. Stock SJ et al. *Nature Medicine*, 13 January 2022, online. <https://doi.org/10.1038/s41591-021-01666-2>.

(Author, edited)

**Full URL:** <https://doi.org/10.1136/bmj.o117>

---

MIDIRS is part of RCM Information Services Limited which is a company incorporated in England and Wales under company no.11914882 with registered office at 10-18 Union Street, London SE1 1SZ  
RCM Information Services Limited is a subsidiary of The Royal College of Midwives



**The Royal College of Midwives Trust**  
10-18 Union Street  
London  
SE1 1SZ



**Open 24 hours a day, 7 days a week**  
T: 0300 303 0444  
F: +44 20 7312 3536  
E: [info@rcm.org.uk](mailto:info@rcm.org.uk)  
W: [www.rcm.org.uk](http://www.rcm.org.uk)



**Chief Executive:**  
Gill Walton,  
MA, PGDip, BSc Hons, RM  
**President:**  
Rebecca Davies, RM  
**Patron:**  
HRH The Princess Royal

## 2021-13447

**Impact of COVID-19 on pregnancy worry in the United States.** Burgess A, Breman RB, Roane LA, et al (2022), Birth 7 January 2022, online

### Background

Several recent studies confirm that the COVID-19 pandemic has increased symptoms of stress, anxiety, and depression in pregnant persons around world. In this study, we aimed to uncover the impact of COVID-19 on worry during pregnancy.

### Methods

This study used a cross-sectional descriptive research design. A link to a survey was emailed to users of the Ovia Pregnancy app. Participants (N = 253) completed the survey, which included the Cambridge Worry in Pregnancy Scale, and answered several free text questions. The free-text questions were included to capture worries not listed on the scale and to allow participants the opportunity to more clearly describe COVID-19-related worries. Descriptive statistics were reported, and content analysis performed to determine themes.

### Results

Overall, respondents reported they were quite or very worried about having their partner with them at birth (31.7%, n = 80), giving birth (28.2%; n = 71), and something being wrong with the baby (27.3%; n = 69). Results on worries also differed by participants' race, parity, and trimester. When comparing White to other racial groups, other racial groups had statistically significantly higher median scores for questions on worries about employment (P = .001), going to the hospital (P = .002), and internal examinations (P = .03). Content analysis revealed isolation, loss of support, anxiety/stress, and grief as major themes.

### Discussion

The worry, isolation, loss of support, anxiety, and grief reported by pregnant persons during the COVID-19 pandemic may impact maternal pre- and postnatal mental health and are not borne equally. Birthing persons of color appear disproportionately impacted. Prenatally, maternity care providers should assess for worry and provide individualized education and resources to pregnant patients, centering individuals and communities made most vulnerable by structural inequality. (Author)

Full URL: <https://doi.org/10.1111/birt.12608>

---

## 2021-13441

**Covid in pregnancy linked to birth-related complications.** Anon (2022), BBC News 13 January 2022

A new study (1) has linked Covid-19 to complications during pregnancy.

1. Stock SJ. Nature Medicine, 13 January 2022, online. <https://doi.org/10.1038/s41591-021-01666-2>.

(Author, edited)

Full URL: <https://www.bbc.co.uk/news/uk-scotland-59986452>

---

MIDIRS is part of RCM Information Services Limited which is a company incorporated in England and Wales under company no.11914882 with registered office at 10-18 Union Street, London SE1 1SZ  
RCM Information Services Limited is a subsidiary of The Royal College of Midwives



**The Royal College of Midwives Trust**  
10-18 Union Street  
London  
SE1 1SZ



**Open 24 hours a day, 7 days a week**  
T: 0300 303 0444  
F: +44 20 7312 3536  
E: [info@rcm.org.uk](mailto:info@rcm.org.uk)  
W: [www.rcm.org.uk](http://www.rcm.org.uk)



**Chief Executive:**  
Gill Walton,  
MA, PGDip, BSc Hons, RM  
**President:**  
Rebecca Davies, RM  
**Patron:**  
HRH The Princess Royal

## 2021-13440

**SARS-CoV-2 infection and COVID-19 vaccination rates in pregnant women in Scotland.** Stock SJ, Carruthers J, Calvert C, et al (2022), Nature Medicine 13 January 2022, online

Population-level data on COVID-19 vaccine uptake in pregnancy and SARS-CoV-2 infection outcomes are lacking. We describe COVID-19 vaccine uptake and SARS-CoV-2 infection in pregnant women in Scotland, using whole-population data from a national, prospective cohort. Between the start of a COVID-19 vaccine program in Scotland, on 8 December 2020 and 31 October 2021, 25,917 COVID-19 vaccinations were given to 18,457 pregnant women. Vaccine coverage was substantially lower in pregnant women than in the general female population of 18–44 years; 32.3% of women giving birth in October 2021 had two doses of vaccine compared to 77.4% in all women. The extended perinatal mortality rate for women who gave birth within 28 d of a COVID-19 diagnosis was 22.6 per 1,000 births (95% CI 12.9–38.5; pandemic background rate 5.6 per 1,000 births; 452 out of 80,456; 95% CI 5.1–6.2). Overall, 77.4% (3,833 out of 4,950; 95% CI 76.2–78.6) of SARS-CoV-2 infections, 90.9% (748 out of 823; 95% CI 88.7–92.7) of SARS-CoV-2 associated with hospital admission and 98% (102 out of 104; 95% CI 92.5–99.7) of SARS-CoV-2 associated with critical care admission, as well as all baby deaths, occurred in pregnant women who were unvaccinated at the time of COVID-19 diagnosis. Addressing low vaccine uptake rates in pregnant women is imperative to protect the health of women and babies in the ongoing pandemic. (Author)

**Full URL:** <https://doi.org/10.1038/s41591-021-01666-2>

---

MIDIRS is part of RCM Information Services Limited which is a company incorporated in England and Wales under company no.11914882 with registered office at 10-18 Union Street, London SE1 1SZ  
RCM Information Services Limited is a subsidiary of The Royal College of Midwives



**The Royal College of Midwives Trust**  
10-18 Union Street  
London  
SE1 1SZ



**Open 24 hours a day, 7 days a week**  
T: 0300 303 0444  
F: +44 20 7312 3536  
E: [info@rcm.org.uk](mailto:info@rcm.org.uk)  
W: [www.rcm.org.uk](http://www.rcm.org.uk)



**Chief Executive:**  
Gill Walton,  
MA, PGDip, BSc Hons, RM  
**President:**  
Rebecca Davies, RM  
**Patron:**  
HRH The Princess Royal

**2021-13078**

**Systematic review and meta-analysis of COVID-19 maternal and neonatal clinical features and pregnancy outcomes up to June 3, 2021.** Marchand G, Patil AS, Masoud AT, et al (2022), *AJOG Global Reports* vol 2, no 1, February 2022, 100049

#### Objectives

COVID-19 is a rapidly changing and developing emergency that requires constant re-evaluation of available data. We report a systematic review and meta analysis based on all published high quality data up to and including June 3rd 2021 on the maternal and neonatal outcomes in pregnant women infected with the Coronavirus Disease 2019 (COVID-19).

#### Data Sources

PubMed, SCOPUS, MEDLINE, ClinicalTrials.gov, and Web of Science databases were queried from inception up to June 3rd 2021.

#### Study Eligibility Criteria

We included all clinical studies (prospective and retrospective cohort studies, case-control studies, case series, and rapid communications) that reported data on any maternal and neonatal outcomes of pregnant women with COVID-19.

#### Study Appraisal and Synthesis Methods

Data were analyzed as pooled proportions or odds ratios (OR) and 95% confidence intervals (95% CI) in meta-analysis models.

#### Results

We included 111 studies enrolling 42754 COVID-19-positive pregnant women. From COVID-19-positive pregnant women, the incidence rate of cesarean section was 53.2% (95% CI: 48%–58.4%), 41.5% (95% CI: 36.3%–46.8%) for spontaneous vaginal delivery and 6.4% (95% CI: 4.5%–9.2%) for operative delivery. The rate of some adverse neonatal events was relatively high in mothers infected with COVID-19 including premature delivery (16.7%, 95% CI: 12.8%–21.5%), and low birthweight (16.7%, 95% CI: 12.8%–21.5%). Vertical transmission (3.5%, 95% CI: 2.7%–4.7%), neonatal death (3%, 95% CI: 2%–4%), stillbirth (1.9%, 95% CI: 1.5–2.4%) and maternal mortality (0.012% 95% CI: 0.010–0.014%) were rare adverse events. Mean birth weight was 3069.7g, 95% CI: 3009.7g–3129.8g). In the comparative analysis, COVID-19 significantly increased the risk of premature delivery (OR= 1.48, [95% CI; 1.22, 1.8]), preeclampsia (OR= 1.6, [95% CI; 1.2, 2.1]), stillbirth (OR= 2.36, [95% CI 1.24, 4.462]), neonatal mortality (OR= 3.35, [95% CI; 1.07, 10.5]), and maternal mortality (OR= 3.08, [95% CI; 1.5, 6.3]). Pooled analyses were homogenous, with mild heterogeneity in premature delivery and preeclampsia outcomes.

#### Conclusion

Data must be interpreted with caution as limited data is available and no complete assessment of bias is possible at this time. Our data suggests that pregnant women who test positive for COVID-19 seem to be at higher risk for lower birth weights and premature delivery. There is no evidence at this time of the sharply increased maternal mortality that was seen with both the previous 2002 Middle East Respiratory Syndrome (MERS) and 2003 Severe Acute Respiratory Syndrome (SARS) pandemics.  
(Author)

**Full URL:** <https://doi.org/10.1016/j.xagr.2021.100049>

---

MIDIRS is part of RCM Information Services Limited which is a company incorporated in England and Wales under company no.11914882 with registered office at 10-18 Union Street, London SE1 1SZ  
RCM Information Services Limited is a subsidiary of The Royal College of Midwives



**The Royal College of Midwives Trust**  
10-18 Union Street  
London  
SE1 1SZ



**Open 24 hours a day, 7 days a week**  
T: 0300 303 0444  
F: +44 20 7312 3536  
E: [info@rcm.org.uk](mailto:info@rcm.org.uk)  
W: [www.rcm.org.uk](http://www.rcm.org.uk)



**Chief Executive:**  
Gill Walton,  
MA, PGDip, BSc Hons, RM  
**President:**  
Rebecca Davies, RM  
**Patron:**  
HRH The Princess Royal

## 2021-09872

**Attitudes Toward COVID-19 Illness and COVID-19 Vaccination among Pregnant Women: A Cross-Sectional Multicenter Study during August–December 2020.** Battarbee AN, Stockwell MS, Varner M, et al (2022), American Journal of Perinatology vol 39, no 1, January 2022, pp 75-83

**Objective** The aim of the study was to evaluate pregnant women's attitudes toward COVID-19 illness and vaccination and identify factors associated with vaccine acceptability.

**Study Design** This was a cross-sectional survey among pregnant women enrolled in a prospective COVID-19 cohort study in Salt Lake City, UT, Birmingham, AL, and New York, NY, from August 9 to December 10, 2020. Women were eligible if they were 18 to 50 years old and <28 weeks of gestation. Upon enrollment, women completed surveys regarding concerns about COVID-19 illness and likelihood of getting COVID-19 vaccine if one were available during pregnancy. Vaccine acceptability was defined as a response of “very likely” or “somewhat likely” on a 4-point Likert scale. Factors associated with vaccine acceptability were assessed with multivariable logistic regression.

**Results** Of 939 pregnant women eligible for the main cohort study, 915 (97%) consented to participate. Among these 915 women, 39% self-identified as White, 23% Black, 33% Hispanic, and 4% Other. Sixty-two percent received an influenza vaccine last season. Seventy-two percent worried about getting sick with COVID-19. If they were to get sick, 92% worried about harm to their pregnancy and 80% about harm to themselves. Only 41% reported they would get a vaccine. Of women who were unlikely to get vaccinated, the most frequently cited concern was vaccine safety for their pregnancy (82%). Non-Hispanic Black and Hispanic women had lower odds of accepting a vaccine compared with non-Hispanic White women (adjusted odds ratios [aOR] 0.4, 95% CI 0.2–0.6 for both). Receipt of influenza vaccine during the previous season was associated with higher odds of vaccine acceptability (aOR 2.1, 95% CI 1.5–3.0).

**Conclusion** Although most pregnant women worried about COVID-19 illness, <50% were willing to get vaccinated during pregnancy. Racial and ethnic disparities in plans to accept COVID-19 vaccine highlight the need to prioritize strategies to address perceived barriers among groups at high risk for COVID-19. (Author)

---

## 2021-09843

**The lethal effect of the second wave of COVID-19 on pregnant women: a matter of concern.** Khoiwal K, Agarwal A, Mittal A, et al (2022), International Journal of Gynecology & Obstetrics vol 156, no 1, January 2022, pp 178-181

The lethal second wave of COVID-19 hit pregnant women terribly with regards to disease severity, requirement of invasive ventilation, and maternal mortality. (Author)

---

## 2021-09741

**SARS-CoV-2 and hypertensive disease in pregnancy.** Madden N, Emeruwa UN, Polin M, et al (2022), American Journal of Obstetrics & Gynecology MFM vol 4, no 1, January 2022, 100496

Correspondence piece aiming to evaluate differences in rates of hypertensive disorders of pregnancy (HDP) and hypertensive disease severity in SARS-CoV-2 patients. Results show that those with SARS-CoV-2 had significantly higher rates of HDP, including gestational hypertension and pre-eclampsia. (LDO)

**Full URL:** <https://doi.org/10.1016/j.ajogmf.2021.100496>

---

MIDIRS is part of RCM Information Services Limited which is a company incorporated in England and Wales under company no.11914882 with registered office at 10-18 Union Street, London SE1 1SZ  
RCM Information Services Limited is a subsidiary of The Royal College of Midwives



**The Royal College of Midwives Trust**  
10-18 Union Street  
London  
SE1 1SZ



**Open 24 hours a day, 7 days a week**  
T: 0300 303 0444  
F: +44 20 7312 3536  
E: [info@rcm.org.uk](mailto:info@rcm.org.uk)  
W: [www.rcm.org.uk](http://www.rcm.org.uk)



**Chief Executive:**  
Gill Walton,  
MA, PGDip, BSc Hons, RM  
**President:**  
Rebecca Davies, RM  
**Patron:**  
HRH The Princess Royal

## 2021-09737

**Neighborhood deprivation and preterm delivery during the coronavirus 2019 pandemic.** Fisher SA, Sakowicz A, Barnard C, et al (2022), American Journal of Obstetrics & Gynecology MFM vol 4, no 1, January 2022, 100493

### Background

Prior studies have reported decreases in preterm delivery (PTD) incidence during the coronavirus 2019 (COVID-19) pandemic, however findings are inconsistent. Given wide disparities in the pandemic's impact across communities, neighborhood deprivation may explain observed variation in the relationship between the COVID-19 pandemic and preterm delivery.

### Objective

To characterize changes in the incidence of PTD during the COVID-19 pandemic with attention to effect modification introduced by neighborhood hardship.

### Study Design

This retrospective cohort study included all pregnant patients who delivered at an urban tertiary care hospital during the pandemic (April-November 2020) or pre-pandemic (April-November 2019). We compared the incidence of PTD, spontaneous PTD, and medically indicated PTD prior to 37 weeks' gestation across epochs. Planned analyses stratified the cohorts by neighborhood deprivation metrics defined by residential zip code, including median neighborhood household income and hardship index (a composite index including dependency, educational attainment, unemployment, poverty, per capita income, and crowded housing). The Breslow-Day test for homogeneity assessed the association of delivery epoch and neighborhood deprivation with PTD outcome.

### Results

Of 16,544 eligible deliveries, 8.7% occurred preterm. Incidences of PTD (8.4% vs. 9.0%,  $p=0.17$ ), spontaneous PTD (5.0 vs 5.4%,  $p=0.27$ ), and medically indicated PTD (3.2% v 3.5%,  $p=0.47$ ) were similar between the pandemic and pre-pandemic epochs. However, PTD (OR 0.78, 95% CI 0.64-0.96) and spontaneous PTD (OR 0.76, 95% CI 0.59-0.99) decreased from the pre-pandemic to pandemic epoch in those living in neighborhoods <50th percentile for median income (Breslow-Day  $p$ -values 0.047 and 0.036, respectively). Similarly, PTD (OR 0.78, 95% CI 0.64-0.97) and spontaneous PTD (OR 0.74, 95% CI 0.57-0.98) decreased for those inhabiting neighborhoods in the highest-hardship quartile (Breslow-Day  $p$ -values 0.045 and 0.029, respectively).

### Conclusion

Populations residing in socioeconomically disadvantaged neighborhoods experienced reductions in preterm delivery during the COVID-19 pandemic. Neighborhood-level social determinants of health offer insight into the complex etiologies that contribute to preterm delivery, and provide opportunities for public health, equity-focused prevention strategies. (Author)

Full URL: <https://doi.org/10.1016/j.ajogmf.2021.100493>

## 2021-09403

**Worse Outcomes of Pregnancy in COVID-19 Infection during Parturition may be due to Referral Bias: Analysis in a Prospective Cohort of 963 pregnancies.** Mohini, Ahmed S, Kasarla V, et al (2022), American Journal of Obstetrics & Gynecology (AJOG) vol 226, no 1, January 2022, pp 144-145.e3

Research letter analysing a prospective cohort of pregnancies to determine if high risk of severe COVID-19 in referral centres was confounded due to other risk factors. Results indicate that COVID-19 infection does not pose additional risk to pregnancy outcomes on its own. (LDO)

Full URL: <https://doi.org/10.1016/j.ajog.2021.08.058>

MIDIRS is part of RCM Information Services Limited which is a company incorporated in England and Wales under company no.11914882 with registered office at 10-18 Union Street, London SE1 1SZ  
RCM Information Services Limited is a subsidiary of The Royal College of Midwives



**The Royal College of Midwives Trust**  
10-18 Union Street  
London  
SE1 1SZ



**Open 24 hours a day, 7 days a week**  
T: 0300 303 0444  
F: +44 20 7312 3536  
E: [info@rcm.org.uk](mailto:info@rcm.org.uk)  
W: [www.rcm.org.uk](http://www.rcm.org.uk)



**Chief Executive:**  
Gill Walton,  
MA, PGDip, BSc Hons, RM  
**President:**  
Rebecca Davies, RM  
**Patron:**  
HRH The Princess Royal

**2021-09398**

**Increasing severity of COVID-19 in pregnancy with Delta (B.1.617.2) variant surge.** Adhikari EH, SoRelle JA, McIntire DD, et al (2022), American Journal of Obstetrics & Gynecology (AJOG) vol 226, no 1, January 2022, pp 149-151

Research letter reporting trends in illness severity among obstetric patients with COVID-19 in the context of the Delta variant. Results indicate increasing rates of hospitalisation and morbidity and highlight the need for prevention measures such as vaccination. (LDO)

**Full URL:** <https://doi.org/10.1016/j.ajog.2021.09.008>

---

**2021-08901**

**Coronavirus disease 2019 and pregnancy is déjà vu all over again.** Rasmussen SA, Jamieson DJ (2022), BJOG: An International Journal of Obstetrics and Gynaecology vol 129, no 2, January 2022, pp 188-191

Commentary on public health emergencies in the last two decades including H1N1 influenza, Ebola virus, Zika virus, SARS and COVID-19. Highlights the need for strategies to address future emergencies and pandemics, particularly for vulnerable groups such as pregnant women. (LDO)

**Full URL:** <https://doi.org/10.1111/1471-0528.16859>

---

MIDIRS is part of RCM Information Services Limited which is a company incorporated in England and Wales under company no.11914882 with registered office at 10-18 Union Street, London SE1 1SZ  
RCM Information Services Limited is a subsidiary of The Royal College of Midwives



**The Royal College of Midwives Trust**  
10-18 Union Street  
London  
SE1 1SZ



**Open 24 hours a day, 7 days a week**  
T: 0300 303 0444  
F: +44 20 7312 3536  
E: [info@rcm.org.uk](mailto:info@rcm.org.uk)  
W: [www.rcm.org.uk](http://www.rcm.org.uk)



**Chief Executive:**  
Gill Walton,  
MA, PGDip, BSc Hons, RM  
**President:**  
Rebecca Davies, RM  
**Patron:**  
HRH The Princess Royal



**2021-07624**

**SARS-COV-2 infection during pregnancy and risk of preeclampsia: a systematic review and meta-analysis.** Conde-Agudelo A, Romero R (2022), American Journal of Obstetrics & Gynecology (AJOG) vol 226, no 1, January 2022, pp 68-89.e3

#### OBJECTIVE

To examine the relationship between severe acute respiratory syndrome coronavirus 2 (SARS-CoV-2) infection during pregnancy and the risk of preeclampsia.

#### DATA SOURCES

MEDLINE, EMBASE, POPLINE, CINAHL, LILACS, and WHO COVID-19, Chinese, and preprint databases (all from December 1, 2019 to May 31, 2021). Google Scholar, bibliographies, and conference proceedings were also searched.

#### STUDY ELIGIBILITY CRITERIA

Observational studies that assessed the association between SARS-CoV-2 infection during pregnancy and preeclampsia and that reported unadjusted and/or adjusted risk estimates and 95% confidence intervals (CIs) or data to calculate them.

#### STUDY APPRAISAL AND SYNTHESIS METHODS

The primary outcome was preeclampsia. Secondary outcomes included preeclampsia with severe features, preeclampsia without severe features, eclampsia, and the hemolysis, elevated liver enzymes, and low platelet count (HELLP) syndrome. Two reviewers independently reviewed studies for inclusion, assessed their risk of bias, and extracted data. Pooled unadjusted and adjusted odds ratios (ORs) with 95% CIs, and 95% prediction interval were calculated. Heterogeneity was quantified with the I<sup>2</sup> statistic, where I<sup>2</sup> ≥30% indicated substantial heterogeneity. Subgroup and sensitivity analyses were performed for testing the robustness of the overall findings.

#### RESULTS

Twenty-eight studies that included 790,954 pregnant women, among which 15,524 were diagnosed with SARS-CoV-2 infection, met the inclusion criteria. The meta-analysis of unadjusted ORs showed that the odds of developing preeclampsia were significantly higher among pregnant women with SARS-CoV-2 infection than among those without SARS-CoV-2 infection (7.0% vs 4.8%; pooled OR 1.62, 95% CI 1.45-1.82; P <0.00001; I<sup>2</sup>=17%; 26 studies; 95% prediction interval of the OR, 1.28-2.05). The meta-analysis of adjusted ORs also showed that SARS-CoV-2 infection during pregnancy was associated with a significant increase in the odds of preeclampsia (pooled OR 1.58, 95% CI 1.39-1.80; P <0.0001; I<sup>2</sup>=0%; 11 studies). There was a statistically significant increase in the odds of preeclampsia with severe features (OR 1.76, 95% CI 1.18-2.63; I<sup>2</sup>=58%; 7 studies), eclampsia (OR 1.97, 95% CI 1.01-3.84; I<sup>2</sup>=0%, 3 studies), and HELLP syndrome (OR 2.10, 95% CI 1.48-2.97; 1 study) among pregnant women with SARS-CoV-2 infection, as compared to those without the infection. Overall, the direction and magnitude of the effect of SARS-CoV-2 infection during pregnancy on preeclampsia was consistent across most pre-specified subgroup and sensitivity analyses. Both asymptomatic and symptomatic SARS-CoV-2 infections significantly increased the odds of preeclampsia although it was higher among patients with symptomatic illness (OR 2.11, 95% CI 1.59-2.81) than among those with asymptomatic illness (OR 1.59, 95% CI 1.21-2.10).

#### CONCLUSIONS

SARS-CoV-2 during pregnancy is associated with higher odds of preeclampsia. (Author)

©2022 MIDIRS All Rights Reserved

This Search Pack is distributed for your personal use, please do not illegally distribute this Search Pack either in part or in its entirety. MIDIRS web site is provided for reference information only. MIDIRS is not responsible or liable for any diagnosis made by a user based on the content of the website. Although great care is taken to ensure reference information is both suitable and accurate, MIDIRS is not liable for the contents of any external internet sites referenced, nor does it endorse any commercial product or service mentioned or advised on any of these sites.

MIDIRS is part of RCM Information Services Limited which is a company incorporated in England and Wales under company no.11914882 with registered office at 10-18 Union Street, London SE1 1SZ  
RCM Information Services Limited is a subsidiary of The Royal College of Midwives



**The Royal College of Midwives Trust**  
10-18 Union Street  
London  
SE1 1SZ



**Open 24 hours a day, 7 days a week**  
T: 0300 303 0444  
F: +44 20 7312 3536  
E: info@rcm.org.uk  
W: www.rcm.org.uk



**Chief Executive:**  
Gill Walton,  
MA, PGDip, BSc Hons, RM  
**President:**  
Rebecca Davies, RM  
**Patron:**  
HRH The Princess Royal